

Real hypersurfaces with Reeb invariant shape operator in the complex hyperbolic quadric

By DOO HYUN HWANG (Daegu), JUAN DE DIOS PÉREZ (Granada)
and YOUNG JIN SUH (Daegu)

Abstract. First, we introduce the notion of Reeb invariant shape operator for real hypersurfaces in the complex hyperbolic quadric $Q^{m*} = SO_{2,m}^0/SO_m SO_2$, $m \geq 3$. Next, we give a complete classification of real hypersurfaces with *Reeb invariant shape operator* in the complex hyperbolic quadrics Q^{m*} .

DOO HYUN HWANG
RESEARCH INSTITUTE OF REAL
AND COMPLEX MANIFOLDS (RIRCM)
KYUNGPOOK NATIONAL UNIVERSITY
41566 DAEGU
REPUBLIC OF KOREA

JUAN DE DIOS PÉREZ
DEPARTMENT OF GEOMETRY AND TOPOLOGY
UNIVERSITY OF GRANADA
18071 GRANADA
SPAIN

YOUNG JIN SUH
DEPARTMENT OF MATHEMATICS AND RIRCM
KYUNGPOOK NATIONAL UNIVERSITY
41566 DAEGU
REPUBLIC OF KOREA

Mathematics Subject Classification: 53C40, 53C55.

Key words and phrases: Reeb invariant, real hypersurface, isometric Reeb flow, \mathfrak{A} -isotropic singular, complex hyperbolic quadric.