

The number of prime factors in h -free and h -full polynomials over function fields

By MATILDE LALÍN (Montreal) and ZHEXING ZHANG (Hong Kong)

Abstract. We study the distribution of the number of prime divisors function Ω over polynomials of the function field $\mathbb{F}_q(T)$ when restricted to h -free polynomials and to h -full polynomials. We use an adaptation of the Selberg–Delange method to arithmetical semigroups due to Warlimont to compute the first and second moments in each case and show that a generalization of the Erdős–Kac Theorem is true.

MATILDE LALÍN
DÉPARTEMENT DE MATHÉMATIQUES
ET DE STATISTIQUE
UNIVERSITÉ DE MONTRÉAL
CP 6128, SUCC. CENTRE-VILLE
MONTREAL, QC H3C 3J7
CANADA

ZHEXING ZHANG
DEPARTMENT OF MATHEMATICS
THE UNIVERSITY OF HONG KONG
ROOM 408, RUN RUN SHAW BUILDING
POKFULAM, HONG KONG
CHINA

Mathematics Subject Classification: Primary: 11N56; Secondary: 11N37, 11B99.

Key words and phrases: h -free polynomial, h -full polynomial, Ω over $\mathbb{F}_q[T]$, Erdős–Kac theorem.