

On a class of generalized Berwald manifolds

By AKBAR TAYEBI (Qom) and FAEZEH ESLAMI (Qom)

Abstract. Let $F = \alpha\phi(s)$, $s := \beta/\alpha$, be a generalized Berwald (α, β) -metric on a two-dimensional manifold. Suppose that F has vanishing S -curvature and $\phi'(0) \neq 0$. We show that if F is regular, then it is a locally Minkowskian metric. If F is an almost regular and non-locally Minkowskian metric, then we explicitly determine the function ϕ which results generalized Berwald metrics not belonging to the classes of Berwald, Landsberg or Douglas metrics. Furthermore, we prove that a left-invariant Finsler metric on a two-dimensional Lie group has vanishing S -curvature if and only if it is a Riemannian metric of constant Gaussian curvature. Finally, we construct a family of Randers type generalized Berwald metrics on an arbitrary odd-dimensional manifold.

AKBAR TAYEBI &
FAEZEH ESLAMI
DEPARTMENT OF MATHEMATICS
FACULTY OF SCIENCE
UNIVERSITY OF QOM
QOM
IRAN

Mathematics Subject Classification: 53C60, 53C25.

Key words and phrases: generalized Berwald metric, Berwald metric, (α, β) -metric, S -curvature.