

## Ordered separation axioms and the Wallman ordered compactification

By HANS-PETER A. KÜNZI (Rondebosch), AISLING E. MCCLUSKEY (Galway)  
and THOMAS A. RICHMOND (Bowling Green)

**Abstract.** Two constructions have been given previously of the Wallman ordered compactification  $w_0X$  of a  $T_1$ -ordered, convex ordered topological space  $(X, \tau, \leq)$ . Both of those papers note that  $w_0X$  is  $T_1$ , but need not be  $T_1$ -ordered. Using this as one motivation, we propose a new version of  $T_1$ -ordered, called  $T_1^K$ -ordered, which has the property that the Wallman ordered compactification of a  $T_1^K$ -ordered topological space is  $T_1^K$ -ordered. We also discuss the  $R_0$ -ordered ( $R_0^K$ -ordered) property, defined so that an ordered topological space is  $T_1$ -ordered ( $T_1^K$ -ordered) if and only if it is  $T_0$ -ordered and  $R_0$ -ordered ( $R_0^K$ -ordered).

### 1. Introduction

Given a set  $X$  with a topology  $\tau$  and a partial order  $\leq$ , we recall the topology  $\tau^\sharp$  on  $X$  as the collection of all  $\tau$ -open  $\leq$ -increasing subsets of  $X$  and the topology  $\tau^\flat$  on  $X$  as the corresponding collection of all  $\tau$ -open  $\leq$ -decreasing subsets of  $X$ . Thus, we may consider the topological space  $(X, \tau)$ , the ordered topological space  $(X, \tau, \leq)$ , or the bitopological space  $(X, \tau^\sharp, \tau^\flat)$ .

As the study of ordered topological spaces and bitopological spaces developed, important topological properties, including separation axioms, were defined

---

*Mathematics Subject Classification:* 54F05, 54D10, 54D35, 54A10, 54E55, 18B30, 06F30.

*Key words and phrases:* ordered topological space,  $T_1$ -ordered,  $T_0$ -ordered,  $R_0$ -ordered,  $T_1^K$ -ordered,  $R_0^K$ -ordered, Wallman ordered compactification, ordered reflection.

Hans-Peter Künzi was partially supported by the South African NRF Grant FA2006022300009 during his stay at the National University of Ireland, Galway, in September 2007.

for these new categories. As a general unifying principle, it seems natural that the chain of implications  $(1) \Rightarrow (2) \Rightarrow (3)$  should hold for the statements

- (1)  $(X, \tau^\sharp, \tau^b)$  satisfies the bitopological (or *pairwise*) property  $P$ ,
- (2)  $(X, \tau, \leq)$  satisfies the ordered property  $P$ , and
- (3)  $(X, \tau)$  satisfies the (topological) property  $P$ .

This scheme is borne out in particular by the complete regularity properties when certain other reasonable necessary conditions are assumed. However, as the ordered theory and bitopological theory were often developed independently, there are exceptions and anomalies, which we address here. In particular, the standard definition of the  $T_1$ -ordered property is seen to fall short of expectations in this regard, as well as in regard to its relation to the Wallman ordered compactification. In Section 2, we review the  $T_0$ -ordered and  $T_1$ -ordered properties and introduce the  $R_0$ -ordered property, noting their relation to the corresponding pairwise properties of  $(X, \tau^\sharp, \tau^b)$ . In Section 3, we introduce the  $T_1^K$ -ordered and  $R_0^K$ -ordered properties, and show that the  $T_1^K$ -ordered property does not have the shortcomings of the standard  $T_1$ -ordered property. Continuing in this theme, Section 4 shows that the Wallman ordered compactification behaves more nicely for  $T_1^K$ -ordered spaces than for  $T_1$ -ordered spaces.

Our notation is that of NACHBIN [15]. If  $(X, \tau, \leq)$  is a partially ordered topological space and  $A \subseteq X$ , then the *increasing hull* of  $A$  is  $i(A) = \{x \in X : \exists a \in A \text{ with } a \leq x\}$ . If  $A = i(A)$ , we say  $A$  is an *increasing set*. The *closed increasing hull* of  $A$ , denoted  $I(A)$ , is the smallest closed increasing set containing  $A$ . The decreasing hull  $d(A)$ , closed decreasing hull  $D(A)$ , and decreasing sets are defined dually. If  $A = \{a\}$ , we write  $i(a)$  for  $i(\{a\})$ . Another useful hull operator, used in the construction of the Wallman ordered compactification, is  $C(A) = I(A) \cap D(A)$ . Following [5] and [6],  $C(A)$  is the *c-set hull* of  $A$ , and if  $A = C(A)$  we say  $A$  is a *c-set*. We say  $(X, \tau, \leq)$  is *convex* if  $\tau$  has a subbase of monotone (i.e., increasing or decreasing) open sets. In other terminology, note that  $D(A) = \text{cl}_{\tau^\sharp}(A)$  and  $I(A) = \text{cl}_{\tau^b}(A)$  where  $\tau^\sharp = \{U \in \tau : U = i(U)\}$  and  $\tau^b = \{U \in \tau : U = d(U)\}$ .

The  $T_0$ (-ordered) *reflection* of a (partially ordered) topological space  $X$  is the  $T_0$ (-ordered) quotient space  $Y$  of  $X$  such that for any continuous (and increasing) function  $f$  from  $X$  into any arbitrary  $T_0$ (-ordered) space  $Z$ , there exists a unique continuous (and increasing) function  $h : Y \rightarrow Z$  with  $f = h \circ q$ , where  $q : X \rightarrow Y$  is the quotient map. The construction of the  $T_0$ -ordered reflection, considered in Section 2, utilizes the equivalence relation defined by  $x \approx y$  if and only if  $I(x) = I(y)$  and  $D(x) = D(y)$ , or equivalently, if and only if  $C(x) = C(y)$  (see [8]). Let  $[x]$  denote the  $\approx$ -equivalence class of  $x$ . The closure operator  $C(\cdot)$

defines a reflexive, transitive relation  $C = \bigcup_{x \in X} \{x\} \times C(x)$ . The inverse relation  $C^{-1}$  is given by  $C^{-1}(x) = \bigcap \{U \in \tau^\# : x \in U\} \cap \bigcap \{V \in \tau^b : x \in V\}$ . (Some authors would call this the intersection of the  $\tau^\#$ -kernel of  $\{x\}$  and the  $\tau^b$ -kernel of  $\{x\}$ .) If  $C^s = C \cap C^{-1}$  is the symmetrization of  $C$ , then  $C^s(x) = [x]$ . Note that if  $X$  is convex, we have  $x \approx y$  if and only if  $\text{cl}\{x\} = \text{cl}\{y\}$ . Also if  $X$  is convex,  $[x] = \text{cl}\{x\} \cap \bigcap \{U \in \tau : x \in U\} \subseteq \text{cl}\{x\} = C(x)$ .

## 2. $T_0$ , $T_1$ , and $R_0$ separation properties

### $T_0$ Properties

We recall the following definitions (see [17]).

*Definition 1.* Suppose  $X$  is a set,  $\tau$  is a topology on  $X$ , and  $\leq$  is a partial order on  $X$ .

- (a)  $(X, \tau)$  is  $T_0$  if  $\text{cl}(x) = \text{cl}(y)$  implies  $x = y$ .
- (b)  $(X, \tau, \leq)$  is  $T_0$ -ordered if for any two distinct points, there exists a monotone open neighborhood of one of the points which does not contain the other, or equivalently, if  $C^s(x) = \{x\}$  for all  $x \in X$ .
- (c)  $(X, \tau_1, \tau_2)$  is weak pairwise  $T_0$  if for distinct points  $x$  and  $y$ , one of the points is not in one of the closures ( $\tau_1$  or  $\tau_2$ ) of the other.
- (d)  $(X, \tau_1, \tau_2)$  is pairwise  $T_0$  if for distinct points  $x$  and  $y$ , either  $x \notin \text{cl}_{\tau_1}(y)$  or  $y \notin \text{cl}_{\tau_2}(x)$ .

Note that pairwise  $T_0$  is defined by  $x \neq y \Rightarrow ([x \notin \text{cl}_{\tau_1}(y) \vee y \notin \text{cl}_{\tau_2}(x)] \wedge [y \notin \text{cl}_{\tau_1}(x) \vee x \notin \text{cl}_{\tau_2}(y)])$ , while weak pairwise  $T_0$  replaces the  $\wedge$  between the bracketed items by  $\vee$ . Clearly pairwise  $T_0$  implies weak pairwise  $T_0$  and, from the open set characterization, it is easily seen that  $(X, \tau, \leq)$  being  $T_0$ -ordered implies  $(X, \tau)$  is  $T_0$ . Now  $(X, \tau^\#, \tau^b)$  is weak pairwise  $T_0$  if and only if  $x \neq y$  implies  $[x \notin D(y) \text{ or } x \notin I(y) \text{ or } y \notin D(x) \text{ or } y \notin I(x)]$ . Since the bracketed condition is equivalent to  $C(x) \neq C(y)$ , we see that  $(X, \tau^\#, \tau^b)$  is weakly pairwise  $T_0$  if and only if  $(X, \tau, \leq)$  is  $T_0$ -ordered. Furthermore, if  $(X, \tau, \leq)$  is convex, these two properties are also equivalent to  $(X, \tau)$  being  $T_0$ .

### $T_1$ Properties

The standard definition of a  $T_1$ -ordered space, given by NACHBIN [15], does not meet many of the expectations one would have for  $T_1$ -ordered spaces. For example, if  $(X, \tau^\#, \tau^b)$  is pairwise  $T_1$ , it does not follow that  $(X, \tau, \leq)$  is  $T_1$ -ordered. Also, the Wallman ordered compactification of a  $T_1$ -ordered topological space is

$T_1$ , but need not be  $T_1$ -ordered. These deficiencies in the standard definition of  $T_1$ -ordered led us to define a new version in the next section. Here we present the standard definitions.

*Definition 2.* Suppose  $X$  is a set,  $\tau$  is a topology on  $X$ , and  $\leq$  is a partial order on  $X$ .

- (a)  $(X, \tau)$  is  $T_1$  if  $\text{cl}(x) = \{x\}$  for all  $x \in X$ .
- (b)  $(X, \tau, \leq)$  is  $T_1$ -ordered if  $d(x)$  and  $i(x)$  are closed for all  $x \in X$ .
- (c)  $(X, \tau_1, \tau_2)$  is pairwise  $T_1$  (see [14]) if for any two distinct points  $x$  and  $y$  in  $X$ , each has either a  $\tau_1$ -open or a  $\tau_2$ -open neighborhood which excludes the other.

Our definition of pairwise  $T_1$ , given by MURDESHWAR and NAIMPALLY [14] is called MN-pairwise- $T_1$  by REILLY [17] and MRŠEVIĆ [12]. Several other versions of pairwise  $T_1$  have been studied.

- (1) (REILLY [17])  $(X, \tau_1, \tau_2)$  is *Reilly pairwise  $T_1$*  if for distinct points  $x$  and  $y$  in  $X$ , there exists a  $\tau_1$ -open neighborhood of  $x$  excluding  $y$  and a  $\tau_2$ -open neighborhood of  $y$  excluding  $x$ .
- (2) (MRŠEVIĆ [12])  $(X, \tau_1, \tau_2)$  is *middle pairwise  $T_1$*  if for distinct points  $x$  and  $y$  in  $X$ ,  $\text{cl}_{\tau_1}(x) \cap \text{cl}_{\tau_2}(y) = \emptyset$  or  $\text{cl}_{\tau_1}(y) \cap \text{cl}_{\tau_2}(x) = \emptyset$ .
- (3) (SWART [19])  $(X, \tau_1, \tau_2)$  is *weak pairwise  $T_1$*  if for distinct points  $x$  and  $y$  in  $X$ , there exists a  $\tau_1$ -open neighborhood of  $x$  excluding  $y$  and a  $\tau_2$ -open neighborhood of  $y$  excluding  $x$ , or there exists a  $\tau_1$ -open neighborhood of  $y$  excluding  $x$  and a  $\tau_2$ -open neighborhood of  $x$  excluding  $y$ .

We note that, given an ordered topological space  $(X, \tau, \leq)$ , the corresponding bitopological space  $(X, \tau^\#, \tau^\flat)$  is Reilly pairwise  $T_1$  only if the order  $\leq$  is discrete (that is, equality), rendering this version not useful in the study of ordered topological spaces. The version due to Mršević is called “middle” since it is implied by Reilly pairwise  $T_1$  and implies Swart’s weak pairwise  $T_1$ . All of these versions of pairwise  $T_1$  are stronger than our definition of (MN) pairwise  $T_1$ . Some elementary facts regarding these definitions are found in [4] and [18]. The fact that many versions of the pairwise  $T_1$  property have been considered suggests that there may be more than one useful version of the  $T_1$ -ordered property.

The new version of  $T_1$ -ordered introduced below, called  $T_1^K$ -ordered, is implied by  $(X, \tau^\#, \tau^\flat)$  being (MN) pairwise  $T_1$ , the weakest of these pairwise  $T_1$  properties.

It is easily seen that  $(X, \tau, \leq)$  being  $T_1$ -ordered implies  $(X, \tau)$  is  $T_1$ , and  $(X, \tau^\sharp, \tau^\flat)$  being pairwise  $T_1$  implies  $(X, \tau)$  is  $T_1$ . However,  $(X, \tau^\sharp, \tau^\flat)$  being pairwise  $T_1$  does not imply  $(X, \tau, \leq)$  is  $T_1$ -ordered. Indeed, the example after Theorem 15 shows that  $(X, \tau^\sharp, \tau^\flat)$  being middle pairwise  $T_1$  (the strongest of these pairwise  $T_1$  properties which can possibly be satisfied by ordered spaces other than antichains) does not imply  $(X, \tau, \leq)$  is  $T_1$ -ordered. That example also shows that  $(X, \tau^\sharp, \tau^\flat)$  being middle pairwise  $T_1$  does not imply that  $(X, \tau^\sharp, \tau^\flat)$  is pairwise  $R_0$  (for the definition of this concept, see below).

The following characterization of the  $T_1$ -ordered property will be used often.

**Lemma 3.**  $(X, \tau, \leq)$  is  $T_1$ -ordered if and only if  $i(x) = \bigcap \{U \in \tau^\sharp : x \in U\}$  and dually for all  $x \in X$ .

PROOF. If  $X$  is  $T_1$ -ordered, then  $y \notin i(x)$  implies  $x \notin d(y) = D(y)$ , so  $i(x) \subseteq X \setminus D(y) \equiv H_y$ , and thus  $i(x) = \bigcap \{H_y : y \notin i(x)\} \supseteq \bigcap \{U \in \tau^\sharp : x \in U\}$ . The reverse inclusion is immediate. Conversely, if  $y \notin i(x)$ , then  $x \notin d(y) = \bigcap \{U \in \tau^\flat : y \in U\}$ , so there exists an open decreasing neighborhood of  $y$  disjoint from  $x$  and therefore from  $i(x)$ , so  $y \notin \text{cl}(i(x))$ , and thus  $i(x)$  is closed.  $\square$

**$R_0$  Properties**

One motivation for  $R_0$  spaces, introduced by DAVIS [3], is that a space is  $T_1$  if and only if it is  $T_0$  and  $R_0$ . The analogous equivalences in the categories of ordered topological spaces and bitopological spaces are used to motivate definitions of the appropriate concepts of  $R_0$  in those categories. Note however, as seen in the example mentioned before Lemma 3, that the weak concept of pairwise  $T_1$  used in this article does not imply pairwise  $R_0$  as defined below. It appears that  $R_0$ -ordered spaces have not been studied previously, so our discussion here will be more thorough. We start with some theorems providing equivalent definitions of the  $R_0$  properties in the three categories in question.

**Theorem 4.** *The following are equivalent:*

- (a)  $(X, \tau)$  is an  $R_0$ -space.
- (b)  $F$  closed,  $x \notin F \Rightarrow \exists$  open  $U$  with  $F \subseteq U, x \notin U$ .
- (c)  $U$  open,  $x \in U \Rightarrow \text{cl}\{x\} \subseteq U$ .
- (d)  $\{\text{cl}\{x\} : x \in X\}$  is a partition of  $X$ .
- (e)  $\tau$  is lattice isomorphic to the topology of a  $T_1$ -space.
- (f)  $\text{cl}\{x\} \neq \text{cl}\{y\} \Rightarrow \exists$  neighborhood of  $x$  not containing  $y$ .
- (g)  $F$  closed,  $\text{cl}\{x\} \cap F \neq \emptyset \Rightarrow x \in F$ .
- (h)  $\bigcap \{U \in \tau : x \in U\} = \text{cl}\{x\}$  for all  $x \in X$ .

- (i) *The  $T_0$ -reflection of  $X$  is  $T_1$ .*
- (j)  $\bigcap\{U \in \tau : A \subseteq U\} \subseteq \text{cl } A$  for all  $A \subseteq X$ .

These equivalences may be found in [3] or [1], except for (j), which is easily shown. (In [1], a  $T_{(\alpha,\beta)}$  space is one whose  $T_\alpha$  reflection is already  $T_\beta$ , so there,  $R_0$  spaces are called  $T_{(0,1)}$  spaces.) Note that the containment in (j) cannot be strengthened to equality: Let  $X = [0, 1]$  with the discrete topology on  $(0, 1]$  and the usual neighborhoods of 0. Now (b) holds, but  $A = (0, 1] = \bigcap\{U \in \tau : A \subseteq U\} \neq \text{cl } A = [0, 1]$ .

It will be interesting to note that many of the characterizations of  $R_0$  given above have direct analogs in either the ordered topological setting or the bitopological setting, but not both.

The definition given below for an  $R_0$ -ordered space arises from the necessary and sufficient conditions in [8] for the  $T_0$ -ordered reflection of an ordered topological space to be  $T_1$ -ordered. For an ordered topological space  $(X, \tau, \leq)$ , we obtain the  $T_0$ -ordered reflection  $(X/\approx, \tau^0, \leq^0)$  as an ordered quotient of  $X$ , modulo the equivalence relation  $x \approx y$  if and only if  $C(x) = C(y)$ . The order  $\leq^0$  on  $X/\approx$  is the “finite step order” given by

$$[z_0] \leq^0 [z_n] \iff \exists [z_1], [z_2], \dots, [z_{n-1}] \text{ and } \exists z'_i, z^*_i \in [z_i] \ (i = 0, 1, \dots, n) \\ \text{with } z'_i \leq z^*_{i+1} \ \forall i = 0, 1, \dots, n - 1.$$

Note that any closed or open monotone set  $S$  in  $X$  is  $\approx$ -saturated (that is,  $x \in S$  implies  $[x] \subseteq S$ ).

We note that MRŠEVIĆ [13] introduced a bitopological “quotient space” which, in the case of  $(X, \tau^\sharp, \tau^\flat)$ , is equivalent to the  $T_0$ -ordered reflection of  $(X, \tau, \leq)$ .

**Lemma 5.** *Suppose  $x, y \in X$ ,  $F \subseteq X$ , and  $f : X \rightarrow X/\approx$  is the quotient map from an ordered topological space  $X$  to its  $T_0$ -ordered reflection  $X/\approx$ .*

- (a) *If  $A$  is closed and increasing in  $X$  then  $f(A)$  is closed and increasing in  $X/\approx$ , and dually. If  $A$  is  $\approx$ -saturated, the converse holds.*
- (b)  *$B$  is closed and increasing in  $X/\approx$  if and only if  $f^{-1}(B)$  is closed and increasing in  $X$ , and dually.*
- (c)  *$f(I(x)) = I_{X/\approx}([x])$  and  $f(D(x)) = D_{X/\approx}([x])$ .*
- (d)  *$f^{-1}(I_{X/\approx}(f(x))) = f^{-1}(I_{X/\approx}([x])) = I(x)$  and dually.*
- (e) *If  $[y] \in f(D(F))$ , and  $[x] \leq [y]$  in  $X/\approx$ , then  $x \in D(F)$ , and dually.*

PROOF. (a), (b), and (c) were stated and justified in the paragraph before Theorem 3.1 of [8]. See also Corollary 2 and Proposition 5 of [13].

(d)  $I(x) \subseteq f^{-1}(I_{X/\approx}([x]))$  since the latter set is a closed increasing set containing  $x$  and the former is the smallest such set. For the reverse inclusion, if  $z \in f^{-1}(I_{X/\approx}([x]))$  but  $z \notin I(x)$ , then since  $I(x)$  is saturated, applying  $f$  (and part (c)) gives  $f(z) \notin f(I(x)) = I_{X/\approx}([x])$ , contrary to  $z \in f^{-1}(I_{X/\approx}([x]))$ .

(e) Suppose  $[y] \in f(D(F))$  with  $[x] \leq [y]$  in  $X/\approx$ . Now  $D(F)$  is closed and decreasing, and thus saturated, and contains  $y$ . Since  $X/\approx$  carries the finite step order, it follows that  $x \in D(F)$ .  $\square$

Any of the equivalent statements below may be taken as the definition of an  $R_0$ -ordered space.

**Theorem 6.** *For an ordered topological space  $(X, \tau, \leq)$  with  $T_0$ -ordered reflection  $X/\approx$ , the following are equivalent:*

- (a)  $(X, \tau, \leq)$  is  $R_0$ -ordered.
- (b)  $\bigcap \{U \in \tau^\# : x \in U\} = i([x])$  and dually for all  $x \in X$ .
- (c) The  $T_0$ -ordered reflection of  $X$  is  $T_1$ -ordered.
- (d)  $[x] \not\leq [y]$  in  $X/\approx$  implies  $x \notin D(y)$  and  $y \notin I(x)$ .
- (e)  $I(x) = f^{-1}(i_{X/\approx}([x]))$  and  $D(x) = f^{-1}(d_{X/\approx}([x]))$  for all  $x \in X$ , where  $f : X \rightarrow X/\approx$  is the natural ordered quotient map.

PROOF. The equivalence of (b) and (c) is Theorem 3.2 of [8], and was the impetus for taking these conditions to be the definition of  $R_0$ -ordered.

(c)  $\Rightarrow$  (d). The proof of this implication is modeled on the proof of the corresponding non-ordered case (Theorem 3.5 (i)  $\Rightarrow$  (ii) in [1]). Suppose (c) holds and  $[x] \not\leq [y]$  in  $X/\approx$ . Then  $[y] \notin i_{X/\approx}([x]) = I_{X/\approx}([x])$ . Applying  $f^{-1}$ , where  $f : X \rightarrow X/\approx$  is the (ordered) quotient map, we have  $y \notin f^{-1}(I_{X/\approx}([x])) = f^{-1}(f(I(x))) \supseteq I(x)$ , so  $y \notin I(x)$ . Similarly,  $[x] \notin d_{X/\approx}([y])$  implies  $x \notin D(y)$ .

(d)  $\Rightarrow$  (c). Suppose (d) holds and  $[y] \notin i_{X/\approx}([x])$ . Then  $x \notin D(y)$  implies that  $X \setminus D(y)$  is an open increasing (and thus, saturated) neighborhood of  $x$  which does not include  $y$ , so  $[y] \notin I_{X/\approx}([x])$ . Similarly,  $y \notin I(x)$  shows  $X \setminus I(x)$  is an open decreasing saturated neighborhood of  $y$  not containing  $x$ .

(c)  $\Rightarrow$  (e). Suppose (c) holds. Then  $i_{X/\approx}([x])$  is closed and increasing, so  $f^{-1}(i_{X/\approx}([x]))$  is closed, increasing, and contains  $x$ , so  $I(x) \subseteq f^{-1}(i_{X/\approx}([x]))$ . For the reverse inclusion, if  $z \in f^{-1}(i_{X/\approx}([x]))$ , then  $[z] = f(z) \in i_{X/\approx}([x]) = I_{X/\approx}([x]) = f(I(x))$ , and thus  $z \in I(x)$  by Lemma 5 (e). Thus, we have  $I(x) = f^{-1}(i_{X/\approx}([x]))$ . The dual argument completes this implication.

(e)  $\Rightarrow$  (c). Suppose (e). Then by Lemma 5 (d), we have  $f^{-1}(I_{X/\approx}([x])) = f^{-1}(i_{X/\approx}([x]))$  and  $f^{-1}(D_{X/\approx}([x])) = f^{-1}(d_{X/\approx}([x]))$  for all  $x \in X$ , and applying  $f$ , which is onto, shows  $I_{X/\approx}([x]) = i_{X/\approx}([x])$  and  $D_{X/\approx}([x]) = d_{X/\approx}([x])$  for all  $[x] \in X/\approx$ , so  $X/\approx$  is  $T_1$ -ordered.  $\square$

We note that items (b), (c), and (d) of Theorem 6 are direct analogs of items (h), (i), and (f), respectively, of Theorem 4.

**Theorem 7.** *An ordered topological space  $(X, \tau, \leq)$  is  $T_1$ -ordered if and only if it is  $T_0$ -ordered and  $R_0$ -ordered.*

PROOF. Suppose  $(X, \tau, \leq)$  is  $T_0$ -ordered and  $R_0$ -ordered. The former condition implies  $[x] = \{x\}$  and the latter condition then implies  $i(x) = \bigcap\{U \in \tau^\# : x \in U\}$  and dually, so  $(X, \tau, \leq)$  is  $T_1$ -ordered by Lemma 3.

Conversely, suppose  $(X, \tau, \leq)$  is  $T_1$ -ordered. Then  $[x] = C^s(x) = D(x) \cap I(x) \cap D^{-1}(x) \cap I^{-1}(x) = d(x) \cap i(x) \cap d^{-1}(x) \cap i^{-1}(x) = \{x\}$ , so  $X$  is  $T_0$ -ordered. Thus,  $i([x]) = i(C^s(x)) = i(x) = \bigcap\{U \in \tau^\# : x \in U\}$ , by Lemma 3. With the dual argument, Theorem 6 (b) shows that  $X$  is  $R_0$ -ordered.  $\square$

MISRA and DUBE [11] give several characterizations of the pairwise  $R_0$  property, including the following. The equivalence of (a) and (b) is taken to be the definition of the pairwise  $R_0$  property.

**Theorem 8** ([11]). *For a bitopological space  $(X, \tau_1, \tau_2)$ , the following are equivalent:*

- (a)  $(X, \tau_1, \tau_2)$  is pairwise  $R_0$ .
- (b) For every  $x \in X$ , if  $U$  is a  $\tau_i$ -open neighborhood of  $x$ , then  $\text{cl}_{\tau_j}(x) \subseteq U$ , where  $\{i, j\} = \{1, 2\}$ .
- (c)  $y \in \text{cl}_{\tau_i}(x) \iff x \in \text{cl}_{\tau_j}(y)$  for  $\{i, j\} = \{1, 2\}$ .
- (d) If  $F$  is  $\tau_i$ -closed and  $x \notin F$ , then there exists a  $\tau_j$ -open set  $U$  with  $F \subseteq U$  and  $x \notin U$ , for  $\{i, j\} = \{1, 2\}$ .
- (e)  $\bigcap\{U \in \tau_i : F \subseteq U\} = F$  for any  $\tau_j$ -closed  $F \subseteq X$ , for  $\{i, j\} = \{1, 2\}$ .
- (f)  $\text{cl}_{\tau_i}(x) \cap \text{cl}_{\tau_j}(F) \neq \emptyset \Rightarrow x \in \text{cl}_{\tau_j}(F)$  for all  $x \in X$ ,  $F \subseteq X$ , and for  $\{i, j\} = \{1, 2\}$ .

We observe that items (b), (d), and (f) of Theorem 8 are pairwise versions of items (c), (b), and (g), respectively, of Theorem 4. Of the standard  $R_0$  defining items from Theorem 4, we found none which had direct analogs to both the ordered setting and the bitopological setting. This fact has also prompted our



reconsideration of the standard definition of  $T_1$ -ordered, and subsequently, of  $R_0$ -ordered. We note that  $(X, \tau_1, \tau_2)$  being pairwise  $R_0$  implies  $(X, \tau_1 \vee \tau_2)$  is  $R_0$  (Proposition 9 of [13]).

### 3. $T_1^K$ -ordered and $R_0^K$ -ordered spaces

Since the usual definitions of  $T_1$ -ordered, and hence  $R_0$ -ordered have some deficiencies as noted above, we propose the following new versions, denoted  $T_1^K$ -ordered and  $R_0^K$ -ordered.

*Definition 9.* An ordered topological space  $(X, \tau, \leq)$  is  $T_1^K$ -ordered if  $C(x) = \{x\}$  for all  $x \in X$ .

We make the following observations.

**Theorem 10.** Suppose  $(X, \tau, \leq)$  is an ordered topological space.

- (a)  $(X, \tau, \leq)$  is  $T_1^K$ -ordered if and only if  $\{x\} = \bigcap \{U \in \tau^\# \cup \tau^b : x \in U\}$  for every  $x \in X$ .
- (b)  $(X, \tau^\#, \tau^b)$  is pairwise  $T_1 \iff (X, \tau, \leq)$  is  $T_1^K$ -ordered  $\Rightarrow (X, \tau)$  is  $T_1$ , and  $(X, \tau, \leq)$  is  $T_1$ -ordered  $\Rightarrow (X, \tau, \leq)$  is  $T_1^K$ -ordered.  $(X, \tau, \leq)$  is  $T_1^K$ -ordered and convex  $\iff (X, \tau)$  is  $T_1$  and  $(X, \tau, \leq)$  is convex.
- (c) For a linearly ordered space  $(X, \tau, \leq)$ ,  $T_1$ -ordered and  $T_1^K$ -ordered are equivalent.

PROOF. (a) is proved analogously to the corresponding characterization of  $T_1$ -ordered given in Lemma 3. The proof of (b) follows immediately from (a), Definitions 2 and 9, and the last sentence of Section 1. For (c), if  $X$  is linearly ordered and  $T_1^K$ -ordered and  $y \notin i(x)$ , then  $y \in d(x) \subseteq D(x)$ , but  $y \notin \{x\} = C(x) = I(x) \cap D(x)$ , so  $y \notin I(x)$ . Thus,  $i(x) = I(x)$ , and with the dual argument,  $X$  is  $T_1$ -ordered. □

The example below shows that  $T_1^K$ -ordered is not equivalent to the middle- or weak- pairwise  $T_1$  properties applied to  $(X, \tau^\#, \tau^b)$ .

*Example 11.* Let  $X = \mathbb{N} \cup \{e, \pi\}$ , where  $\mathbb{N}$  is the set of natural numbers. Take all points in  $\mathbb{N}$  to be topologically isolated and take the neighborhoods of  $x \in \{e, \pi\}$  to be cofinite sets in  $X$  containing  $x$ . Now  $X$  is a  $T_1$  space. Let  $\leq = \Delta \cup \{(2n, e) : n \in \mathbb{N}\} \cup \{(2n - 1, \pi) : n \in \mathbb{N}\}$ . This is a partial order. Note that  $e$  and  $\pi$  are the only nonisolated points in  $X$  and for each  $x \in X$ ,  $d(x)$  and  $i(x)$  are finite and thus closed, except for  $d(\pi)$  and  $d(e)$ . Observe that  $\pi \in D(e)$

and  $e \in D(\pi)$ . It immediately follows that  $C(x) = \{x\}$  for every  $x \in X$ . Thus,  $X$  is a  $T_1^K$ -ordered space.

Consider  $e$  and  $\pi$ . There exists no increasing (open) neighborhood of  $e$  that does not contain  $\pi$  and no increasing (open) neighborhood of  $\pi$  that does not contain  $e$ . Thus, the associated bispace  $(X, \tau^\sharp, \tau^\flat)$  is not a weak pairwise  $T_1$  space, and hence is not middle pairwise  $T_1$ . Noting that  $\pi \in \bigcap \{U \in \tau^\sharp : e \in U\} \neq i(e) = \{e\}$ , Lemma 3 shows that  $X$  is not  $T_1$ -ordered. This is also easily seen since  $d(\pi)$  is not closed.

Now we define  $R_0^K$ -ordered spaces and subsequently list some of their properties.

*Definition 12.* An ordered topological space  $(X, \tau, \leq)$  is  $R_0^K$ -ordered if, for all  $x, y \in X$ , we have  $y \in C(x)$  implies  $x \in C(y)$ .

**Theorem 13.** Suppose  $(X, \tau, \leq)$  is an ordered topological space.

- (a) The following are equivalent:
  - (i)  $(X, \tau, \leq)$  is  $R_0^K$ -ordered.
  - (ii)  $C = C^{-1}$ .
  - (iii)  $C$  is an equivalence relation.
  - (iv)  $x \in C(y)$  if and only if  $C(x) = C(y)$  for all  $x, y \in X$ .
  - (v)  $x \in C(y)$  if and only if  $[x] = [y]$  for all  $x, y \in X$ .
  - (vi)  $\{C(x) : x \in X\}$  is a partition of  $X$ .
- (b)  $(X, \tau^\sharp, \tau^\flat)$  is pairwise  $R_0 \Rightarrow (X, \tau, \leq)$  is  $R_0^K$ -ordered.  $(X, \tau, \leq)$  is  $R_0$ -ordered  $\Rightarrow (X, \tau, \leq)$  is  $R_0^K$ -ordered.  $(X, \tau, \leq)$  is  $R_0^K$ -ordered and convex  $\iff (X, \tau)$  is  $R_0$  and  $(X, \tau, \leq)$  is convex.
- (c) If  $(X, \tau, \leq)$  is a linearly ordered space, then  $R_0$ -ordered and  $R_0^K$ -ordered are equivalent properties.

PROOF. (a) is immediate.

(b) The first implication follows from a comparison of part (iv) of (a) above and Theorem 8 (c).  $X$  is  $R_0$ -ordered if and only if  $X/\approx$  is  $T_1$ -ordered, which implies  $X/\approx$  is  $T_1^K$ -ordered, and hence (by Theorem 15 below)  $X$  is  $R_0^K$ -ordered.

Suppose  $X$  is  $R_0^K$ -ordered and convex. We will show that  $\{\text{cl}(x) : x \in X\}$  partitions  $X$ . Suppose  $z \in \text{cl}(x) \cap \text{cl}(y)$ . Now  $z \in \text{cl}(x) \subseteq C(x)$  implies, by part (v) of (a), that  $[x] = [z]$ , and similarly  $[y] = [z] = [x]$ . Recalling that convexity implies  $[w] \subseteq \text{cl}(w)$  for any  $w \in X$ , we have  $x \in [x] = [y] \subseteq \text{cl}(y)$  and  $y \in [y] = [x] \subseteq \text{cl}(x)$ . Applying the closure operator now shows  $\text{cl}(x) \subseteq \text{cl}(y) \subseteq \text{cl}(x)$ , so  $\text{cl}(x) = \text{cl}(y)$ , and thus  $(X, \tau)$  is  $R_0$  by Theorem 4 (d). Conversely, suppose  $(X, \tau, \leq)$  is convex

and  $(X, \tau)$  is  $R_0$ . We will show that  $z \in C(x)$  implies  $C(z) = C(x)$ , from which it easily follows that  $\{C(x) : x \in X\}$  partitions  $X$  and thus  $X$  is  $R_0^K$ -ordered. If  $z \in C(x)$  and  $x \in C(z)$ , applying the closure operator  $C$  shows  $C(x) = C(z)$ . Thus, suppose  $z \in C(x)$  and  $x \notin C(z)$ . Then  $x \notin \text{cl}(z)$ , so  $\text{cl}(x) \neq \text{cl}(z)$ . Theorem 4 (f) implies the existence of a neighborhood  $N$  of  $z$  which does not contain  $x$ , and by convexity, we may assume  $N$  is monotone open. If  $N$  is increasing, then  $X \setminus N$  is a closed decreasing set containing  $x$  and excluding  $z$ . This shows  $z \notin D(x)$ , giving the contradiction that  $z \notin C(x)$ . The dual argument applies if  $N$  is decreasing.

(c) If  $X$  is a linearly ordered  $R_0^K$ -ordered space, by Theorem 15 below,  $X/\approx$  is  $T_1^K$ -ordered. It is easy to see that the finite step order on  $X/\approx$  is also linear, in which case  $X/\approx$  is  $T_1$ -ordered, and therefore  $X$  is  $R_0$ -ordered. The converse follows from part (b).  $\square$

Observing the appearance of convexity in (b) above, we note that  $R_0^K$ -ordered need not imply  $R_0$  if the topology is not convex. For example, consider  $X = \{\perp, a, b, \top\}$  where  $a$  and  $b$  are noncomparable and  $\perp \leq x \leq \top$  for all  $x \in X$ . Give  $X$  the topology having  $\{\{\top, \perp\}, \{a, b\}, \{a\}\}$  as base of closed sets. It is easy to check that  $C(x) = X$  for each  $x \in X$ , so  $X$  is  $R_0^K$ -ordered. In fact,  $X$  is  $R_0$ -ordered since  $i([x]) = i(X) = X = \bigcap\{U \in \tau^\# : x \in U\}$  and dually for each  $x \in X$ . However,  $\{\text{cl}(x) : x \in X\}$  does not partition  $X$ , so  $X$  is not  $R_0$ .

The next theorems show that the  $R_0^K$ -ordered and  $T_0^K$ -ordered properties interact as one would hope.

**Theorem 14.**  $(X, \tau, \leq)$  is  $T_1^K$ -ordered if and only if it is  $T_0$ -ordered and  $R_0^K$ -ordered.

PROOF. If  $X$  is  $T_0$ -ordered and  $R_0^K$ -ordered, then for all  $x \in X$  we have  $\{x\} = C^{-1}(x) \cap C(x) = C(x) \cap C(x) = C(x)$ , where the first equality follows from the  $T_0$ -ordered property and the second equality from the  $R_0^K$ -ordered property. Thus,  $X$  is  $T_1^K$ -ordered. Conversely, if  $X$  is  $T_1^K$ -ordered, then  $C(x) = \{x\}$  implies  $C(x) \cap C^{-1}(x) = \{x\}$ , so that  $X$  is  $T_0$ -ordered. Also,  $y \in C(x) = \{x\}$  implies  $x \in C(y)$ , so  $X$  is  $R_0^K$ -ordered.  $\square$

**Theorem 15.** The  $T_0$ -ordered reflection  $X/\approx$  of an ordered topological space  $X$  is  $T_1^K$ -ordered if and only if  $X$  is  $R_0^K$ -ordered.

PROOF. If  $X/\approx$  is  $T_1^K$ -ordered, then  $I_{X/\approx}([x]) \cap D_{X/\approx}([x]) = \{[x]\}$  for all  $[x] \in X/\approx$ . Applying  $f^{-1}$  as in Lemma 5 (d) gives  $C(x) = I(x) \cap D(x) = [x]$ . Now  $y \in C(x) = [x] \iff C(x) = C(y)$ , so  $X$  is  $R_0^K$ -ordered by Theorem 13 (a).

Conversely, suppose  $X$  is  $R_0^K$ -ordered and  $[y] \in C_{X/\approx}([x]) = I_{X/\approx}([x]) \cap D_{X/\approx}([x])$ . Applying  $f^{-1}$  as in Lemma 5 (d) gives  $[y] \subseteq I(x) \cap D(x) = C(x)$ ,

and now  $y \in C(x)$  implies  $[y] = [x]$ . It follows that  $C_{X/\approx}([x]) = \{[x]\}$  for any  $[x] \in X/\approx$ , so  $X/\approx$  is  $T_1^K$ -ordered. This direction of the proof also follows from the bitopological quotient construction of Theorem 3.1 of [20], which also appears as Corollary 6 of [13].  $\square$

| $(X, \tau^\sharp, \tau^\flat)$  |                   | $(X, \tau, \leq)$            |                   | $(X, \tau)$  |
|---|-------------------|------------------------------|-------------------|--|
| pairwise $T_0$  | $\Rightarrow$     | $T_0$ -ordered               | $\Rightarrow$     | $T_0$  |
| weak pairwise $T_0$   | $\Leftrightarrow$ | $T_0$ -ordered               | $\Rightarrow$     | $T_0$  |
|   |                   | $T_0$ -ordered + convex      | $\Leftrightarrow$ | $\left\{ \begin{array}{l} T_0 + \\ (X, \tau, \leq) \text{ convex} \end{array} \right.$ |
| pairwise $R_0$  | $\not\Rightarrow$ | $R_0$ -ordered               | $\not\Rightarrow$ | $R_0$  |
|   |                   | $\Downarrow$                 |                   |  |
| pairwise $R_0$  | $\Rightarrow$     | $R_0^K$ -ordered             | $\not\Rightarrow$ | $R_0$  |
|   |                   | $R_0^K$ -ordered + convex    | $\Leftrightarrow$ | $\left\{ \begin{array}{l} R_0 + \\ (X, \tau, \leq) \text{ convex} \end{array} \right.$ |
| pairwise $T_1$  | $\not\Rightarrow$ | $T_1$ -ordered               | $\Rightarrow$     | $T_1$  |
|   |                   | $\Downarrow$                 |                   |  |
| pairwise $T_1$  | $\Leftrightarrow$ | $T_1^K$ -ordered             | $\Rightarrow$     | $T_1$  |
|   |                   | $T_1^K$ -ordered + convex    | $\Leftrightarrow$ | $\left\{ \begin{array}{l} T_1 + \\ (X, \tau, \leq) \text{ convex} \end{array} \right.$ |
| $\left. \begin{array}{l} \text{pairwise} \\ \text{completely regular} \\ + (X, \tau, \leq) \text{ convex} \\ + (X, \tau, \leq) T_1\text{-ordered} \end{array} \right\}$ | $\Rightarrow$     | completely regularly ordered | $\Rightarrow$     | completely regular   |

We note that  $T_1$ -ordered is a strictly stronger property than  $T_1^K$ -ordered. For example, consider the interval  $[0, 1] \subseteq \mathbb{R}$  with the usual topology. Impose the usual order on  $(0, 1]$ , with 0 noncomparable to all other points. This ordered topological space is easily seen to be  $(\tau^\sharp, \tau^\flat)$ -pairwise  $T_1$ , and hence  $T_1^K$ -ordered, but since, for example,  $d(1) = (0, 1]$  is not closed, it is not  $T_1$ -ordered. Furthermore, since  $T_1^K$ -ordered implies  $T_0$ -ordered, this space is its own  $T_0$ -ordered reflection. Now the characterizations in Theorem 6 and Theorem 15 show that this space is  $R_0^K$ -ordered but not  $R_0$ -ordered. Thus,  $R_0$ -ordered is a strictly stronger property than  $R_0^K$ -ordered.

The table above summarizes the implications  $(1) \Rightarrow (2) \Rightarrow (3)$  suggested in the introduction. A bitopological space is *pairwise completely regular* (see [9]) if and only if it admits a compatible quasi-uniformity  $\mathcal{U}$  in the sense that  $\tau(\mathcal{U})$  is the first topology and  $\tau(\mathcal{U}^{-1})$  is the second topology. An ordered topological space is

completely regularly ordered (see [7], [16]) if and only if there is a quasi-uniformity  $\mathcal{U}$  with  $\bigcap \mathcal{U} = \leq$  and  $\tau(\mathcal{U}) \vee \tau(\mathcal{U}^{-1}) = \tau$ . Such spaces are always convex and  $T_1$ -ordered. Completely regular topological spaces are the uniformizable spaces.

The following result is analogous to Theorem 4 (e).

**Theorem 16.** *An ordered topological space  $(X, \tau, \leq)$  is  $R_0^K$ -ordered if and only if there is a lattice isomorphism  $\varphi$  between  $\tau$  and the topology  $\tau_1$  of some  $T_1^K$ -ordered topological space  $(X_1, \tau_1, \leq_1)$  such that the restrictions  $\varphi|_{\tau^\sharp}$  and  $\varphi|_{\tau^\flat}$  are lattice isomorphisms from  $\tau^\sharp$  onto  $\tau_1^\sharp$  and from  $\tau^\flat$  onto  $\tau_1^\flat$ , respectively.*

PROOF. Suppose  $(X, \tau, \leq)$  is an ordered topological space,  $(X_1, \tau_1, \leq_1)$  is a  $T_1^K$ -ordered topological space, and there exists a lattice isomorphism from  $\tau$  to  $\tau_1$  which induces lattice isomorphisms from  $\tau^\sharp$  to  $\tau_1^\sharp$  and from  $\tau^\flat$  to  $\tau_1^\flat$ . Now  $\{x\} = C(x) = I(x) \cap D(x)$  for every  $x \in X_1$ , so every  $\tau_1$ -closed set  $F$  is a disjoint union of minimal nonempty closed sets, each of which is the intersection of a closed increasing set and a closed decreasing set. Correspondingly, in  $X$ , each closed set, and in particular,  $X$ , is partitioned into minimal nonempty closed sets, each of which is the intersection of a closed increasing set and a closed decreasing set. For  $x \in X$ , let  $\hat{x}$  be the member of this partition of  $X$  which contains  $x$ . Then  $\hat{x} = C(x)$ , so  $\{C(x) : x \in X\}$  partitions  $X$ , and thus  $X$  is  $R_0^K$ -ordered.

For the converse, take  $(X_1, \tau_1, \leq_1)$  to be the  $T_0$ -ordered reflection of  $X$ . The result follows from Lemma 5 (a) and (b), noting that closed or open monotone sets in  $X$  are  $\approx$ -saturated. □

As additional evidence that the standard definition of  $T_1$ -ordered and the corresponding definition of  $R_0$ -ordered are not optimal, we show that Theorem 16 does not remain valid if  $R_0^K$ -ordered and  $T_1^K$ -ordered are replaced by  $R_0$ -ordered and  $T_1$ -ordered. Let  $N = \{a, b, c, d\}$  with the order  $\Delta_N \cup \{(a, b), (c, d), (c, b)\}$ , and the topology  $\tau = \tau^\sharp \cup \tau^\flat$  where  $\tau^\sharp = \{\emptyset, \{a, b\}, N\}$  and  $\tau^\flat = \{\emptyset, \{c, d\}, N\}$ . Now  $i(\{c\}) = i(\{c, d\}) \neq N = \bigcap \{U \in \tau^\sharp : c \in U\}$ . Thus  $N$  is not  $R_0$ -ordered. However, if  $(Y, \tau_1, \leq_1)$  is the subspace  $\{a, d\}$  of  $N$ , then  $Y$  is  $T_1$ -ordered and there is a natural lattice isomorphism between  $\tau$  and  $\tau_1$  which induces a lattice isomorphism between  $\tau^\sharp$  ( $\tau^\flat$ ) and  $\tau_1^\sharp$  ( $\tau_1^\flat$ ). Further note that both  $N$  and  $Y$  are convex. One direction of Theorem 16, however, does remain valid if the  $K$  is dropped from  $R_0^K$ -ordered and  $T_1^K$ -ordered. The argument of the last paragraph of the proof still holds. (See also Proposition 6 of [13].)

We also note that for  $(X, \tau, \leq)$  to be  $R_0^K$ -ordered, it is not sufficient to have  $\tau^\sharp$  and  $\tau^\flat$  individually lattice isomorphic to the upper and lower topologies  $\tau_1^\sharp$  and  $\tau_1^\flat$  of a  $T_1^K$ -ordered topological space, even if both spaces are convex. Consider

$X = \{a, b, c, d\}$  with the order  $\Delta_X \cup \{(a, b), (c, d)\}$  and with subbase of open sets  $\{\{b\}, \{c\}, X\}$ . Note that  $\tau^\sharp = \{\emptyset, \{b\}, X\}$  is a 3-element nested chain, as is  $\tau^b = \{\emptyset, \{c\}, X\}$ , and  $(X, \tau, \leq)$  is convex. Now if  $X_1 = \{1, 2\}$  with the discrete topology  $\tau_1$  and usual order  $\leq_1$ , then it is easy to see that  $(X_1, \tau_1, \leq_1)$  is  $T_1$ -ordered and thus  $T_1^K$ -ordered,  $\tau_1^\sharp$  and  $\tau_1^b$  are individually lattice isomorphic to  $\tau^\sharp$  and  $\tau^b$ , yet  $(X, \tau, \leq)$  is not  $R_0^K$ -ordered since, for example,  $a \in C(b)$  but  $b \notin C(a)$ .

#### 4. The Wallman ordered compactification

The Wallman ordered compactification of a  $T_1$ -ordered convex ordered topological space  $(X, \tau, \leq)$  was first constructed by CHOE and PARK [2] using maximal bifilters, and later by KENT [5] using maximal  $c$ -filters. A  $c$ -filter is a filter having a base of  $c$ -sets. Let  $w_0X$  be the set of maximal  $c$ -filters on  $(X, \tau, \leq)$ . For  $A \subseteq X$ , let  $A^* = \{\mathcal{F} \in w_0X : A \in \mathcal{F}\}$ . Then  $\{A^* : A \text{ is a } c\text{-set in } X\}$  is a closed subbase for a topology on  $w_0X$ . A partial order on  $w_0X$  is defined by  $\mathcal{F} \leq \mathcal{G}$  if and only if  $I(\mathcal{F}) \subseteq \mathcal{G}$  and  $D(\mathcal{G}) \subseteq \mathcal{F}$ , where  $I(\mathcal{F})$  is the filter generated by  $\{I(F) : F \in \mathcal{F}\}$  and  $D(\mathcal{G})$  is defined similarly. Now  $w_0X$  with this topology and order is the Wallman ordered compactification. If  $H$  is a monotone open or closed set in  $X$ , then  $H^*$  has the same properties in  $w_0X$ . Choe and Park's earlier construction uses maximal bifilters  $(\mathcal{F}, \mathcal{G})$  where a bifilter is defined to be a pair of filters  $(\mathcal{F}, \mathcal{G})$  such that  $\mathcal{F} \vee \mathcal{G}$  exists,  $\mathcal{F}$  has a base of closed decreasing sets, and  $\mathcal{G}$  has a base of closed increasing sets. The set of maximal bifilters is given the topology having a subbase of closed sets of form  $\{(\mathcal{F}, \mathcal{G}) \in w_0X : D(A) \in \mathcal{F} \text{ and } I(A) \in \mathcal{G}\}$  where  $A$  is any subset of  $X$ , and they order the set of maximal bifilters by  $(\mathcal{F}, \mathcal{G}) \leq (\mathcal{F}', \mathcal{G}')$  if and only if  $\mathcal{F} \subseteq \mathcal{F}'$  and  $\mathcal{G} \subseteq \mathcal{G}'$ . The equivalence of the bifilter construction and the  $c$ -set construction of  $w_0X$  is seen by the bijection  $(\mathcal{F}, \mathcal{G}) \rightarrow \mathcal{F} \vee \mathcal{G}$  from the set of maximal bifilters on  $X$  to the set of maximal  $c$ -filters on  $X$ .

Both constructions of the Wallman ordered compactification use the hypothesis that  $(X, \tau, \leq)$  be  $T_1$ -ordered only to insure that  $X$  is embedded in  $w_0X$ , that is, only to insure that  $\{\mathcal{S}(x)\}$  is a maximal  $c$ -filter or  $(\mathcal{S}(d(x)), \mathcal{S}(i(x)))$  is a maximal bifilter, where  $\mathcal{S}(A)$  represents the collection of supersets of  $A$ . Thus, the  $c$ -filter construction of  $w_0X$  only uses  $T_1$ -ordered to imply the  $T_1^K$ -ordered property, that  $\{x\} = C(x)$  so that  $\mathcal{S}(x)$  is a maximal  $c$ -filter. It follows that Kent's  $c$ -filter construction of  $w_0X$  remains valid for any  $T_1^K$ -ordered space.

If  $X$  is  $T_1$ -ordered, Choe and Park consider the points of  $X \subseteq w_0X$  to be the maximal bifilters  $(\mathcal{S}(d(x)), \mathcal{S}(i(x))) = (\mathcal{S}(D(x)), \mathcal{S}(I(x)))$ . This equality need not hold if  $X$  is only assumed to be  $T_1^K$ -ordered, so to ensure that the bifilters have

bases of closed sets, we must use  $(\mathcal{S}(D(x)), \mathcal{S}(I(x)))$  for the construction of the Wallman ordered compactification of a  $T_1^K$ -ordered space. With this modification, the proof in [2] that  $(\mathcal{S}(d(x)), \mathcal{S}(i(x)))$  is a maximal bifilter for all  $x \in X$  does not show that the bifilter  $(\mathcal{S}(D(x)), \mathcal{S}(I(x)))$  is maximal if  $X$  is  $T_1^K$ -ordered. We remedy this situation with the following Lemma.

**Lemma 17.** *If  $X$  is a convex  $T_1^K$ -ordered topological space and  $x \in X$ , then  $(\mathcal{S}(D(x)), \mathcal{S}(I(x)))$  is a maximal bifilter.*

PROOF. It is easy to see that  $(\mathcal{S}(D(x)), \mathcal{S}(I(x)))$  is a bifilter. If it is not maximal, then there exists a bifilter  $(\mathcal{F}, \mathcal{G}) \supseteq (\mathcal{S}(D(x)), \mathcal{S}(I(x)))$  with either  $\mathcal{S}(D(x)) \neq \mathcal{F}$  or  $\mathcal{S}(I(x)) \neq \mathcal{G}$ . The cases are dual, so we will only consider the case  $\mathcal{S}(D(x)) \neq \mathcal{F}$ . Since  $\mathcal{F}$  has a base of closed decreasing sets, there exists a closed decreasing set  $F \in \mathcal{F}$  such that  $F \notin \mathcal{S}(D(x))$ , or equivalently,  $D(x) \not\subseteq F$ . Since  $D(x) \in \mathcal{F}$ , we have  $A = D(x) \cap F \in \mathcal{F}$ . Now  $A \subseteq F$  implies  $D(x) \not\subseteq A$ , so  $x \notin A$ . We also have  $I(x) \subseteq X \setminus A$ , for if  $y \in I(x) \cap A \subseteq I(x) \cap D(x)$ , then we have the contradiction that  $y = x \notin A$  since, by the  $T_1^K$ -ordered property,  $I(x) \cap D(x) = \{x\}$ . Thus, we have  $X \setminus A \in \mathcal{S}(I(x)) \subseteq \mathcal{G}$  and  $A \in \mathcal{F}$ , contradicting that  $\mathcal{F} \vee \mathcal{G}$  exists. This shows that the bifilter  $(\mathcal{S}(D(x)), \mathcal{S}(I(x)))$  is maximal.  $\square$

The Wallman ordered compactification of a  $T_1$ -ordered space need not be  $T_1$ -ordered, but the theorem below shows the advantage of the  $T_1^K$ -ordered property.

**Theorem 18.** *If  $X$  is any convex  $T_1^K$ -ordered topological space, the Wallman ordered compactification  $w_0X$  is  $T_1^K$ -ordered.*

PROOF. We will show that  $w_0X$  satisfies the characterization of  $T_1^K$ -ordered given in Theorem 10 (a). Suppose  $\mathcal{F}$  and  $\mathcal{G}$  are distinct maximal  $c$ -filters on  $X$ , that is,  $\mathcal{F}$  and  $\mathcal{G}$  are distinct points in  $w_0X$ . Now either  $I(\mathcal{G}) \not\subseteq \mathcal{F}$  or  $I(\mathcal{G}) \subseteq \mathcal{F}$ . In case  $I(\mathcal{G}) \not\subseteq \mathcal{F}$ , there exists  $I(G) \in I(\mathcal{G}) \subseteq \mathcal{G}$  with  $I(G) \notin \mathcal{F}$ . Since  $\mathcal{G}$  is a filter and  $\mathcal{F}$  is maximal, it follows that  $X \setminus I(G) \notin \mathcal{G}$  and  $X \setminus I(G) \in \mathcal{F}$ , so that  $\mathcal{G} \notin (X \setminus I(G))^*$  and  $\mathcal{F} \in (X \setminus I(G))^*$ . Now  $(X \setminus I(G))^*$  is an open decreasing neighborhood of  $\mathcal{F}$  in  $w_0X$  which excludes  $\mathcal{G}$ . In case  $I(\mathcal{G}) \subseteq \mathcal{F}$ , we have  $D(\mathcal{G}) \not\subseteq \mathcal{F}$ , for otherwise  $I(\mathcal{G}) \vee D(\mathcal{G}) = C(\mathcal{G}) = \mathcal{G} \subseteq \mathcal{F}$ , contradicting the maximality of  $\mathcal{G}$ . Now there exists  $D(G) \in \mathcal{G}$  with  $D(G) \notin \mathcal{F}$ , and the dual argument of the previous case shows that  $(X \setminus D(G))^*$  is an open increasing neighborhood of  $\mathcal{F}$  in  $w_0X$  which excludes  $\mathcal{G}$ . In either case, we have found that  $\mathcal{G} \neq \mathcal{F}$  implies  $\mathcal{G} \notin \bigcap \{ \text{open monotone neighborhoods of } \mathcal{F} \}$ , and it follows that  $w_0X$  is  $T_1^K$ -ordered.  $\square$

We mention that the expected properties of the Wallman ordered compactification remain valid even when the construction is applied to  $T_1^K$ -ordered topological spaces. For example, if  $X$  is convex and  $T_1^K$ -ordered,  $\varphi : X \rightarrow w_0X$  is the natural embedding, and  $f : X \rightarrow Y$  is a continuous increasing function from  $X$  into an arbitrary compact  $T_2$ -ordered space  $Y$  (i.e.,  $Y$  is compact and the graph of its order is closed in  $Y \times Y$ ), then there exist a unique continuous increasing function  $\bar{f} : w_0X \rightarrow Y$  such that  $\bar{f} \circ \varphi = f$ .

### References

- [1] K. BELAID, O. ECHI, and S. LAZAAR,  $T_{(\alpha,\beta)}$ -spaces and the Wallman compactification, *Internat. J. Math. & Math. Sci.* **68** (2004), 3717–3735.
- [2] T. H. CHOE and Y. S. PARK, Wallman's Type Order Compactification, *Pacific J. Math.* **82** (2) (1979), 339–347.
- [3] A. S. DAVIS, Indexed Systems of Neighborhoods for General Topological Spaces, *The Amer. Math. Monthly* **68** (9) (1961), 886–893.
- [4] B. P. DVALISHVILI, Bitopological Spaces: Theory, Relations With Generalized Algebraic Structures, and Applications, Vol. 199, North-Holland Mathematics Studies, *Elsevier Science B. V., Amsterdam*, 2005.
- [5] D. C. KENT, On the Wallman Order Compactification, *Pacific J. Math.* **118** (1985), 159–163.
- [6] D. C. KENT and T. A. RICHMOND, Separation properties of the Wallman ordered compactification, *Internat. J. Math. & Math. Sci.* **13** (2) (1990), 209–222.
- [7] H.-P. A. KÜNZI and T. A. RICHMOND, Completely regularly ordered spaces versus  $T_2$ -ordered spaces which are completely regular, *Topology Appl.* **135** (1–3) (2004), 185–196.
- [8] H.-P. A. KÜNZI and T. A. RICHMOND,  $T_i$ -ordered reflections, *Applied General Topology* **6** (2) (2005), 207–216.
- [9] E. P. LANE, Bitopological spaces and quasi-uniform spaces, *Proc. London Math. Soc.* (3) **17** (1967), 241–256.
- [10] S. D. MCCARTAN, Separation axioms for topological ordered spaces, *Proc. Cambridge Philos. Soc.* **64** (1968), 965–973.
- [11] D. N. MISRA and K. K. DUBE, Pairwise  $R_0$ -space, *Annales de la Société Scientifique de Bruxelles* **87**, I (1973), 3–15.
- [12] M. MRŠEVIĆ, On bitopological separation axioms, *Math. Vesnik* **38** (1986), 313–318.
- [13] M. MRŠEVIĆ, On pairwise  $R_0$  and pairwise  $R_1$  bitopological spaces, *Bull. Math. Soc. Sci. Math. R. S. Roumanie* **30** (78) (1986), 141–148.
- [14] M. G. MURDESHWAR and S. A. NAIMPALLY, Quasi-Uniform Topological Spaces, *Noordhoof, Groningen*, 1966.
- [15] L. NACHBIN, Topology and Order., *Van Nostrand Math. Studies 4, Princeton, N. J.*, 1965.
- [16] K. R. NAILANA, Strict complete regularity in the categories of bitopological spaces and ordered topological spaces, *Publ. Math. Debrecen* **58** (4) (2001), 693–705.
- [17] I. REILLY, On Pairwise  $R_0$  Spaces, *Annales de la Société Scientifique de Bruxelles* **88**, III (1974), 293–296.



- [18] M. J. SAEGROVE, Pairwise completely regularity and compactification in bitopological spaces, *J. London Math. Soc. (2)* **7** (1973), 286–290.
- [19] J. SWART, Total disconnectedness in bitopological spaces and product bitopological spaces, *Nederl. Akad. Wetensch. Proc. Ser. A* 74 Indag. Math. **33** (1971), 135–145.
- [20] M. R. ŽIŽOVIC, Some properties of bitopological spaces, *Math. Vesnik* **11** (26) (1974), 233–237.

HANS-PETER A. KÜNZI  
DEPARTMENT OF MATHEMATICS AND APPLIED MATHEMATICS  
UNIVERSITY OF CAPE TOWN  
RONDEBOSCH 7701  
SOUTH AFRICA

*E-mail:* [hans-peter.kunzi@uct.ac.za](mailto:hans-peter.kunzi@uct.ac.za)

AISLING E. MCCLUSKEY  
DEPARTMENT OF MATHEMATICS  
NATIONAL UNIVERSITY OF IRELAND, GALWAY  
GALWAY  
IRELAND

*E-mail:* [aisling.mccluskey@nuigalway.ie](mailto:aisling.mccluskey@nuigalway.ie)

THOMAS A. RICHMOND  
DEPARTMENT OF MATHEMATICS  
WESTERN KENTUCKY UNIVERSITY  
BOWLING GREEN, KY 42101  
USA

*E-mail:* [tom.richmond@wku.edu](mailto:tom.richmond@wku.edu)

*(Received March 3, 2008; revised June 25, 2008)*