

Year: 2010

Vol.: 77

Fasc.: 3-4

**Title:** On  $(a, b)$ -balancing numbers

**Author(s):** Tünde Kovács , Kálmán Liptai and Péter Olajos

A positive integer  $n$  is called a balancing number if  $1 + \dots + (n - 1) = (n + 1) + \dots + (n + r)$  for some positive integer  $r$ . Balancing numbers and their generalizations have been investigated by several authors, from many aspects. In this paper we introduce the concept of balancing numbers in arithmetic progressions, and prove several effective finiteness and explicit results about them. In the proofs of our results, among others, we combine Baker's method, the modular method developed by Wiles and others, a result of Bennett about the diophantine equation  $|ax^n - by^n| = 1$ , the Chabauty method and the theory of elliptic curves.

**Address:**

Tünde Kovács  
Institute of Mathematics  
University of Debrecen  
H-4010 Debrecen, P.O. Box 12  
Hungary

**Address:**

Kálmán Liptai  
Institute of Mathematics and Informatics  
Eszterházy Károly College  
H-3300 Eger, Eszterházy tér 1  
Hungary

**Address:**

Péter Olajos  
Department of Applied Mathematics  
University of Miskolc  
H-3515 Miskolc-Egyetemváros  
Hungary