

## Schur power convexity of Stolarsky means

By ZHEN-HANG YANG (Hangzhou)

**Abstract.** In this paper, the Schur convexity is generalized to Schur  $f$ -convexity, which contains the Schur geometrical convexity, Schur harmonic convexity and so on. When  $f : \mathbb{R}_+ \rightarrow \mathbb{R}$  is defined by  $f(x) = (x^m - 1)/m$  if  $m \neq 0$  and  $f(x) = \ln x$  if  $m = 0$ , the necessary and sufficient conditions for  $f$ -convexity (is called Schur  $m$ -power convexity) of Stolarsky means are given, which generalized and unified certain known results.

### 1. Introduction and main results

Let  $p, q \in \mathbb{R}$  and  $a, b \in \mathbb{R}_+ := (0, \infty)$  with  $a \neq b$ . The so-called Stolarsky means  $S_{p,q}(a, b)$  are defined by

$$S_{p,q}(a, b) = \begin{cases} \left( \frac{q(a^p - b^p)}{p(a^q - b^q)} \right)^{\frac{1}{p-q}} & \text{if } pq(p-q) \neq 0, \\ \left( \frac{a^p - b^p}{p(\ln a - \ln b)} \right)^{\frac{1}{p}} & \text{if } p \neq 0, q = 0, \\ \left( \frac{a^q - b^q}{q(\ln a - \ln b)} \right)^{\frac{1}{q}} & \text{if } q \neq 0, p = 0, \\ \exp \left( \frac{a^p \ln a - b^p \ln b}{a^p - b^p} - \frac{1}{p} \right) & \text{if } p = q \neq 0, \\ \sqrt{ab} & \text{if } p = q = 0. \end{cases} \quad (1.1)$$

---

*Mathematics Subject Classification:* Primary: 26B25, 26E60; Secondary: 26D15, 26D99.

*Key words and phrases:* Schur convexity, Schur  $f$ -convexity, Schur power convexity, Stolarsky means.

This paper is in final form and no version of it will be submitted for publication elsewhere.

Also,  $S_{p,q}(a, a) = a$ . It is known that the Stolarsky means  $S_{p,q}(a, b)$  are  $C^\infty$  function on the domain  $\{(p, q, a, b) : p, q \in \mathbb{R}, a, b \in \mathbb{R}_+\}$  (see [19, Lemma 1]), and obviously symmetric with respect to  $a, b$  and  $p, q$ .

Most of the classical two variable means are special cases of  $S_{p,q}(a, b)$ , for example,  $S_{1,2} = A$  is the arithmetic means,  $S_{0,0} = G$  is the geometric mean,  $S_{-1,-2} = H$  is the harmonic mean,  $S_{1,0} = L$  is the logarithmic mean,  $S_{1,1} = I$  is the identric mean (exponential mean), and more generally, the  $r$ -th power mean is equal to  $S_{r,2r}$ . The basic properties of Stolarsky means, as well as their comparison theorems, log-convexity, and inequalities were studied in papers [3], [8], [12], [15], [16], [18], [19], [20], [23], [24], [25], [26], [27], [28], [36], [37], [42], [43], [44], [46], [47].

Schur convexity was introduced by Schur in 1923 [21], and it has many important applications in analytic inequalities [2], [11], [49], linear regression [35], graphs and matrices [7], combinatorial optimization [14], information-theoretic topics [9], Gamma functions [22], stochastic orderings [32], reliability [13], and other related fields.

In recent years, the Schur convexity and Schur geometrical convexity of  $S_{p,q}(a, b)$  have attracted the attention of a considerable number of mathematicians [4], [5, 17], [29], [30], [31], [33]. QI [30] first proved that the Stolarsky means  $S_{p,q}(a, b)$  are Schur convex on  $(-\infty, 0] \times (-\infty, 0]$  and Schur concave on  $[0, \infty) \times [0, \infty)$  with respect to  $(p, q)$  for fixed  $a, b > 0$  with  $a \neq b$ . YANG [45] improved Qi's result and proved that Stolarsky means  $S_{p,q}(a, b)$  are Schur convex with respect to  $(p, q)$  for fixed  $a, b > 0$  with  $a \neq b$  if and only if  $p + q < 0$  and Schur concave if and only if  $p + q > 0$ .

QI *et al.* [29] tried to obtain the Schur convexity of  $S_{p,q}(a, b)$  with respect to  $(a, b)$  for fixed  $(p, q)$  and declared an incorrect conclusion. SHI *et al.* [33] observed that the above conclusion is wrong and obtained a sufficient condition for the Schur convexity of  $S_{p,q}(a, b)$  with respect to  $(a, b)$ . CHU and ZHANG [5] improved Shi's results and gave a necessary and sufficient condition. This perfectly solved the Schur convexity of Stolarsky means with respect to  $(a, b)$ .

The Schur geometrical convexity was introduced by ZHANG [50], and there has many interesting results [10], [34], [39], [40]. For the Schur geometrical convexity of Stolarsky means  $S_{p,q}(a, b)$ , CHU and ZHANG [4] proved that they are Schur geometrically convex with respect to  $(a, b) \in \mathbb{R}_+^2$  if  $p + q \geq 0$  and Schur geometrically concave if  $p + q \leq 0$ . LI *et al.* [17] also investigated the Schur geometrical convexity of generalized exponent mean  $I_p(a, b)$ . In 2010, a necessary and sufficient condition for Schur geometrical convexity of the four-parameter

means with respect to a pair of parameters was given in [48]. This gives a unified treatment for Schur geometrical convexity of Stolarsky and Gini means.

Recently, ANDERSON *et al.* [1] discussed an attractive class of inequalities, which arise from the notation of harmonic convexity. And then it was started to research for *Schur harmonic convexity*. CHU *et al.* [6] showed that the Hamy symmetric function is Schur harmonic convex and obtained some analytic inequalities including the well-known Weierstrass inequalities. XIAO [41] proved that the Lehmer mean values  $L_p(a, b)$  are Schur harmonic convex (Schur harmonic concave) with respect to  $(a, b) \in \mathbb{R}_+^2$  if and only if  $p \geq (\leq) 0$ .

The purpose of this paper is to generalize the notion of Schur convexity and investigate the so-called *Schur power convexity* of Stolarsky means  $S_{p,q}(a, b)$ . Our main results are as follows.

**Theorem 1.** For  $m > 0$  and fixed  $(p, q) \in \mathbb{R}^2$ , Stolarsky mean  $S_{p,q}(a, b)$  is Schur  $m$ -power convex with respect to  $(a, b) \in \mathbb{R}_+^2$  if and only if  $p + q \geq 3m$  and  $\min(p, q) \geq m$ .

**Theorem 2.** For  $m > 0$  and fixed  $(p, q) \in \mathbb{R}^2$ , Stolarsky mean  $S_{p,q}(a, b)$  is Schur  $m$ -power concave with respect to  $(a, b) \in \mathbb{R}_+^2$  if and only if  $p + q \leq 3m$  and  $\min(p, q) \leq m$ .

**Theorem 3.** For  $m < 0$  and fixed  $(p, q) \in \mathbb{R}^2$ , Stolarsky mean  $S_{p,q}(a, b)$  is Schur  $m$ -power convex with respect to  $(a, b) \in \mathbb{R}_+^2$  if and only if  $p + q \geq 3m$  and  $\max(p, q) \geq m$ .

**Theorem 4.** For  $m < 0$  and fixed  $(p, q) \in \mathbb{R}^2$ , Stolarsky mean  $S_{p,q}(a, b)$  is Schur  $m$ -power concave with respect to  $(a, b) \in \mathbb{R}_+^2$  if and only if  $p + q \leq 3m$  and  $\max(p, q) \leq m$ .

**Theorem 5.** For  $m = 0$  and fixed  $(p, q) \in \mathbb{R}^2$ , Stolarsky mean  $S_{p,q}(a, b)$  is Schur  $m$ -power convex (Schur  $m$ -power concave) with respect to  $(a, b) \in \mathbb{R}_+^2$  if and only if  $p + q \geq (\leq) 0$ .

The organization of the paper is as follows. In Section 2, based on the notion and lemmas of Schur convexity, we introduce the definition of Schur  $f$ -convex and Schur  $f$ -concave function, and prove decision theorem for Schur  $f$ -convexity. As a special case, the definition and decision theorem of Schur power convexity are deduced. In Section 3, some lemmas are given. In Section 4, our main results are proved.

## 2. Schur $f$ -convexity and Schur power convexity

For convenience of readers, we recall some definitions as follows.

*Definition 1* ([21], [38]). Let  $\mathbf{x} = (x_1, x_2, \dots, x_n)$  and  $\mathbf{y} = (y_1, y_2, \dots, y_n) \in \mathbb{R}^n (n \geq 2)$ .

(i)  $x$  is said to be majorized by  $y$  (in symbol  $\mathbf{x} \prec \mathbf{y}$ ) if

$$\sum_{i=1}^k x_{[i]} \leq \sum_{i=1}^k y_{[i]} \quad \text{for } 1 \leq k \leq n-1, \quad \sum_{i=1}^n x_{[i]} = \sum_{i=1}^n y_{[i]}, \quad (2.1)$$

where  $x_{[1]} \geq x_{[2]} \cdots \geq x_{[n]}$  and  $y_{[1]} \geq y_{[2]} \cdots \geq y_{[n]}$  are rearrangements of  $\mathbf{x}$  and  $\mathbf{y}$  in a decreasing order.

- (ii)  $\mathbf{x} \geq \mathbf{y}$  means  $x_i \geq y_i$  for all  $i = 1, 2, \dots, n$ . Let  $\Omega \subseteq \mathbb{R}^n (n \geq 2)$ . The function  $\phi : \Omega \rightarrow \mathbb{R}$  is said to be increasing if  $\mathbf{x} \geq \mathbf{y}$  implies that  $\phi(\mathbf{x}) \geq \phi(\mathbf{y})$ .  $\phi$  is said to be decreasing if and only if  $-\phi$  is increasing.
- (iii)  $\Omega \subseteq \mathbb{R}^n$  is called a convex set if  $(\alpha x_1 + \beta y_1, \dots, \alpha x_n + \beta y_n) \in \Omega$  for all  $\mathbf{x}, \mathbf{y}$  and all  $\alpha, \beta \in [0, 1]$  with  $\alpha + \beta = 1$ .
- (iv) Let  $\Omega \subseteq \mathbb{R}^n (n \geq 2)$  be a set with nonempty interior. Then  $\phi : \Omega \rightarrow \mathbb{R}$  is said to be Schur convex if  $\mathbf{x} \prec \mathbf{y}$  on  $\Omega$  implies that  $\phi(\mathbf{x}) \leq \phi(\mathbf{y})$ .  $\phi$  is said to be Schur concave if  $-\phi$  is Schur convex.

*Definition 2* ([21]). (i)  $\Omega \subseteq \mathbb{R}^n (n \geq 2)$  is called symmetric set, if  $\mathbf{x} \in \Omega$  implies that  $\mathbf{xP} \in \Omega$  for every  $n \times n$  permutation matrix  $\mathbf{P}$ .

(ii) The function  $\phi : \Omega \rightarrow \mathbb{R}^n$  is called symmetric if for every permutation matrix  $\mathbf{P}$ ,  $\phi(\mathbf{xP}) = \phi(\mathbf{x})$  for all  $\mathbf{x} \in \Omega$ .

For the Schur convexity, there is the following well-known result.

**Lemma 1** ([21], [38]). Let  $\Omega \subseteq \mathbb{R}^n$  be a symmetric set with nonempty interior  $\Omega^0$  and  $\phi : \Omega \rightarrow \mathbb{R}$  be continuous on  $\Omega$  and differentiable in  $\Omega^0$ . Then  $\phi$  is Schur convex (Schur concave) on  $\Omega$  if and only if  $\phi$  is symmetric on  $\Omega$  and

$$(x_1 - x_2) \left( \frac{\partial \phi}{\partial x_1} - \frac{\partial \phi}{\partial x_2} \right) \geq (\leq) 0 \quad (2.2)$$

Next let us define the Schur  $f$ -convexity as follows.

*Definition 3.* Let  $\Omega = \mathbb{U}^n (\mathbb{U} \subseteq \mathbb{R})$  and  $f$  be a strictly monotone function defined on  $\mathbb{U}$ . Denote by

$$f(\mathbf{x}) = (f(x_1), f(x_2), \dots, f(x_n)) \quad \text{and} \quad f(\mathbf{y}) = (f(y_1), f(y_2), \dots, f(y_n)).$$

- (i)  $\Omega$  is called a  $f$ -convex set if  $(f^{-1}(\alpha f(x_1) + \beta f(y_1)), \dots, f^{-1}(\alpha f(x_n) + \beta f(y_n))) \in \Omega$  for all  $\mathbf{x}, \mathbf{y} \in \Omega$  and all  $\alpha, \beta \in [0, 1]$  with  $\alpha + \beta = 1$ .
- (ii) Let  $\Omega$  be a set with nonempty interior. Then function  $\phi : \Omega \rightarrow \mathbb{R}$  is said to be Schur  $f$ -convex on  $\Omega$  if  $f(\mathbf{x}) \prec f(\mathbf{y})$  on  $\Omega$  implies that  $\phi(\mathbf{x}) \leq \phi(\mathbf{y})$ .

$\phi$  is said to be Schur  $f$ -concave if  $-\phi$  is Schur  $f$ -convex.

*Remark 1.* Let  $\Omega = \mathbb{U}^n$  ( $\mathbb{U} \subseteq \mathbb{R}$ ) and  $f$  be a strictly monotone function defined on  $\mathbb{U}$  and  $f(\Omega) = \{f(\mathbf{x}) : \mathbf{x} \in \Omega\}$ . Then function  $\phi : \Omega \rightarrow \mathbb{R}$  is Schur  $f$ -convex (Schur  $f$ -concave) if and only if  $\phi \circ f^{-1}$  is Schur convex (Schur concave) on  $f(\Omega)$ .

Indeed, if function  $\phi : \Omega \rightarrow \mathbb{R}$  is Schur  $f$ -convex, then  $\forall \mathbf{x}', \mathbf{y}' \in f(\Omega)$ , there are  $\mathbf{x}, \mathbf{y} \in \Omega$  such that  $\mathbf{x}' = f(\mathbf{x}), \mathbf{y}' = f(\mathbf{y})$ . If  $f(\mathbf{x}) \prec f(\mathbf{y})$ , that is,  $\mathbf{x}' \prec \mathbf{y}'$ , then  $\phi(\mathbf{x}) \leq \phi(\mathbf{y})$ , that is,  $\phi((f^{-1}(\mathbf{x}')) \leq \phi((f^{-1}(\mathbf{y}'))$ . This shows that  $\phi \circ f^{-1}$  is Schur convex on  $f(\Omega)$ . Conversely, if  $\phi \circ f^{-1}$  is Schur convex on  $f(\Omega)$ , then  $\forall \mathbf{x}, \mathbf{y} \in \Omega$  such that  $f(\mathbf{x}) \prec f(\mathbf{y})$ , we have  $\phi((f^{-1}(f(\mathbf{x}))) \leq \phi((f^{-1}(f(\mathbf{y})))$ , that is,  $\phi(\mathbf{x}) \leq \phi(\mathbf{y})$ . This indicates  $\phi$  is Schur  $f$ -convex on  $\Omega$ .

In the same way, we can show that  $\phi$  is Schur  $f$ -concave on  $\Omega$  if and only if  $\phi \circ f^{-1}$  is Schur concave on  $f(\Omega)$ .

*Remark 2.* Let  $\Omega \subseteq \mathbb{R}^n$  ( $n \geq 2$ ) be a symmetric set and the function  $\phi : \Omega \rightarrow \mathbb{R}$  be Schur  $f$ -convex (Schur  $f$ -concave). Then  $\phi$  is symmetric on  $\Omega$ .

In fact, for any  $\mathbf{x} \in \Omega$  and every permutation matrix  $\mathbf{P}$ , we have  $\mathbf{xP} \in \Omega$ . Note  $\mathbf{xP}$  is another permutation of  $\mathbf{x}$ , hence  $f(\mathbf{x}) \prec f(\mathbf{xP}) \prec f(\mathbf{x})$ . Since  $\phi$  is Schur  $f$ -convex (Schur  $f$ -concave), we have  $\phi(\mathbf{x}) \leq (\geq) \phi(\mathbf{xP}) \leq (\geq) \phi(\mathbf{x})$ , that is,  $\phi(\mathbf{xP}) = \phi(\mathbf{x})$  for all  $\mathbf{x} \in \Omega$ . This shows that  $\phi$  is symmetric on  $\Omega$ .

By Lemma 1 and Remark 1, 2, we have the following

**Theorem 6.** Assume that  $\Omega = \mathbb{U}^n$  ( $\mathbb{U} \subseteq \mathbb{R}$ ) is a symmetric set with nonempty interior  $\Omega^0$ ,  $f$  is a strictly monotone and derivable function defined on  $\mathbb{U}$ , and  $\phi : \Omega \rightarrow \mathbb{R}$  is continuous on  $\Omega$  and differentiable in  $\Omega^0$ . Then  $\phi$  is Schur  $f$ -convex (Schur  $f$ -concave) on  $\Omega$  if and only if  $\phi$  is symmetric on  $\Omega$  and

$$(f(x_1) - f(x_2)) \left( \frac{1}{f'(x_1)} \frac{\partial \phi}{\partial x_1} - \frac{1}{f'(x_2)} \frac{\partial \phi}{\partial x_2} \right) \geq (\leq) 0 \quad (2.3)$$

holds for any  $\mathbf{x} = (x_1, x_2, \dots, x_n) \in \Omega^0$  with  $x_1 \neq x_2$ .

PROOF. We easily check that  $\phi \circ f^{-1}$  is symmetric on  $f(\Omega)$  if and only if  $\phi$  is symmetric on  $\Omega$ .

By Remark 1 and Lemma 1,  $\phi \circ f^{-1}$  is Schur convex (Schur concave) if and only if  $\phi \circ f^{-1}$  is symmetric on  $f(\Omega)$  and

$$(y_1 - y_2) \left( \frac{\partial(\phi \circ f^{-1})}{\partial y_1} - \frac{\partial(\phi \circ f^{-1})}{\partial y_2} \right) \geq (\leq) 0$$

holds for any  $\mathbf{y} \in f(\Omega)^0$  with  $y_1 \neq y_2$ . Substituting  $f^{-1}(y) = x$  yields (2.3), where  $\mathbf{x} = (x_1, x_2, \dots, x_n) \in \Omega^0$  with  $x_1 \neq x_2$ .

This proof is finished.  $\square$

Putting  $f(x) = 1, \ln x, x^{-1}$  in Definition 3 yield the Schur convexity, Schur geometrical convexity and Schur harmonic convexity. It is clear that the Schur  $f$ -convexity is a generalization of the Schur convexity mentioned above. In general, we have

*Definition 4.* Let  $f : \mathbb{R}_+ \rightarrow \mathbb{R}$  be defined by  $f(x) = (x^m - 1)/m$  if  $m \neq 0$  and  $f(x) = \ln x$  if  $m = 0$ . Then function  $\phi : \Omega(\subseteq \mathbb{R}_+^n) \rightarrow \mathbb{R}$  is said to be Schur  $m$ -power convex on  $\Omega$  if  $f(\mathbf{x}) \prec f(\mathbf{y})$  on  $\Omega$  implies that  $\phi(\mathbf{x}) \leq \phi(\mathbf{y})$ .

$\phi$  is said to be Schur  $m$ -power concave if  $-\phi$  is Schur  $m$ -power convex.

For Schur power convexity, by Theorem 6 we have

**Corollary 1.** Let  $\Omega \subseteq \mathbb{R}_+^n$  be a symmetric set with nonempty interior  $\Omega^0$  and  $\phi : \Omega \rightarrow \mathbb{R}$  be continuous on  $\Omega$  and differentiable in  $\Omega^0$ . Then  $\phi$  is Schur  $m$ -power convex (Schur  $m$ -power concave) on  $\Omega$  if and only if  $\phi$  is symmetric on  $\Omega$  and

$$\frac{x_1^m - x_2^m}{m} \left( x_1^{1-m} \frac{\partial \phi}{\partial x_1} - x_2^{1-m} \frac{\partial \phi}{\partial x_2} \right) \geq (\leq) 0 \quad \text{if } m \neq 0, \quad (2.4)$$

$$(\ln x_1 - \ln x_2) \left( x_1 \frac{\partial \phi}{\partial x_1} - x_2 \frac{\partial \phi}{\partial x_2} \right) \geq (\leq) 0 \quad \text{if } m = 0 \quad (2.5)$$

holds for any  $\mathbf{x} = (x_1, x_2, \dots, x_n) \in \Omega^0$  with  $x_1 \neq x_2$ .

### 3. Lemmas

**Lemma 2.** For fixed  $(p, q) \in \mathbb{R}^2$ , Stolarsky mean  $S_{p,q}(a, b)$  is Schur  $m$ -power convex (Schur  $m$ -power concave) with respect to  $(a, b) \in \mathbb{R}_+^2$  if and only if  $g(t) \geq (\leq) 0$  for all  $t > 0$ , where

$$g(t) = g_{p,q}(t) = \begin{cases} \frac{(p-q) \sinh At - p \sinh Bt - q \sinh Ct}{pq(p-q)} & \text{if } pq(p-q) \neq 0, \\ \frac{\sinh(p-m)t + \sinh(p+m)t - 2pt \cosh(p-m)t}{-p^2} & \text{if } p \neq 0, q = 0, \\ \frac{\sinh(q-m)t + \sinh(q+m)t - 2qt \cosh(q-m)t}{-q^2} & \text{if } q \neq 0, p = 0, \\ \frac{\sinh(2p-m)t + \sinh mt - 2pt \cosh mt}{p^2} & \text{if } p = q \neq 0, \\ -2t^2 \sinh mt & \text{if } p = q = 0, \end{cases} \quad (3.1)$$

and

$$A = p + q - m, \quad B = p - q - m, \quad C = p - q + m, \quad (3.2)$$

PROOF. Let  $m \neq 0$  and  $S = S_{p,q} := S_{p,q}(a, b)$  defined by (1.1).

In the case of  $pq(p - q) \neq 0$ . We have

$$\begin{aligned} \frac{\partial \ln S}{\partial a} &= \frac{1}{S} \frac{\partial S}{\partial a} = \frac{1}{p - q} \left( \frac{pa^{p-1}}{a^p - b^p} - \frac{qa^{q-1}}{a^q - b^q} \right), \\ \frac{\partial \ln S}{\partial b} &= \frac{1}{S} \frac{\partial S}{\partial b} = \frac{1}{p - q} \left( \frac{-pb^{p-1}}{a^p - b^p} - \frac{-qb^{q-1}}{a^q - b^q} \right), \end{aligned}$$

hence,

$$a^{1-m} \frac{\partial S}{\partial a} - b^{1-m} \frac{\partial S}{\partial b} = \frac{S}{p - q} \left( p \frac{a^{p-m} + b^{p-m}}{a^p - b^p} - q \frac{a^{q-m} + b^{q-m}}{a^q - b^q} \right).$$

Substituting  $\ln \sqrt{a/b} = t$  and using  $\sinh x = \frac{1}{2}(e^x - e^{-x})$ ,  $\cosh x = \frac{1}{2}(e^x + e^{-x})$ , the right hand side above can be written as

$$\begin{aligned} a^{1-m} \frac{\partial S}{\partial a} - b^{1-m} \frac{\partial S}{\partial b} &= \frac{S(ab)^{-m/2}}{p - q} \left( p \frac{\cosh(p - m)t}{\sinh pt} - q \frac{\cosh(q - m)t}{\sinh qt} \right) \\ &= \frac{S}{2(ab)^{m/2}} \frac{p}{\sinh pt} \frac{q}{\sinh qt} \\ &\quad \cdot \frac{p \cosh(p - m)t \sinh qt - q \cosh(q - m)t \sinh pt}{pq(p - q)}. \end{aligned}$$

Using the “product into sum” formula for hyperbolic functions and (3.2), we have

$$\begin{aligned} \Delta &:= \frac{a^m - b^m}{m} \left( a^{1-m} \frac{\partial S_{p,q}}{\partial a} - b^{1-m} \frac{\partial S_{p,q}}{\partial b} \right) \\ &= \frac{a^m - b^m}{m(a - b)} \frac{(a - b)S_{p,q}}{2(ab)^{m/2}} \frac{p}{\sinh pt} \frac{q}{\sinh qt} \frac{(p - q) \sinh At - p \sinh Bt - q \sinh Ct}{pq(p - q)} \\ &= d_{p,q}(t) \cdot g_{p,q}(t), \end{aligned}$$

where

$$d_{p,q}(t) = \frac{a^m - b^m}{m(a - b)} \frac{(a - b)S_{p,q}}{2(ab)^{m/2}} \frac{p}{\sinh pt} \frac{q}{\sinh qt} \quad (pq(p - q) \neq 0) \quad (3.3)$$

and  $g_{p,q}(t)$  is defined by (3.1).

In the case of  $p \neq q = 0$ . Since  $S_{p,q}(a, b) \in C^\infty$  we have

$$\frac{\partial S_{p,0}}{\partial a} = \lim_{q \rightarrow 0} \frac{\partial S_{p,q}}{\partial a}, \quad \frac{\partial S_{p,0}}{\partial b} = \lim_{q \rightarrow 0} \frac{\partial S_{p,q}}{\partial b},$$

$$\begin{aligned}\frac{\partial S_{p,p}}{\partial a} &= \lim_{q \rightarrow p} \frac{\partial S_{p,q}}{\partial a}, & \frac{\partial S_{p,p}}{\partial b} &= \lim_{q \rightarrow p} \frac{\partial S_{p,q}}{\partial b}, \\ \frac{\partial S_{0,0}}{\partial a} &= \lim_{p \rightarrow 0} \frac{\partial S_{p,p}}{\partial a}, & \frac{\partial S_{0,0}}{\partial b} &= \lim_{p \rightarrow 0} \frac{\partial S_{p,p}}{\partial b}.\end{aligned}$$

It follows that

$$\begin{aligned}\Delta &= \frac{a^m - b^m}{m} \left( a^{1-m} \frac{\partial S_{p,0}}{\partial a} - b^{1-m} \frac{\partial S_{p,0}}{\partial b} \right) \\ &= \lim_{q \rightarrow 0} \left( \frac{a^m - b^m}{m} \left( a^{1-m} \frac{\partial S_{p,q}}{\partial a} - b^{1-m} \frac{\partial S_{p,q}}{\partial b} \right) \right) \\ &= \lim_{q \rightarrow 0} (d_{p,q}(t) g_{p,q}(t)) = g_{p,0}(t) \lim_{q \rightarrow 0} d_{p,q}(t).\end{aligned}$$

Likewise, in the case of  $q \neq p = 0$ , we have

$$\Delta = \frac{a^m - b^m}{m} \left( a^{1-m} \frac{\partial S_{0,q}}{\partial a} - b^{1-m} \frac{\partial S_{0,q}}{\partial b} \right) = g_{0,q}(t) \lim_{p \rightarrow 0} d_{p,q}(t).$$

In the case of  $p = q \neq 0$ , we have

$$\Delta = \frac{a^m - b^m}{m} \left( a^{1-m} \frac{\partial S_{p,p}}{\partial a} - b^{1-m} \frac{\partial S_{p,p}}{\partial b} \right) = g_{p,p}(t) \lim_{q \rightarrow p} d_{p,q}(t),$$

In the case of  $p = q = 0$ , we have

$$\Delta = \frac{a^m - b^m}{m} \left( a^{1-m} \frac{\partial S_{0,0}}{\partial a} - b^{1-m} \frac{\partial S_{0,0}}{\partial b} \right) = g_{0,0}(t) \lim_{p \rightarrow 0} d_{p,p}(t),$$

Summarizing all cases above yield

$$\begin{aligned}\Delta &= \frac{a^m - b^m}{m} \left( a^{1-m} \frac{\partial S}{\partial a} - b^{1-m} \frac{\partial S}{\partial b} \right) \tag{3.4} \\ &= \begin{cases} g_{p,q}(t) \cdot d_{p,q}(t) & \text{if } pq(p-q) \neq 0, \\ g_{p,0}(t) \lim_{q \rightarrow 0} d_{p,q}(t) & \text{if } p \neq 0, q = 0, \\ g_{0,q}(t) \lim_{p \rightarrow 0} d_{p,q}(t) & \text{if } q \neq 0, p = 0, \\ g_{p,p}(t) \lim_{q \rightarrow p} d_{p,q}(t) & \text{if } p = q \neq 0, \\ g_{0,0}(t) \lim_{p \rightarrow 0} d_{p,0}(t) & \text{if } p = q = 0. \end{cases} \tag{3.5}\end{aligned}$$

Since  $\Delta$  is symmetric with respect to  $a$  and  $b$ , without loss of generality we assume that  $a > b$ . It is easy to verify that  $\frac{a^m - b^m}{m(a-b)} > 0$ ,  $\frac{(a-b)S_{p,q}}{2(ab)^{m/2}} > 0$ ,  $\frac{p}{\sinh pt} \cdot \frac{q}{\sinh qt} > 0$  if  $pq(p-q) \neq 0$  for  $t = \ln \sqrt{a/b} > 0$ , which implies that  $d_{p,q}(t)$  and its limits at  $(p, q) \in \{(p, q) : pq(p-q) = 0\}$  are all positive. Thus by Corollary 1 Stolarsky mean  $S_{p,q}(a, b)$  is Schur  $m$ -power convex (Schur  $m$ -power concave) with respect to  $(a, b) \in \mathbb{R}_+^2$  if and only if  $\Delta \geq (\leq) 0$  if and only if  $g(t) = g_{p,q}(t) \geq (\leq) 0$  for all  $t > 0$ .

It is easy to check that for  $m = 0$  this lemma is also true.

This Lemma is proved.  $\square$

**Lemma 3.** *Both the  $g(t) = g_{p,q}(t)$  defined by (3.1) and  $g'(t) := \partial g_{p,q}(t)/\partial t$  are symmetric with respect to  $p$  and  $q$ , and continuous with respect to  $(p, q)$  on  $\mathbb{R}^2$ .*

PROOF. Firstly, it is easy to check  $g_{p,q}(t)$  is symmetric with respect to  $p$  and  $q$ . Hence,  $\partial g_{p,q}(t)/\partial t = \partial g_{q,p}(t)/\partial t$ , which implies that  $\partial g_{p,q}(t)/\partial t$  is also symmetric with respect to  $p$  and  $q$ .

Secondly, by the proof of Lemma 2, it is easy to see that  $g(t) = g_{p,q}(t)$  is continuous with respect to  $(p, q)$  on  $\mathbb{R}^2$ .

Lastly, we prove  $g'(t) = \partial g_{p,q}(t)/\partial t$  is also continuous with respect to  $(p, q)$  on  $\mathbb{R}^2$ .

A simple calculation yields

$$g'(t) = \frac{\partial g_{p,q}(t)}{\partial t} = \begin{cases} \frac{(p-q)A \cosh At - pB \cosh Bt - qC \cosh Ct}{pq(p-q)} & \text{if } pq(p-q) \neq 0, \\ \frac{(p+m) \cosh(p+m)t - (p+m) \cosh(p-m)t - 2p(p-m)t \sinh(p-m)t}{-p^2} & \text{if } p \neq 0, q = 0, \\ \frac{(q+m) \cosh(q+m)t - (q+m) \cosh(q-m)t - 2q(q-m)t \sinh(q-m)t}{-q^2} & \text{if } q \neq 0, p = 0, \\ \frac{(2p-m) \cosh(2p-m)t - (2p-m) \cosh mt - 2pmt \sinh mt}{p^2} & \text{if } p = q \neq 0, \\ -4t \sinh mt - 2mt^2 \cosh mt & \text{if } p = q = 0. \end{cases} \quad (3.6)$$

It is obvious that  $\partial g_{p,q}(t)/\partial t$  is continuous with respect to  $(p, q) \in \{(p, q) : p, q \in \mathbb{R}, pq(p-q) \neq 0\}$ . We have also to verify that  $\partial g_{p,q}(t)/\partial t$  is continuous on  $(p, q) \in$

$\{(p, 0) : p \in \mathbb{R}, p \neq 0\}, \{(0, q) : q \in \mathbb{R}, q \neq 0\}, \{(p, p) : p \in \mathbb{R}, p \neq 0\}, \{(0, 0)\}$ .  
In fact, some simple calculations yield

$$\begin{aligned} \lim_{q \rightarrow 0} \frac{\partial g_{p,q}(t)}{\partial t} &= \frac{\partial g_{p,0}(t)}{\partial t}, & \lim_{p \rightarrow 0} \frac{\partial g_{p,q}(t)}{\partial t} &= \frac{\partial g_{0,q}(t)}{\partial t}, \\ \lim_{q \rightarrow p} \frac{\partial g_{p,q}(t)}{\partial t} &= \frac{\partial g_{p,p}(t)}{\partial t}, & \lim_{p \rightarrow 0} \frac{\partial g_{p,p}(t)}{\partial t} &= \frac{\partial g_{0,0}(t)}{\partial t}, \end{aligned}$$

which completes this proof.  $\square$

**Lemma 4.** *We have*

$$\lim_{t \rightarrow 0, t > 0} \frac{3g(t)}{2t^3} = p + q - 3m. \quad (3.7)$$

PROOF. It is easy to check that  $g(0) = g'(0) = g''(0) = 0$ .

In the case of  $pq(p - q) \neq 0$ . Applying L'Hospital's rule (three times) we have

$$\begin{aligned} \lim_{t \rightarrow 0, t > 0} \frac{3g(t)}{2t^3} &= \lim_{t \rightarrow 0, t > 0} \frac{g'(t)}{2t^2} = \dots \\ &= \frac{(p - q)A^3 - pB^3 - qC^3}{4pq(p - q)} = p + q - 3m. \end{aligned} \quad (3.8)$$

In the case of  $pq(p - q) = 0$ . Likewise, some simple calculations also lead to (3.7).

This completes the proof.  $\square$

**Lemma 5.** *Let  $m > 0$  and  $\beta = \max(|A|, |B|, |C|)$  where  $A, B, C$  are defined by (3.2). Then*

(i) *If  $pq(p - q) \neq 0$  and  $p > q$ , then*

$$\lim_{t \rightarrow \infty} \frac{2\beta g(t)}{e^{\beta t}} = \begin{cases} \frac{p + q - m}{pq} & \text{if } p > q > m \text{ or } q < p < 0, \\ \frac{p - q - m}{q(p - q)} & \text{if } p > q = m, \\ -\frac{p - q + m}{p(p - q)} & \text{if } p > 0, q < m, p > q. \end{cases} \quad (3.9)$$

(ii) *If  $p \neq q = 0$ , then*

$$\lim_{t \rightarrow \infty} \frac{2\beta g(t)}{e^{\beta t}} = \begin{cases} -\infty & \text{if } p < 0, \\ -(p + m)p^{-2} & \text{if } p > 0. \end{cases} \quad (3.10)$$

(iii) If  $p = q \neq 0$ , then

$$\lim_{t \rightarrow \infty} \frac{2\beta g(t)}{e^{\beta t}} = \begin{cases} (2p - m)p^{-2} & \text{if } p > m \text{ or } p < 0, \\ -\infty & \text{if } 0 < p \leq m. \end{cases} \quad (3.11)$$

(iv) If  $p = q = 0$ , then

$$\lim_{t \rightarrow \infty} \frac{2\beta g(t)}{e^{\beta t}} = -\infty. \quad (3.12)$$

PROOF. (3.9)–(3.12) easily follow from the following limit relations:

$$\lim_{t \rightarrow \infty} \frac{2 \cosh \alpha t}{e^{\beta t}} = \begin{cases} 1 & \text{if } \beta = |\alpha|, \\ 0 & \text{if } \beta > |\alpha|; \end{cases} \quad (3.13)$$

$$\lim_{t \rightarrow \infty} \frac{2\alpha t \sinh \alpha t}{e^{\beta t}} = \begin{cases} \infty & \text{if } \beta = |\alpha|, \\ 0 & \text{if } \beta > |\alpha|. \end{cases} \quad (3.14)$$

(i) If  $pq(p - q) \neq 0$  and  $p > q$ , then  $\beta = \max(|A|, |B|, |C|) = \max(|A|, |C|)$  because  $|C|^2 - |B|^2 = 4m(p - q) > 0$ . By (3.6) and (3.13) we have

$$\begin{aligned} pq(p - q) \lim_{t \rightarrow \infty} \frac{2\beta g(t)}{e^{\beta t}} &= pq(p - q) \lim_{t \rightarrow \infty} \frac{2g'(t)}{e^{\beta t}} \\ &= \lim_{t \rightarrow \infty} 2 \frac{(p - q)A \cosh At - pB \cosh Bt - qC \cosh Ct}{e^{\beta t}} \\ &= \begin{cases} (p - q)A & \text{if } |A| > |C|, \text{ i.e. } p(q - m) > 0, \\ (p - q)A - qC & \text{if } |A| = |C|, \text{ i.e. } p(q - m) = 0, \\ -qC & \text{if } |A| < |C|, \text{ i.e. } p(q - m) < 0. \end{cases} \end{aligned} \quad (3.15)$$

Taking into account  $pq(p - q) \neq 0$  and  $p > q$ , we obtain

$$pq(p - q) \lim_{t \rightarrow \infty} \frac{2\beta g(t)}{e^{\beta t}} = \begin{cases} (p - q)(p + q - m) & \text{if } p > q > m \text{ or } q < p < 0, \\ p(p - q - m) & \text{if } p > q = m, \\ -q(p - q + m) & \text{if } p > 0, q < m, p > q. \end{cases}$$

Divided by  $pq(p - q)$  in the above limit relation yields (3.9).

(ii) If  $p \neq q = 0$ , then  $\beta = \max(|A|, |B|, |C|) = \max(|p - m|, |p + m|)$ . By (3.6) and (3.13), (3.14) we have

$$\lim_{t \rightarrow \infty} \frac{2\beta g(t)}{e^{\beta t}} = \lim_{t \rightarrow \infty} \frac{2g'(t)}{e^{\beta t}}$$

$$\begin{aligned}
&= \lim_{t \rightarrow \infty} 2 \frac{(p+m) \cosh(p+m)t - (p+m) \cosh(p-m)t - 2p(p-m)t \sinh(p-m)t}{-p^2 e^{\beta t}} \\
&= \begin{cases} -\infty & \text{if } |p-m| > |p+m|, \text{ i.e. } p < 0, \\ -(p+m)p^{-2} > 0 & \text{if } |p-m| < |p+m|, \text{ i.e. } p > 0. \end{cases}
\end{aligned}$$

(iii) If  $p = q \neq 0$ , then  $\beta = \max(|A|, |B|, |C|) = \max(|2p-m|, m)$ . By (3.6) and (3.13), (3.14) we have

$$\begin{aligned}
\lim_{t \rightarrow \infty} \frac{2\beta g(t)}{e^{\beta t}} &= \lim_{t \rightarrow \infty} \frac{2g'(t)}{e^{\beta t}} \\
&= \lim_{t \rightarrow \infty} 2 \frac{(2p-m) \cosh(2p-m)t - (2p-m) \cosh mt - 2pmt \sinh mt}{p^2 e^{\beta t}} \\
&= \begin{cases} (2p-m)p^{-2} & \text{if } |2p-m| > m, \text{ i.e. } p > m \text{ or } p < 0, \\ -\infty & \text{if } |2p-m| = m, \text{ i.e. } p = m, \\ -\infty & \text{if } |2p-m| < m, \text{ i.e. } 0 < p < m. \end{cases}
\end{aligned}$$

(iv) If  $p = q = 0$ , then  $\beta = \max(|A|, |B|, |C|) = \max(m, m, m) = m$ . By (3.6) and (3.13), (3.14) we have

$$\lim_{t \rightarrow \infty} \frac{2\beta g(t)}{e^{\beta t}} = \lim_{t \rightarrow \infty} 2 \frac{-2mt^2 \sinh mt}{e^{mt}} = -\infty.$$

This proof is complete.  $\square$

**Lemma 6.** *Suppose that  $|t_1|, |t_2|, |t_3|$  are pairwise distinct numbers. Then the following identities*

$$\operatorname{sgn}(u(t_1, t_2, t_3)) = \operatorname{sgn}(\cosh t_1 - \cosh t_3) = \operatorname{sgn}(|t_1| - |t_3|) \quad (3.16)$$

hold, where

$$u(t_1, t_2, t_3) = \frac{t_1 \sinh t_1 - t_2 \sinh t_2}{\cosh t_1 - \cosh t_2} - \frac{t_2 \sinh t_2 - t_3 \sinh t_3}{\cosh t_2 - \cosh t_3} \quad (3.17)$$

PROOF. To prove the first identity of (3.16), we note that both the function  $t \rightarrow \cosh t$  and  $t \rightarrow t \sinh t$  are even on  $\mathbb{R}$ , and so we have

$$u(t_1, t_2, t_3) = u(|t_1|, |t_2|, |t_3|).$$

Put  $\cosh |t_i| = x_i$ ,  $i = 1, 2, 3$ , then  $x_1, x_2, x_3 > 1$  and are also pairwise distinct, and

$$|t_i| = \ln \left( x_i + \sqrt{x_i^2 - 1} \right), \quad \sinh |t_i| = \sqrt{x_i^2 - 1}.$$

Thus, the first identity of (3.16) is equivalent to

$$\operatorname{sgn} \left( \frac{f(x_1) - f(x_2)}{x_1 - x_2} - \frac{f(x_2) - f(x_3)}{x_2 - x_3} \right) = \operatorname{sgn}(x_1 - x_3), \quad (3.18)$$

where

$$f(x) = \sqrt{x^2 - 1} \ln(x + \sqrt{x^2 - 1}), \quad x > 1.$$

By simple calculations, we get

$$f'(x) = 1 + \frac{x}{\sqrt{x^2 - 1}} \ln(x + \sqrt{x^2 - 1}),$$

$$f''(x) = \frac{x\sqrt{x^2 - 1} - \ln(x + \sqrt{x^2 - 1})}{(\sqrt{x^2 - 1})^3} := \frac{h(x)}{(\sqrt{x^2 - 1})^3}.$$

Since  $h'(x) = 2(\sqrt{x^2 - 1})^{-1} > 0$ , we have  $h(x) > h(1) = 0$ , which yields  $f''(x) > 0$ , and so  $f$  is convex on  $(1, \infty)$ . From the properties of convex functions it follows that

$$\frac{1}{x_1 - x_3} \left( \frac{f(x_1) - f(x_2)}{x_1 - x_2} - \frac{f(x_2) - f(x_3)}{x_2 - x_3} \right) > 0,$$

which implies that the first identity of (3.16) holds.

Next we show that the second identity of (3.16) holds. Since the function  $t \rightarrow \cosh t$  is even on  $\mathbb{R}$  and strictly increasing on  $\mathbb{R}_+$ , we have

$$\cosh t_1 - \cosh t_3 = (|t_1| - |t_3|) \frac{\cosh |t_1| - \cosh |t_3|}{|t_1| - |t_3|},$$

from which the second identity of (3.16) follows.

This proof is ended. □

**Lemma 7.** *Let*

$$g'(t) = \frac{\partial g_{p,q}(t)}{\partial t} = g_1(t) \cdot g_2(t) \quad \text{for } pq(p-q) \neq 0, \quad (3.19)$$

where

$$g_1(t) = \frac{\cos Bt - \cos Ct}{p - q}, \quad (3.20)$$

$$g_2(t) = \frac{(p - q)A \frac{\cosh At - \cosh Ct}{\cos Bt - \cos Ct} - pB}{pq} \quad (3.21)$$

and  $A, B, C$  are defined by (3.2). Then for all  $t > 0$ , we have

- (i)  $\operatorname{sgn}(g_1(t)) = -\operatorname{sgn}(m)$ .
- (ii)  $\operatorname{sgn}(g_2(t)) = -\operatorname{sgn}(m)\operatorname{sgn}(g'(t))$ .
- (iii)  $g_2(t)$  is monotone with  $t > 0$ .

PROOF. (i) By the second identity of (3.16) we have

$$\operatorname{sgn}(g_1(t)) = \frac{\operatorname{sgn}(|Bt| - |Ct|)}{\operatorname{sgn}(p - q)} = -\operatorname{sgn}(m)$$

for all  $t > 0$ .

(ii) Using (3.19) and the first result of this lemma yield

$$\operatorname{sgn}(g_2(t)) = \frac{\operatorname{sgn}(g'(t))}{\operatorname{sgn}(g_1(t))} = \frac{\operatorname{sgn}(g'(t))}{-\operatorname{sgn}(m)} = -\operatorname{sgn}(m)\operatorname{sgn}(g'(t)).$$

(iii) To prove that  $g_2(t)$  is monotone with  $t > 0$ , it is enough to show that  $\operatorname{sgn}(g_2'(t))$  does not depend on all  $t > 0$ . In fact, we have

$$\operatorname{sgn}(g_2'(t)) = -\operatorname{sgn}(m)\operatorname{sgn}(p - m)\operatorname{sgn}(q - m)\operatorname{sgn}(p + q - m) \quad (3.22)$$

holds for  $pq(p - q) \neq 0$ .

A simple derivative computation yields

$$\begin{aligned} pqg_2'(t) &= (p - q)A \frac{\cosh At - \cosh Ct}{\cos Bt - \cos Ct} \\ &\quad \times \left( \frac{A \sinh At - C \sinh Ct}{\cosh At - \cosh Ct} - \frac{B \sinh Bt - C \sinh Ct}{\cos Bt - \cos Ct} \right) \\ &= t^{-1}(p - q)A \frac{\cosh At - \cosh Ct}{\cos Bt - \cos Ct} u(At, Ct, Bt), \end{aligned}$$

where  $u(t_1, t_2, t_3)$  is defined by (3.17). From (3.16) and  $t > 0$  it follows that

$$\begin{aligned} \operatorname{sgn}(pqg_2'(t)) &= \operatorname{sgn}(t^{-1}(p - q)) \operatorname{sgn}(A) \frac{\operatorname{sgn}(|At| - |Ct|)}{\operatorname{sgn}(|Bt| - |Ct|)} \operatorname{sgn}(\cosh |At| - \cosh |Bt|) \\ &= \operatorname{sgn}(p - q) \operatorname{sgn}(p + q - m) \frac{\operatorname{sgn}(p(q - m))}{\operatorname{sgn}(-m(p - q))} \operatorname{sgn}(q(p - m)) \\ &= -\operatorname{sgn}(m) \operatorname{sgn}(p) \operatorname{sgn}(q) \operatorname{sgn}(p - m) \operatorname{sgn}(q - m) \operatorname{sgn}(p + q - m), \end{aligned}$$

which is equivalent to (3.22) for  $pq(p - q) \neq 0$ .

This accomplishes the proof.  $\square$

#### 4. Proofs of main results

PROOF OF THEOREM 1. Denote by

$$D = \{(p, q) : p + q - 3m \geq 0, \min(p, q) \geq m\} \quad (m > 0).$$

By Lemma 3.1, to prove Theorem 1, it suffices to prove that  $g_{p,q}(t) \geq 0$  for all  $t > 0$  if and only if  $(p, q) \in D$ .

**Necessity.** We prove that  $(p, q) \in D$  is the necessary conditions for  $g(t) = g_{p,q}(t) \geq 0$  for all  $t > 0$ . It is obvious that

$$\lim_{t \rightarrow 0, t > 0} \frac{3g(t)}{2t^3} \geq 0 \quad \text{and} \quad \lim_{t \rightarrow \infty} \frac{2\beta g(t)}{e^{\beta t}} \geq 0. \quad (4.1)$$

The necessary conditions will be obtained from (4.1) together with (3.7) and (3.9)–(3.12). We divide the proof of necessity into six cases.

(i) *Case 1:*  $pq(p - q) \neq 0$  and  $p > q$ .

*Subcase 1:*

$$\begin{cases} p + q - 3m \geq 0, \\ \frac{p + q - m}{pq} \geq 0, \\ p > q > m \text{ or } q < p < 0 \end{cases} \implies \begin{cases} p + q - 3m \geq 0, \\ p > q > m, \end{cases}$$

which implies that  $(p, q) \in \{(p, q) : p > q > m\} = D_{11}$ .

*Subcase 2:*

$$\begin{cases} p + q - 3m \geq 0, \\ \frac{p - q - m}{q(p - q)} \geq 0, \\ p > q = m \end{cases} \implies \begin{cases} p \geq 2m, \\ q = m, \end{cases}$$

which implies that  $(p, q) \in \{(p, q) : p \geq 2m, q = m\} = D_{12}$ .

*Subcase 3:*

$$\begin{cases} p + q - 3m \geq 0, \\ -\frac{p - q + m}{p(p - q)} \geq 0, \\ p > 0, \\ q < m, \\ p > q, \end{cases} \implies \text{which is impossible.}$$

(i') *Case 1'*:  $pq(p-q) \neq 0$  and  $p < q$ .

Since  $g_{p,q}(t)$  is symmetric with respect to  $p$  and  $q$ , so  $(p, q) \in D'_{11} \cup D'_{12}$  if (4.1) holds, where

$$D'_{11} = \{(p, q) : q > p > m\}, \quad D'_{12} = \{(p, q) : q \geq 2m, p = m\}.$$

(ii) *Case 2*:  $p \neq q = 0$ .

*Subcase 1*:

$$\begin{cases} p + q - 3m \geq 0, \\ -\infty \geq 0, \\ p < 0 = q, \end{cases} \implies \text{which is impossible.}$$

*Subcase 2*:

$$\begin{cases} p + q - 3m \geq 0, \\ -(p+m)p^{-2} \geq 0, \\ p > 0 = q, \end{cases} \implies \text{which is impossible.}$$

(ii') *Case 2'*:  $q \neq p = 0$ .

Since  $g_{p,q}(t)$  is symmetric with respect to  $p$  and  $q$ , so this case is also impossible if (4.1) holds.

(iii) *Case 3*:  $p = q \neq 0$ .

*Subcase 1*:

$$\begin{cases} p + q - 3m \geq 0, \\ (2p - m)p^{-2} \geq 0, \\ p > m \text{ or } p < 0, \\ p = q \neq 0 \end{cases} \implies \begin{cases} p + q - 3m \geq 0, \\ p = q > m, \end{cases}$$

which implies that  $(p, q) \in \{(p, q) : p + q - 3m \geq 0, p = q > m\} = D_{31}$ .

*Subcase 2*:

$$\begin{cases} p + q - 3m \geq 0, \\ -\infty \geq 0, \\ 0 < p \leq m, \\ p = q \neq 0, \end{cases} \implies \text{which is impossible.}$$

(iv) *Case 4*:  $p = q = 0$ .

$$\begin{cases} p + q - 3m \geq 0, \\ -\infty \geq 0, \\ p = q = 0, \end{cases} \implies \text{which is impossible.}$$

Summarizing all the cases yield

$$(p, q) \in (D_{11} \cup D_{12}) \cup (D'_{11} \cup D'_{12}) \cup D_{31} = D.$$

**Sufficiency.** We prove the condition  $(p, q) \in D$  is sufficient for  $g(t) = g_{p,q}(t) \geq 0$  for all  $t > 0$ . Since  $g(0) = 0$ , it is enough to prove  $g'(t) \geq 0$  if  $(p, q) \in D$ .

(i) In the case of  $(p, q) \in D$  with  $pq(p - q) \neq 0$ . By (3.8) and (3.15), we see that

$$\operatorname{sgn}(g'(0)) = \operatorname{sgn}(g(0)) \geq 0 \quad \text{and} \quad \operatorname{sgn}(g'(\infty)) = \operatorname{sgn}(g(\infty)) \geq 0$$

if  $(p, q) \in D$  with  $pq(p - q) \neq 0$ .

On the other hand, noting  $m > 0$  and by (ii) and (iii) of Lemma 7, we have

$$\operatorname{sgn}(g_2(0)) = -\operatorname{sgn}(m) \operatorname{sgn}(g'(0)) \leq 0,$$

$$\operatorname{sgn}(g_2(\infty)) = -\operatorname{sgn}(m) \operatorname{sgn}(g'(\infty)) \leq 0$$

and  $g_2(t)$  is monotone with  $t > 0$ , which mean that  $g_2(t) \leq 0$  for all  $t > 0$ . Taking into account  $\operatorname{sgn}(g_1(t)) = -\operatorname{sgn}(m) < 0$ , we obtain that  $g'(t) = g_1(t)g_2(t) \geq 0$  for all  $t > 0$ .

(ii) In the case of  $(p, q) \in D$  with  $pq(p - q) = 0$ . Form Lemma 3 it follows that

$$g'(t) = \frac{\partial g_{p,0}(t)}{\partial t} = \lim_{q \rightarrow 0} \frac{\partial g_{p,q}(t)}{\partial t} \geq 0 \quad \text{if } (p, q) \in D \text{ with } p \neq q = 0.$$

Similarly, we have

$$g'(t) = \frac{\partial g_{0,q}(t)}{\partial t} \geq 0 \quad \text{if } (p, q) \in D \text{ with } q \neq p = 0,$$

$$g'(t) = \frac{\partial g_{p,p}(t)}{\partial t} \geq 0 \quad \text{if } (p, q) \in D \text{ with } p = q \neq 0,$$

$$g'(t) = \frac{\partial g_{0,0}(t)}{\partial t} \geq 0 \quad \text{if } (p, q) \in D \text{ with } p = q = 0.$$

Therefore,  $g'(t) = \partial g_{p,q}(t)/\partial t \geq 0$  if  $(p, q) \in D$ .

This completes the proof of Theorem 1.  $\square$

PROOF OF THEOREM 2. Denote by

$$E = \{(p, q) : p + q - 3m \leq 0, p \geq q, q \leq m\} \quad (m > 0),$$

Then

$$E' = \{(p, q) : p + q - 3m \leq 0, q \geq p, p \leq m\} \quad (m > 0).$$

$$E \cup E' = \{(p, q) : p + q - 3m \leq 0 \text{ and } \min(p, q) \leq m\} \quad (m > 0).$$

By Lemma 3.1, to prove Theorem 2, it suffices to show that  $g_{p,q}(t) \leq 0$  for all  $t > 0$  if and only if  $(p, q) \in E \cup E'$ .

**Necessity.** We prove  $(p, q) \in E \cup E'$  is the necessary conditions for  $g(t) = g_{p,q}(t) \leq 0$  for all  $t > 0$ . It is clear that

$$\lim_{t \rightarrow 0, t > 0} \frac{3g(t)}{2t^3} \leq 0 \quad \text{and} \quad \lim_{t \rightarrow \infty} \frac{2\beta g(t)}{e^{\beta t}} \leq 0. \quad (4.2)$$

We derive the necessary conditions from (4.2) together with (3.7) and (3.9)–(3.12). To this aim, we divide the proof of necessity into six cases.

(i) *Case 1:*  $pq(p - q) \neq 0$  and  $p > q$ .

*Subcase 1:*

$$\begin{cases} p + q - 3m \leq 0, \\ \frac{p + q - m}{pq} \leq 0, \\ p > q > m \text{ or } q < p < 0 \end{cases} \implies 0 > p > q,$$

which implies that  $(p, q) \in \{(p, q) : 0 > p > q\} = E_{11}$ .

*Subcase 2:*

$$\begin{cases} p + q - 3m \leq 0, \\ \frac{p - q - m}{q(p - q)} \leq 0, \\ p > q = m \end{cases} \implies \begin{cases} p \leq 2m, \\ q = m, \\ p > q, \end{cases}$$

which implies that  $(p, q) \in \{(p, q) : q = m, p \leq 2m\} = E_{12}$ .

*Subcase 3:*

$$\begin{cases} p + q - 3m \leq 0, \\ -\frac{p - q + m}{p(p - q)} \leq 0, \\ p > 0, \\ q < m, \\ p > q \end{cases} \implies \begin{cases} p + q - 3m \leq 0, \\ p > 0, \\ q < m, \\ p > q, \end{cases}$$

which implies that  $(p, q) \in \{(p, q) : p + q - 3m \leq 0, p > 0, q < m, p > q\} = E_{13}$ .

(i') *Case 1'*:  $pq(p - q) \neq 0$  and  $p < q$ .

Since  $g_{p,q}(t)$  is symmetric with respect to  $p$  and  $q$ , so  $(p, q) \in E'_{11} \cup E'_{12} \cup E'_{13}$  if (4.2) holds, where

$$E'_{11} = \{(p, q) : 0 > q > p\},$$

$$E'_{12} = \{(p, q) : p = m, q \leq 2m, q > p\},$$

$$E'_{13} = \{(p, q) : p + q - 3m \leq 0, q > 0, p < m, q > p\}.$$

(ii) *Case 2*:  $p \neq q = 0$ .

*Subcase 1*:

$$\begin{cases} p + q - 3m \leq 0, \\ -\infty \leq 0, \\ p < 0 = q \end{cases} \implies \begin{cases} p + q - 3m \leq 0, \\ p < 0 = q, \end{cases}$$

which implies that  $(p, q) \in \{(p, q) : p + q - 3m \leq 0, p < 0 = q\} = E_{21}$ .

*Subcase 2*:

$$\begin{cases} p + q - 3m \leq 0, \\ -(p + m)p^{-2} \leq 0, \\ p > 0 = q \end{cases} \implies \begin{cases} p + q - 3m \leq 0, \\ p > 0 = q, \end{cases}$$

which implies that  $(p, q) \in \{(p, q) : p + q - 3m \leq 0, p > 0 = q\} = E_{22}$ .

(ii') *Case 2'*:  $q \neq p = 0$ .

Since  $g_{p,q}(t)$  is symmetric with respect to  $p$  and  $q$ , so  $(p, q) \in E'_{21} \cup E'_{22}$  if (4.2) holds, where

$$E'_{21} = \{(p, q) : p + q - 3m \leq 0, q < 0 = p\},$$

$$E'_{22} = \{(p, q) : p + q - 3m \leq 0, q > 0 = p\}.$$

(iii) *Case 3*:  $p = q \neq 0$ .

*Subcase 1*:

$$\begin{cases} p + q - 3m \leq 0, \\ (2p - m)p^{-2} \leq 0, \\ p > m \text{ or } p < 0, \\ p = q \neq 0 \end{cases} \implies \begin{cases} p + q - 3m \leq 0, \\ p = q < 0, \end{cases}$$

which implies that  $(p, q) \in \{(p, q) : p + q - 3m \leq 0, p = q < 0\} = E_{31}$ .

Subcase 2:

$$\begin{cases} p + q - 3m \leq 0, \\ -\infty \leq 0, \\ 0 < p \leq m, \\ p = q \neq 0 \end{cases} \implies \begin{cases} p + q - 3m \leq 0, \\ 0 < p = q \leq m, \end{cases}$$

which implies that  $(p, q) \in \{(p, q) : p + q - 3m \leq 0, 0 < p = q \leq m\} = E_{32}$ .

(iv) Case 4:  $p = q = 0$ .

$$\begin{cases} p + q - 3m \leq 0, \\ -\infty \leq 0, \\ p = q = 0, \end{cases} \implies \text{which implies that } (p, q) \in \{(0, 0)\} = E_4.$$

Summarizing all the cases yield

$$\begin{aligned} (p, q) \in & (E_{11} \cup E_{12} \cup E_{13}) \cup (E'_{11} \cup E'_{12} \cup E'_{13}) \\ & \cup (E_{21} \cup E_{22}) \cup (E'_{21} \cup E'_{22}) \cup (E_{31} \cup E_{32}) \cup E_{24} = E \cup E'. \end{aligned}$$

**Sufficiency.** Similarly to proof of sufficiency of Theorem 1, we can prove  $g'(t) \leq 0$  if  $(p, q) \in E \cup E'$ . Hence  $g(t) = g_{p,q}(t) \leq g(0) = 0$  for all  $t > 0$ .

The proof of Theorem 2 is completed.  $\square$

PROOF OF THEOREM 3. Let  $g_{p,q,m}(t) := g_{p,q}(t)$  defined by (3.1) and

$$p' = -p, \quad q' = -q, \quad m' = -m.$$

We easily verify that, for  $p, q, p', q', m, m' \in \mathbb{R}$ ,

$$g_{p,q,m}(t) = -g_{p',q',m'}(t).$$

From this and Lemma 2, for  $m < 0$  Stolarsky mean  $S_{p,q}(a, b)$  is Schur  $m$ -power convex if and only if  $S_{p',q'}(a, b)$  is Schur  $m'$ -power concave with respect to  $(a, b) \in \mathbb{R}_+^2$ , which, by Theorem 2, if and only if

$$p' + q' \leq 3m' \quad \text{and} \quad \min(p', q') \leq m',$$

that is,

$$p + q \geq 3m \quad \text{and} \quad \max(p, q) \geq m.$$

Theorem 3 follows.  $\square$

PROOF OF THEOREM 4. Similarly to the proof of Theorem 3, we have that for  $m < 0$  Stolarsky mean  $S_{p,q}(a,b)$  is Schur  $m$ -power concave if and only if  $S_{p',q'}(a,b)$  is Schur  $m'$ -power convex with respect to  $(a,b) \in \mathbb{R}_+^2$ , which, by Theorem 1, if and only if

$$p' + q' \geq 3m' \quad \text{and} \quad \min(p', q') \geq m',$$

that is,

$$p + q \leq 3m \quad \text{and} \quad \max(p, q) \leq m,$$

The proof of Theorem 4 ends.  $\square$

PROOF OF THEOREM 5. By Lemma 3.1, to prove Theorem 5, it is enough to prove that  $g_{p,q}(t) \geq (\leq) 0$  for all  $t > 0$  if and only if  $p + q \geq (\leq) 0$  for  $m = 0$ . For this end, we divide the proof into four cases.

(i) *Case 1:*  $pq(p - q) \neq 0$ . By (3.1) we have

$$\begin{aligned} g_{p,q}(t) &= \frac{(p - q) \sinh(p + q)t - (p + q) \sinh(p - q)t}{pq(p - q)} \\ &= t(p + q) \frac{k((p + q)t) - k((p - q)t)}{pq}. \end{aligned}$$

Denote by  $k(x) = (\sinh x)/x$  if  $x \neq 0$  and  $k(0) = 1$ . We easily check that  $k(-x) = k(x)$  and  $k'(x) > (<) 0$  for  $x > (<) 0$ . In fact,  $k'(x) = x^{-2}w(x)$ ,  $w(x) = x \cosh x - \sinh x > (<) 0$  for  $x > (<) 0$  because  $w'(x) = x \sinh x > 0$  for  $x \neq 0$ . Thus,

$$\begin{aligned} &\operatorname{sgn} \left( \frac{k((p + q)t) - k((p - q)t)}{pq} \right) \\ &= \operatorname{sgn} \left( \frac{|(p + q)t| - |(p - q)t|}{pq} \right) \operatorname{sgn} \left( \frac{k(|(p + q)t|) - k(|(p - q)t|)}{|(p + q)t| - |(p - q)t|} \right) \\ &= \operatorname{sgn} \left( \frac{t}{|p + q| + |p - q|} \frac{(p + q)^2 - (p - q)^2}{pq} \right) = 1, \end{aligned}$$

it follows that

$$\operatorname{sgn}(g_{p,q}(t)) = \operatorname{sgn}(t(p + q)) \operatorname{sgn} \left( \frac{k((p + q)t) - k((p - q)t)}{pq} \right) = \operatorname{sgn}(p + q).$$

This shows that  $g_{p,q}(t) \geq (\leq) 0$  for all  $t > 0$  if and only if  $p + q \geq (\leq) 0$ .

(ii) *Case 2:*  $pq = 0$ ,  $p \neq q$ . By (3.1) we have

$$g_{p,0}(t) = \frac{2}{p^2}(pt \cosh(pt) - \sinh(pt)) \quad (p \neq 0).$$

Since  $w(x) = x \cosh x - \sinh x > (<)0$  for  $x > (<)0$ ,  $g_{p,0}(t) \geq (\leq)0$  ( $p \neq 0$ ) for all  $t > 0$  if and only if  $pt > (<)0$ , that is,  $p > (<)0$ .

In the same way, we can prove that  $g_{0,q}(t) \geq (\leq)0$  ( $q \neq 0$ ) for all  $t > 0$  if and only if  $q > (<)0$ .

(iii) *Case 3:*  $p = q \neq 0$ . By (3.1) we have

$$g_{p,p}(t) = \frac{\sinh(2pt) - 2pt}{p^2} = \frac{2t}{p} \left( \frac{\sinh(2pt)}{2pt} - 1 \right) = \frac{2t}{p} (k(2pt) - k(0)).$$

Since  $k'(x) > (<)0$  for  $x > (<)0$ , we get  $k(2pt) > k(0)$ . It follows that  $g_{p,p}(t) \geq (\leq)0$  ( $p \neq 0$ ) for all  $t > 0$  if and only if  $2t/p > (<)0$ , that is,  $p > (<)0$ .

(iv) *Case 4:*  $p = q = 0$ . Clearly,  $g_{0,0}(t) = 0$ .

To sum up, for  $m = 0$ ,  $g_{p,q}(t) \geq (\leq)0$  for all  $t > 0$  if and only if  $p + q \geq (\leq)0$ .

The proof of Theorem 5 is completed.  $\square$

## References

- [1] G. D. ANDERSON, M. K. VAMANAMURTHY and M. VUORINEN, Generalized convexity and inequalities, *J. Math. Anal. Appl.* **335**(2) (2007), 1294–1308.
- [2] J. S. AUJLA and F. C. SILVA, Weak majorization inequalities and convex functions, *Linear Algebra Appl.* **369** (2003), 217–233.
- [3] CH.-P. CHEN, The monotonicity of the ratio between Stolarsky means, *RGMA Res. Rep. Coll.* **11**(4) (2008), Art. 15.
- [4] Y.-M. CHU, X.-M. ZHANG and G.-D. WANG, The Schur geometrical convexity of the extended mean values, *J. Convex. Anal.* **15**(4) (2008), 707–718.
- [5] Y. M. CHU and X. M. ZHANG, Necessary and sufficient conditions such that extended mean values are Schur-convex or Schur-concave, *J. Math. Kyoto Univ.* **48**(1) (2008), 229–238.
- [6] Y.-M. CHU and Y.-P. LV, The Schur harmonic convexity of the Hamy symmetric function and its applications, *J. Inequal. Appl.* **2009** (2009), Art. ID 838529.
- [7] G. M. CONSTANTINE, Schur convex functions on the spectra of graphs, *Disc. Math.* **45**(2–3) (1983), 181–188.
- [8] P. CZINDER and ZS. PÁLES, An extension of the Hermite–Hadward inequality and an application for Gini and stolarsky means, *J. Inequal. Pure Appl. Math.* **5**(2) (2004), Art. 42.
- [9] A. FORCINA and A. GIOVAGNOLI, Homogeneity indices and Schur-convex functions, *Statistica* **42**(4) (1982), 529–542.
- [10] CH. GU and H. N. SHI, Schur-convexity and Schur-geometric convexity of Lehmer Means, *Math. Prac. Theory* **39**(12) (2009), 183–188.
- [11] G. H. HARDY, J. E. LITTLEWOOD and G. PÓLYA, Some simple inequalities satisfied by convex functions, *Messenger Math.* **58** (1929), 145–152.
- [12] PETER. A. HÄSTÖ, A monotonicity property of ratios of symmetric homogeneous means, *J. Inequal. Pure Appl. Math.* **35** (2002), Art. 71.

- [13] F. K. HWANG, U. G. ROTHBLUM and L. SHEPP, Monotone optimal multipartitions using Schur convexity with respect to partial orders, *SIAM J. Discrete Math.* **6**(4) (1993), 533–547.
- [14] F. K. HWANG and U. G. ROTHBLUM, Partition-optimization with Schur convex sum objective functions, *SIAM J. Discrete Math.* **18**(3) (2004), 512–524.
- [15] E. B. LEACH and M. C. SHOLANDER, Extended mean values, *Amer. Math. Monthly* **85** (1978), 84–90.
- [16] E. B. LEACH and M. C. SHOLANDER, Extended mean values II, *J. Math. Anal. Appl.* **92** (1983), 207–223.
- [17] D.-M. LI and H.-N. SHI, Schur convexity and Schur-geometrically concavity of generalized exponent mean, *J. Math. Inequal.* **3**(2) (2009), 217–225.
- [18] ZH. LIU, Minkowski's inequality for extended mean values, Proceedings of the Second ISAAC Congress, Vol. 1 (Fukuoka, 1999), 585–592, *Int. Soc. Anal. Appl. Comput.* **7**, Kluwer Acad. Publ., Dordrecht, 2000.
- [19] L. LOSONCZI and ZS. PÁLES, Minkowski's inequality for two variable difference means, *Proc. Amer. Math. Soc.* **126** (1998), 779–789.
- [20] L. LOSONCZI, Ratio of Stolarsky means: monotonicity and comparison, *Publ. Math. Debrecen* **75** (2009), 221–238.
- [21] A. W. MARSHALL and I. OLKIN, Inequalities: Theory of Majorization and Its Applications, *Academic Press, New York*, 1979.
- [22] M. MERKLE, Convexity, Schur-convexity and bounds for the gamma function involving the digamma function, *Rocky Mountain J. Math.* **28**(3) (1998), 1053–1066.
- [23] E. NEUMAN and J. SÁNDOR, Inequalities involving Stolarsky and Gini means, *Math. Pannon.* **14**(1) (2003), 29–44.
- [24] E. NEUMAN and ZS. PÁLES, On comparison of Stolarsky and Gini means, *J. Math. Anal. Appl.* **278**(2) (2003), 274–284.
- [25] ZS. PÁLES, Inequalities for differences of powers, *J. Math. Anal. Appl.* **131** (1988), 271–281.
- [26] ZS. PÁLES, Comparison of two variable homogeneous means, *General Inequal. 6. Proc. 6th Internat. Conf. Math. Res. Inst. Oberwolfach, Birkhäuser Verlag Basel* (1992), 59–69.
- [27] F. QI, Logarithmically convexities of the Extended Mean values, *Proc. Amer. Math. Soc.* **130**(6) (2002), 1787–1796.
- [28] F. QI, The extended mean values: definition, properties, monotonicities, comparison, convexities, generalizations, and applications, *Cubo Mat. Educ.* **5**(3) (2003), 63–90.
- [29] F. QI, J. SÁNDOR and S. S. DRAGOMIR, Notes on the Schur-convexity of the extended mean values, *Taiwanese J. Math.* **9**(3) (2005), 411–420.
- [30] F. QI, A note on Schur-convexity of extended mean values, *Rocky Mountain J. Math.* **35**(5) (2005), 1787–1793.
- [31] J. SÁNDOR, The Schur-convexity of Stolarsky and Gini means, *Banach J. Math. Anal.* **1**(2) (2007), 212–215.
- [32] M. SHAKED, J. G. SHANTHIKUMAR and Y. L. TONG, Parametric Schur convexity and arrangement monotonicity properties of partial sums, *J. Multivariate Anal.* **53**(2) (1995), 293–310.
- [33] H. N. SHI, S. H. WU and F. QI, An alternative note on the Schur-convexity of the extended mean values, *Math. Inequal. Appl.* **9**(2) (2006), 219–224.
- [34] H.-N. SHI, Y.-M. JIANG and W.-D. JIANG, Schur-convexity and Schur-geometrically concavity of Gini means, *Comp. Math. Appl.* **57**(2) (2009), 266–274.

- [35] C. STEPNIAK, Stochastic ordering and Schur-convex functions in comparison of linear experiments, *Metrika* **36**(5) (1989), 291–298.
- [36] K. B. STOLARSKY, Generalizations of the Logarithmic Mean, *Math. Mag.* **48** (1975), 87–92.
- [37] K. B. STOLARSKY, The power and generalized Logarithmic Means, *Amer. Math. Monthly* **87** (1980), 545–548.
- [38] B.-Y. WANG, Foundations of Majorization Inequalities, *Beijing Normal Univ. Press, Beijing, China*, 1990 (in *Chinese*).
- [39] Z.-H. WANG and X.-M. ZHANG, Necessary and sufficient conditions for Schur convexity and Schur-geometrically convexity of Gini means, *Communications of inequalities researching* **14**(2) (2007), 193–197 (in *Chinese*).
- [40] Z.-H. WANG, The necessary and sufficient condition for  $S$ -convexity and  $S$ -geometrically convexity of Gini mean, *J. Beijing Ins. Edu. (Natural Science)* **2**(5) (2007), 1–3.
- [41] W.-F. XIA, The Schur harmonic convexity of Lehmer means, *Int. Math. Forum* **4**(41) (2009), 2009–2015.
- [42] ZH.-H. YANG, ON the homogeneous functions with two parameters and its monotonicity, *J. Inequal. Pure Appl. Math.* **6**(4) (2005), Art. 101.
- [43] ZH.-H. YANG, ON the log-convexity of two-parameter homogeneous functions, *Math. Inequal. Appl.* **10**(3) (2007), 499–516.
- [44] ZH. -H. YANG, On the monotonicity and log-convexity of a four-parameter homogeneous mean, *J. Inequal. Appl.* **2008** (2008), Art. ID 149286.
- [45] ZH.-H. YANG, Necessary and sufficient conditions for Schur convexity the two-parameter symmetric homogeneous means, *Applied Mathematical Sciences* **5**(64) (2011), 3183–3190.
- [46] ZH.-H. YANG, Some monotonicity results for the ratio of two-parameter symmetric homogeneous functions, *Int. J. Math. Math. Sci.* **2009** (2009), Art. ID 591382.
- [47] ZH.-H. YANG, Log-convexity of ratio of the two-parameter symmetric homogeneous functions and an application, *J. Inequal. Spec. Funct.* **1**(1) (2010), 16–29.
- [48] ZH.-H. YANG, Necessary and sufficient conditions for Schur geometrical convexity of the four-parameter homogeneous means, *Abstr. Appl. Anal.* **2010** (2010), Art. ID 830163.
- [49] X.-M. ZHANG, Schur-convex functions and isoperimetric inequalities, *Proc. Amer. Math. Soc.* **126**(2) (1998), 461–470.
- [50] X.-M. ZHANG, Geometrically Convex Functions, *Hefei, Anhui University Press*, 2004 (in *Chinese*).

ZHEN-HANG YANG  
 SYSTEM DIVISION  
 ZHEJIANG PROVINCE ELECTRIC POWER TEST  
 AND RESEARCH INSTITUTE  
 HANGZHOU, ZHEJIANG  
 CHINA, 310014

*E-mail:* yzhkm@163.com

*(Received January 11, 2010; revised February 7, 2011)*