

Inclusion relations of hyperbolic type metric balls II

By RIKU KLÉN (Turku) and MATTI VUORINEN (Turku)

Abstract. Inclusion relations of metric balls defined by the hyperbolic, the quasihyperbolic, the j -metric and the chordal metric will be studied. The hyperbolic metric, the quasihyperbolic metric and the j -metric are considered in the unit ball.

1. Introduction

The most important metrics in the classical complex analysis are the Euclidean and the hyperbolic metric. Studying quasiconformal mappings in \mathbb{R}^n , F. W. GEHRING and B. P. PALKA [6] introduced the quasihyperbolic metric, which plays the role of the hyperbolic metric in the higher dimensions. The quasihyperbolic metric has recently been studied in [2], [11]. There are also other hyperbolic type metrics like the distance ratio metric and the Apollonian metric, which have lately been studied by various authors [3], [4].

Suppose that (X, d_j) , $j = 1, 2$, are two metric spaces with $X \subset \mathbb{R}^n$ such that both metrics determine the same Euclidean topology. In order to understand the geometric structure of the spaces, it is a fundamental question to study the corresponding balls and inclusion relations among balls with the same center with respect to both of these two metrics. In several classical cases such relations are well-known, comparing e.g. the Euclidean balls and hyperbolic balls (cf. [14, Section 2]). But this comparison problem makes sense in numerous non-classical cases as well, as pointed out in [15]. Very recently, such non-classical inclusion problems have been studied in [8], [9] and our goal here is to investigate the inclusion problem for hyperbolic type metrics in the unit ball.

Mathematics Subject Classification: 30F45, 51M10.

Key words and phrases: hyperbolic ball, j -metric ball, quasihyperbolic ball, chordal ball.

In a metric space (X, m) we define *metric ball* or *m-ball* with center $x \in X$ and radius $r > 0$ by

$$B_m(x, r) = \{y \in X : m(x, y) < r\} \quad (1.1)$$

and *metric sphere* with center $x \in X$ and radius $r > 0$ by

$$S_m(x, r) = \{y \in X : m(x, y) = r\}.$$

For Euclidean balls and spheres we use notation $B^n(x, r)$ and $S^{n-1}(x, r)$. For $x, y \in \mathbb{R}^n$ we use notation $l = [x, y]$ for the line segment joining x to y , similarly $[x, y) = l \setminus \{y\}$ and $(x, y] = l \setminus \{x\}$. For $x \in G \subsetneq \mathbb{R}^n$ we denote by $d(x)$ the Euclidean distance between x and ∂G .

The n -dimensional unit ball will be denoted by \mathbb{B}^n and half-space by $\mathbb{H}^n = \{x \in \mathbb{R}^n : x_n > 0\}$. The *hyperbolic length* of a rectifiable curve $\gamma \subset \mathbb{B}^n$ is defined by

$$\ell_{\rho_{\mathbb{B}^n}}(\gamma) = \int_{\gamma} \frac{2|dz|}{1-|z|^2}$$

and $\gamma \subset \mathbb{H}^n$ by

$$\ell_{\rho_{\mathbb{H}^n}}(\gamma) = \int_{\gamma} \frac{|dz|}{z_n}.$$

The *hyperbolic metric* in $G \in \{\mathbb{B}^n, \mathbb{H}^n\}$ is

$$\rho_G(x, y) = \inf_{\gamma} \ell_{\rho_G}(\gamma),$$

where the infimum is taken over all rectifiable curves in G joining x and y .

For a domain $G \subsetneq \mathbb{R}^n$, $n \geq 2$ the *quasihyperbolic length* of a rectifiable arc $\gamma \subset G$ is given by

$$\ell_k(\gamma) = \int_{\gamma} \frac{|dz|}{d(z)}$$

and the *quasihyperbolic metric* by

$$k_G(x, y) = \inf_{\gamma} \ell_k(\gamma), \quad (1.2)$$

where the infimum is taken over all rectifiable curves in G joining x and y . Note that $k_{\mathbb{H}^n} = \rho_{\mathbb{H}^n}$.

The *distance ratio metric* or *j-metric* in a proper subdomain G of the Euclidean space \mathbb{R}^n , $n \geq 2$, is defined by

$$j_G(x, y) = \log \left(1 + \frac{|x - y|}{\min\{d(x), d(y)\}} \right).$$

The distance ratio metric satisfies the triangle inequality by [12, Lemma 2.2]. If the domain G is understood from the context we use the notation j instead of j_G and k instead of k_G . The distance ratio metric was first introduced by F. W. GEHRING and B. G. OSGOOD [5], and in the above form by M. VUORINEN [14]. The metric space (G, j_G) is not geodesic for any domain G [7, Theorem 2.10].

The *chordal metric* in $\overline{\mathbb{R}^n} = \mathbb{R}^n \cup \{\infty\}$ is defined by

$$q(x, y) = \begin{cases} \frac{|x - y|}{\sqrt{1 + |x|^2} \sqrt{1 + |y|^2}}, & x \neq \infty \neq y, \\ \frac{1}{\sqrt{1 + |x|^2}}, & y = \infty. \end{cases}$$

The metric space $(\overline{\mathbb{R}^n}, q)$ is not geodesic

We find radii $m(r)$ and $M(r)$ for $d_1, d_2 \in \{q, k_G, j_G\}$ such that

$$B_{d_1}(x, m(r)) \subset B_{d_2}(x, r) \subset B_{d_1}(x, M(r))$$

for all $x \in G$. This kind of estimates can be used to compare metrics, because $B_{m_2}(x, r) \subset B_{m_1}(x, r)$ for all r implies $m_1(x, y) \leq m_2(x, y) < r$ for all $y \in B_{m_2}(x, r)$.

The geometry of the j -metric balls is easily described in \mathbb{H}^n , $\mathbb{R}^n \setminus \{0\}$ and polygons in the case $n = 2$, as we will point out in Lemma 2.1, Remark 2.2 and Figure 1. However, when the boundary of the domain does not consist of lines and isolated points the situation becomes more complicated. Already in the unit ball the geometry of the j -metric balls differ significantly from the other cases.

The following theorem is our main result.

Theorem 1.3. *Let $G = \mathbb{B}^n$, $x \in G$ and $r > 0$. Then*

$$\begin{aligned} B_j(x, m_1(r)) &\subset B_\rho(x, r), & B_j(x, m_2(r)) &\subset B_k(x, r), \\ B_j(x, m_3(r)) &\subset B_q(x, r), & r &< (1 - |x|)/\sqrt{2(1 + |x|^2)}, \end{aligned}$$

where

$$\begin{aligned} m_1(r) &= \log \left(1 + 2 \sinh \frac{r}{2} \right), \\ m_2(r) &= \log \left(1 + 2 \sinh \frac{r}{4} \right), \\ m_3(r) &= \log \left(1 + \frac{r}{\sqrt{1 - r^2}} \right). \end{aligned}$$

Remark 1.4. We show that

$$\frac{r}{2} < m_1(r) < r, \quad \frac{r}{4} < m_2(r) < \frac{r}{2}, \quad \frac{4r}{5} < m_3(r),$$

where $m_1(r)$, $m_2(r)$ and $m_3(r)$ are as in Theorem 1.3.

By a simple computation we obtain

$$m_1'(r) = \frac{\cosh \frac{r}{2}}{1 + 2 \sinh \frac{r}{2}} < 1,$$

where the inequality follows from the fact $\cosh(r/2) < 1 + 2 \sinh(r/2)$. Since $m_1(0) = 0$ we obtain $m_1(r) < r$.

The lower bound for $m_1(r)$ follows from

$$m_1(r) = \log(1 + e^{r/2} - e^{-r/2}) > \log(e^{r/2}) = \frac{r}{2}.$$

The upper and lower bounds for $m_2(r)$ follow from the bounds of $m_1(r)$, because $m_2(r) = m_1(r/2)$.

Since

$$m_3''(r) = \frac{1 - 3r(r + \sqrt{1 - r^2})}{(1 - r^2)^2(r + \sqrt{1 - r^2})^2}$$

we know that $m_3'(r)$ attains its minimum on the interval $(0, 1)$ at $r_0 = ((5 - \sqrt{17})/3)^{1/2}/2$ and

$$m_3'(r_0) = \frac{24\sqrt{3}}{(7 + \sqrt{17})(\sqrt{5 - \sqrt{17}} + \sqrt{7 + \sqrt{17}})} > \frac{4}{5}.$$

Thus $4r/5 < m_3$.

2. Preliminary results

In this section we introduce preliminary results such as properties of hyperbolic type metric balls and relations between hyperbolic type metrics.

The *curvature* of a plane curve parameterized in polar coordinates is defined by

$$\kappa = \frac{r(\theta)^2 + 2r'(\theta)^2 - r(\theta)r''(\theta)}{(r(\theta)^2 + r'(\theta)^2)^{3/2}}.$$

Note that if a curve has a constant curvature, then it is a circular arc.

The following lemma describes the geometric shape of the j -metric balls. Some examples of j -metric disks are shown in Figure 1.

Lemma 2.1. *Let $G \subset \mathbb{R}^2$, $x \in G$ and $r > 0$. Then curvature of $S_j(x, r)$*

(1) *in $G = \mathbb{R}^2 \setminus \{0\}$ is*

$$\kappa = \begin{cases} \frac{1}{|x|(e^r - 1)}, & \text{in } \mathbb{R}^2 \setminus B^2(0, x), \\ \frac{e^r |2 - e^r|}{|x|(e^r - 1)}, & \text{in } B^2(0, x). \end{cases}$$

(2) *in $G = \mathbb{H}^2$ with $x_1 = 0$ is*

$$\kappa = \begin{cases} \frac{1}{(e^r - 1)x_2}, & \text{in } \{y \in \mathbb{H}^2 : y_2 > x_2\}, \\ \frac{|x|^2}{(e^r - 1)(|x|^2 + t^2)}, & \text{in } \{y \in \mathbb{H}^2 : y_2 < x_2\}, \end{cases}$$

where $|t| < |x|(e^r - 1)$.

(3) *in $G = \mathbb{B}^2$ is*

$$\kappa = \begin{cases} \frac{1}{(e^r - 1)(1 - |x|)}, & \text{in } B^2(0, |x|), \\ \frac{f(\alpha)^2 + 2f'(\alpha) - f(\alpha)f''(\alpha)}{(f(\alpha)^2 + f'(\alpha)^2)^{3/2}}, & \text{in } \mathbb{B}^2 \setminus \overline{B^2(0, |x|)}, \end{cases}$$

where α is the angle $\sphericalangle(x, 0, y)$ for $y \in S_j(x, r)$ and

$$f(\alpha) = \frac{(1 - e^r)^2 - \beta - \sqrt{1 + (e^{2r} - 2e^r)(1 + |x|^2) - 2(e^r - 1)^2\beta + \beta^2}}{e^r(e^r - 2)}$$

for $\beta = |x| \cos \alpha$ and $\alpha \in [0, \gamma]$, where

$$\gamma = \begin{cases} \pi, & r \geq \log((1 + |x|)/(1 - |x|)), \\ 2 \arcsin \frac{(e^r - 1)(1 - |x|)}{2|x|}, & r < \log((1 + |x|)/(1 - |x|)). \end{cases}$$

(4) *in $\{y \in G : d(y) \geq d(x)\}$ for any domain G is*

$$\kappa = \frac{1}{(e^r - 1)d(x)}.$$

PROOF. Let $G \subset \mathbb{R}^2$, $x \in G$ and $r > 0$.

- (1) By definition of the j -metric the j -sphere consists of two circular arcs, or in the case $r = \log 2$, a circular arc and a line segment. The assertion follows from proof of [7, Theorem 3.1].
- (2) The case $S_j(x, r) \cap \{y \in \mathbb{H}^2 : y_2 > x_2\}$ is similar to (2). Therefore, we consider $S = S_j(x, r) \cap \{y \in \mathbb{H}^2 : y_2 < x_2\}$. By the definition of the j -metric we obtain that $S = \{y \in \mathbb{H}^2 : y = (t, f(t)), t \in (-|x|(e^r - 1), |x|(e^r - 1))\}$ for the function

$$f(t) = \left| \frac{\sqrt{|x|^2 + e^r(e^r - 2)(|x|^2 + t^2)} - |x|}{e^r(e^r - 2)} \right|.$$

By a straightforward computation we obtain that the curvature of S at point $(t, f(t))$, $t \in (-|x|(e^r - 1), |x|(e^r - 1))$, is

$$\kappa(t) = \frac{|f''(t)|}{(1 + f'(t)^2)^{3/2}} = \frac{|x|^2}{(e^r - 1)(|x|^2 + t^2)^{3/2}}.$$

- (3) To simplify notation we may assume $x = (x_1, 0)$ for $x_1 \in [0, 1)$. We divide $S_j(x, r)$ into two cases $S_1 = S_j(x, r) \cap B^2(0, |x|)$ and $S_2 = S_j(x, r) \cap (\mathbb{B}^2 \setminus \overline{B^2(0, |x|)})$. The set S_2 is always nonempty, whereas the set $S_1 = \emptyset$ whenever $r > \log((1 + |x|)/(1 - |x|))$.

Let us first consider S_1 . By the definition of the j -metric for all $y \in S_1$ we have $|x - y| = (e^r - 1)(1 + |x|^2)$, and thus

$$\kappa(t) = \frac{1}{(e^r - 1)(1 - |x|)}.$$

Let us finally consider S_2 . The assertion follows by the definition of curvature, if the function $f(\alpha) = |y|$, for the point $y \in S_2$ and $\angle(x, 0, y) = \alpha$. By the definition of the j -metric for $y \in S_2$ we obtain $|x - y| = (e^r - 1)(1 - |y|)$ and by the law of cosines we obtain

$$(e^r - 1)^2(1 - |y|)^2 = |x|^2 + |y|^2 - 2|x||y|\cos \alpha,$$

which is equivalent to $|y| = f(\alpha)$. The sign in $f(\alpha)$ was chosen to be minus, because otherwise the values would have been greater than or equal to 1.

- (4) The assertion follows from the definition of the j -metric as in the case (3). □

Remark 2.2. (1) By Lemma 2.1 the boundary $\partial B_j(x, r)$ consists of two circular arcs in the case $G = \mathbb{R}^2 \setminus \{0\}$ and a circular arc and a part of a conic section (hyperbola if $r > \log 2$, parabola if $r = \log 2$ or ellipse if $r < \log 2$) in the case $G = \mathbb{H}^2$, see Figure 1. In the case $G = \mathbb{B}^2$ the boundary $\partial B_j(x, r)$ is more complicated as it may not even contain circular arc.

(2) By Lemma 2.1 (2) the boundary $\partial B_j(x, r)$ can be formed in a polygonal domain $P \subset \mathbb{R}^2$. First the medial axis of P needs to be formed. In a convex polygon the medial axis consists of line segments and can be found as Voronoi diagram [10]. If the polygon is not convex, then the medial axis can contain parts of conic sections. However, the medial axis is unique and it divides P into smaller domains A_i . In each A_i the boundary $\partial B_j(x, r)$ consists of circular arcs, when $A_i \cap (\partial B_j(x, r)) \subset \{z \in P : d(z) \geq d(x)\}$, and parts of a conic section similarly as above in the case $G = \mathbb{H}^2$, see Figure 1.




Figure 1. Examples of j -disks $B_j(x, r)$ in punctured plane (left), half-plane (middle) and in a rectangle (right). Gray line is the medial axis and dashed gray line is the set $\{z \in G : d(z) = d(x)\}$.

For the sake of easy reference we recapitulate a few basic facts in the next result. For part (1) and (4), see e.g. [1, Section 7], for (2) and (3) [14, Section 2].

Proposition 2.3. For all $x, y \in \mathbb{B}^n$

(1)

$$\rho_{\mathbb{B}^n}(x, y) \leq 2k_{\mathbb{B}^n}(x, y) \leq 2\rho_{\mathbb{B}^n}(x, y),$$

(2)

$$\rho_{\mathbb{B}^n}(x, y) = 2 \operatorname{arcsinh} \frac{|x - y|}{\sqrt{1 - |x|^2} \sqrt{1 - |y|^2}},$$

(3) and for $r > 0$

$$B_{\rho_{\mathbb{B}^n}}(x, r) = B^n \left(\frac{x(1 - t^2)}{1 - |x|^2 t^2}, \frac{(1 - |x|^2)t}{1 - |x|^2 t^2} \right),$$

where $t = \tanh(r/2)$,

(4) and for $r \in (0, 1/\sqrt{1+|x|^2})$

$$B_q(x, r) = B^n \left(\frac{x}{1 - r^2(1 + |x|^2)}, \frac{r(1 + |x|^2)\sqrt{1 - r^2}}{1 - r^2(1 + |x|^2)} \right).$$

Note that in (4) we have $B_q(x, r) \subset \mathbb{B}^n$, if $r \in (0, (1 - |x|)/\sqrt{2(1 + |x|^2)})$.

Lemma 2.4. *Let $G = \mathbb{B}^n$, $x = te_1$, $t \in [0, 1)$, $r > 0$ and*

$$\partial B_j(x, r) \cap \{z \in \mathbb{R}^n : z = se_1, s \in \mathbb{R}\} = \{y_1, y_2\}$$

with $|y_2| \leq |y_1|$. Then

$$B^n(x, |x - y_1|) \subset B_j(x, r) \subset B^n(x, |x - y_2|).$$

PROOF. By the selection of y_1 and y_2 we have $j(x, y_1) = j(x, y_2) = r$.

We show that $B^n(x, |x - y_1|) \subset B_j(x, r)$. Let $y \in B^n(x, |x - y_1|)$. Because $d(y_1) < \min\{d(x), d(y)\}$ we have

$$\begin{aligned} j(x, y) &= \log \left(1 + \frac{|x - y|}{\min\{d(x), d(y)\}} \right) < \log \left(1 + \frac{|x - y_1|}{\min\{d(x), d(y)\}} \right) \\ &\leq \log \left(1 + \frac{|x - y_1|}{d(y_1)} \right) = j(x, y_1) = r. \end{aligned}$$

We show that $B_j(x, r) \subset B^n(x, |x - y_2|)$. Let $z \in B_j(x, r)$. We divide the proof into two cases: $y_2 \in [-x, x)$ and $y_2 \in (-x/|x|, -x)$.

If $y_2 \in [-x, x)$, then $d(x) \leq d(y_2)$ and thus $j(x, z) < j(x, y_2)$ is equivalent to

$$\frac{|x - z|}{\min\{d(x), d(z)\}} < \frac{|x - y_2|}{d(x)}$$

implying $|x - z| < |x - y_2|$.

If $y_2 \in (-x/|x|, -x)$, then $d(y_2) < d(x)$. Inequality $j(x, z) < j(x, y_2)$ is equivalent to

$$\frac{|x - z|}{\min\{d(x), d(z)\}} < \frac{|x - y_2|}{d(y_2)}$$

implying $|x - z| < |x - y_2|$, if additionally $d(z) \leq d(y_2)$. If $d(z) > d(y_2)$, then immediately $|x - z| < |x - y|$.

In both cases we obtain that $|x - z| > |x - y_2|$ and thus the assertion follows. \square

Lemma 2.5. *Let $G \in \{\mathbb{R}^n \setminus \{0\}, \mathbb{H}^n, \mathbb{B}^n\}$, $r > 0$ and $x \in G$. Then*

$$B = B^n \left(\frac{y+z}{2}, \frac{|y-z|}{2} \right) \subset B_j(x, r),$$

where $y, z \in l \cap \partial B_j(x, r)$ with $d(y) \leq d(z)$ and l is the line that contains x and a boundary point of G that is closest to x . Moreover, B is the largest Euclidean ball contained in $B_j(x, r)$.

PROOF. The cases $G \in \{\mathbb{R}^n \setminus \{0\}, \mathbb{B}^n\}$ follow easily from Lemma 2.1 (1) and (2).

Let us consider $G = \mathbb{H}^n$. Now $y = x(1 - e^{-r}(1 - |x|)/|x|)$ and

$$z = \begin{cases} x \frac{1 - e^r(1 - |x|)}{|x|}, & \text{if } r \leq \log \frac{1}{1 - |x|}, \\ x \frac{e^r(1 - |x|) - 1}{|x|}, & \text{if } r > \log \frac{1}{1 - |x|} \text{ and } \log \frac{1 + |x|}{1 - |x|} \geq r, \\ x \frac{1 - e^{-r}(1 + |x|)}{|x|}, & \text{if } r > \log \frac{1}{1 - |x|} \text{ and } \log \frac{1 + |x|}{1 - |x|} < r. \end{cases}$$

Thus

$$\frac{y+z}{2} = \begin{cases} \frac{x(1 - (1 - |x|) \cosh r)}{|x|}, & r \leq \log \frac{1 + |x|}{1 - |x|}, \\ -\frac{xe^{-r}}{|x|}, & r > \log \frac{1 + |x|}{1 - |x|} \end{cases}$$

and

$$\frac{|y-z|}{2} = \begin{cases} (1 - |x|) \sinh r, & r \leq \log \frac{1 + |x|}{1 - |x|}, \\ 1 - e^{-r}, & r > \log \frac{1 + |x|}{1 - |x|}. \end{cases} \quad \square$$

Remark 2.6. Lemma 2.5 is not true in general. For $G = \mathbb{R}^2 \setminus \{-e_1, e_1\}$ does not hold for $x = e_2$ and $r = \log 2$. In this case $\partial B_j(x, r)$ consists of two perpendicular line segments and a circular arc. The line segments are $[0, a(e_1 + e_2)]$ and $[0, a(-e_1 + e_2)]$, where $a = (1 + \sqrt{3})/2$. See Figure 2.

However, the following question is open: Is Lemma 2.5 true in convex domains?

Lemma 2.7. *For $a, b \in [0, 1]$ we have*

$$(1) \min\{1 - a, 1 - b\}(1 + \max\{a, b\}) \leq \sqrt{1 - a^2} \sqrt{1 - b^2}.$$




Figure 2. Boundary $\partial B_j(e_2, \log 2)$ in $G = \mathbb{R}^2 \setminus \{-e_1, e_1\}$.

For $x, y \in \mathbb{B}^n$ and $r \geq \operatorname{arcsinh}(2|x|/(1 - |x|^2))$ we have

- (2) $\min\{d(x), d(y)\}(1 + |x|) \leq \sqrt{1 - |x|^2}\sqrt{1 - |y|^2}$,
(3) $\frac{2|x|}{1 - |x|} - \frac{1 + |x|}{1/\tanh(r/2) - |x|} \leq (1 + |x|)\frac{e^r - 1}{2}$.

PROOF. We consider first (1). We easily obtain

$$\min\{1 - a, 1 - b\}(1 + \max\{a, b\}) = 1 - \max\{a, b\}^2 \leq \sqrt{1 - a^2}\sqrt{1 - b^2}.$$

Part (2) follows from (1).

Let us then consider (3), which is equivalent to showing that the function

$$f(r) = (1 + |x|)\frac{e^r - 1}{2} - \frac{2|x|}{1 - |x|} + \frac{1 + |x|}{1/\tanh(r/2) - |x|}$$

is nonnegative. Since

$$f'(r) = \frac{1 + |x|}{2} \left(e^r + \frac{1}{(\cosh(r/2) - |x|\sinh(r/2))^2} \right) > 0$$

and $f(\operatorname{arcsinh}(2|x|/(1 - |x|^2))) = |x|^2/(1 - |x|) \geq 0$ the assertion follows. \square

3. Inclusion relations of metric balls

In this section we consider metric balls in unit ball $G = \mathbb{B}^n$. Since we do not know the exact form of the quasihyperbolic ball we need to use the hyperbolic balls.

Theorem 3.1. *Let $G = \mathbb{B}^n$, $x \in G$ and $r > 0$. Then*

$$B_j(x, m) \subset B_\rho(x, r) \subset B_j(x, M),$$

where $m = \max\{m_1, m_2\}$

$$m_1 = \log \left(1 + (1 + |x|) \sinh \frac{r}{2} \right), \quad m_2 = \log \left(1 + (1 - |x|) \frac{e^r - 1}{2} \right)$$

and

$$M = \log \left(1 + (1 + |x|) \frac{e^r - 1}{2} \right).$$

Moreover, the inclusions are sharp and $M/m \rightarrow 1$ as $r \rightarrow 0$.

PROOF. We prove inclusion $B_j(x, m) \subset B_\rho(x, r)$. Let us first assume $y \in B_j(x, m_1)$, which is equivalent to

$$|x - y| < \min\{d(x), d(y)\}(1 + |x|) \sinh(r/2). \quad (3.2)$$

Since \sinh and $\operatorname{arcsinh}$ are increasing we obtain by Proposition 2.3 (2) and (3.2)

$$\begin{aligned} \rho(x, y) &\leq 2 \operatorname{arcsinh} \frac{\min\{d(x), d(y)\}(1 + |x|) \sinh(r/2)}{\sqrt{1 - |x|^2} \sqrt{1 - |y|^2}} \\ &\leq 2 \operatorname{arcsinh}(\sinh(r/2)) \leq r, \end{aligned}$$

where the second inequality follows from Lemma 2.7 (2). Now $y \in B_\rho(x, r)$ and thus $B_j(x, m_1) \subset B_\rho(x, r)$.

Let us then assume $y \in \partial B_j(x, m_2)$ and $m_1 < m_2$. Since $m_1 < m_2$ is equivalent to $r > 4 \operatorname{arctanh} |x|$, we obtain by Lemma 2.3 (2) that $S^{n-1}(0, |x|) \subset B_\rho(x, r)$. Thus $|x| < |y|$, and $j(x, y) = m_2$ is equivalent to

$$\frac{|x - y|}{1 - |y|} = (1 - |x|) \frac{e^r - 1}{2}.$$

Now

$$\begin{aligned} \rho(x, y) &= 2 \operatorname{arcsinh} \left(\frac{|x - y|}{(1 - |y|) \sqrt{1 - |x|^2}} \sqrt{\frac{1 - |y|}{1 + |y|}} \right) \\ &= 2 \operatorname{arcsinh} \left(\frac{(1 - |x|)(e^r - 1)}{2 \sqrt{1 - |x|^2}} \sqrt{\frac{1 - |y|}{1 + |y|}} \right) \end{aligned}$$

and $\rho(x, y) \leq \rho(x, z_1)$ for $z_1 \in \partial B_j(x, m_2)$ with $|z_1| \leq |y|$. In other words $z_1 = l \cap \partial B_j(x, r)$, where $l = \{u \in \mathbb{B}^n : u = sx, s < |x|\}$, and

$$|z_1| = 1 - \frac{2(1 + |x|)}{1 + |x| + e^r(1 - |x|)}.$$

Thus

$$\rho(x, y) \leq \rho(x, z_1) = 2 \operatorname{arcsinh} \frac{|x| + |z_1|}{\sqrt{1 - |x|^2} \sqrt{1 - |z_1|^2}}$$

and $y \in B_\rho(x, r)$ implying $B_j(x, m_2) \subset B_\rho(x, r)$.

We show that m is sharp. If $S^{n-1}(0, |x|) \cap (\partial B_j(x, r)) = \emptyset$, then $j(x, z_1) = m_2 = m$. Otherwise we can choose $z \in S^{n-1}(0, |x|) \cap (\partial B_j(x, r))$ and we obtain $j(x, z) = m_1 = m$.

We prove next the inclusion $B_\rho(x, r) \subset B_j(x, M)$. We assume first that $y \in B_\rho(x, r)$ and $d(x) \leq d(y)$, which is equivalent to $|y| \leq |x|$. By Lemma 2.1 (3) and Proposition 2.3 (3) $y \in B_j(x, M)$, if $j(x, z_1) \leq M$ for $z_1 = l \cap \partial B_\rho(x, r)$, where $l = \{u \in \mathbb{B}^n : u = sx, s < |x|\}$. If $|x - z_1| = |x| - |z_1|$, then by Proposition 2.3 (2) we have $r \leq \operatorname{arcsinh}(2|x|/(1 - |x|^2))$,

$$|z_1| = \frac{2|x| + (|x|^2 - 1) \sinh r}{1 + |x|^2 - (|x|^2 - 1) \cosh r}$$

and

$$\frac{|x - z_1|}{\min\{d(x), d(z_1)\}} = \frac{|x| - |z_1|}{1 - |x|} = \frac{1 + |x|}{1/\tanh(r/2) - |x|} \leq (1 + |x|) \frac{e^r - 1}{2}$$

implying $j(x, z_1) \leq M$. If $|x - z_1| = |x| + |z_1|$, then by Proposition 2.3 (2) we have $r \geq \operatorname{arcsinh}(2|x|/(1 - |x|^2))$,

$$|z_1| = \frac{2|x| + (|x|^2 - 1) \sinh r}{-1 - |x|^2 + (|x|^2 - 1) \cosh r}$$

and by Lemma 2.7 (3)

$$\frac{|x - z_1|}{\min\{d(x), d(z_1)\}} = \frac{|x| + |z_1|}{1 - |x|} = \frac{2|x|}{1 - |x|} - \frac{1 + |x|}{1/\tanh(r/2) - |x|} \leq (1 + |x|) \frac{e^r - 1}{2}$$

implying $j(x, z_1) \leq M$.

We assume then that $y \in \partial B_\rho(x, r)$ and $d(y) \leq d(x)$. Now $y \in \partial B_\rho(x, r)$ is equivalent to

$$\frac{|x - y|}{\sqrt{1 - |y|}} = \sqrt{1 + |y|} \sqrt{1 - |x|^2} \sinh \frac{r}{2}$$

and thus by Lemma 2.3 (3)

$$j(x, y) = \log \left(1 + \frac{\sqrt{1+|y|}}{\sqrt{1-|y|}} \sqrt{1-|x|^2} \sinh \frac{r}{2} \right) \leq j(x, z_2)$$

for $z_2 = l \cap \partial B_\rho(x, r)$, where $l = \{u \in \mathbb{B}^n : u = sx, s > |x|\}$. By Proposition 2.3 (2) we obtain

$$|z_2| = 1 - \frac{2(1-|x|)}{1-|x| + e^r(1+|x|)}$$

and

$$j(x, z_2) = \log \left(1 + \frac{|z_2| - |x|}{1 - |z_2|} \right) = M$$

implying the claim. This also shows that M is sharp.

By the l'Hôpital rule we obtain

$$\lim_{r \rightarrow 0} \frac{M}{m} = \lim_{r \rightarrow 0} \frac{(1 + (e^r - 1)(1 + |x|)/2) \cosh(r/2)}{e^r(1 + (1 + |x|) \sinh(r/2))} = 1$$

and the assertion follows. \square




Figure 3. An example of inclusions of hyperbolic disks (black) and j -metric disks (gray) in the unit disk. The black dot is the center of the disks and the black thin circle is the unit circle.

Corollary 3.3. *Let $G = \mathbb{B}^n$, $x \in G$ and $r > 0$. Then*

$$B_\rho(x, m) \subset B_j(x, r) \subset B_\rho(x, M),$$

where

$$m = \log \left(1 + \frac{2(e^r - 1)}{1 + |x|} \right)$$

and

$$M = \min \left\{ 2 \operatorname{arcsinh} \frac{e^r - 1}{1 + |x|}, \log \frac{2e^r - 1 - |x|}{1 - |x|} \right\}.$$

Moreover, the inclusions are sharp and $M/m \rightarrow 1$ as $r \rightarrow 0$.

PROOF. Assertion follows from Theorem 3.1. \square

Corollary 3.4. *Let $G = \mathbb{B}^n$, $x \in G$ and $r > 0$. Then*

$$B_j(x, m) \subset B_k(x, r) \subset B_j(x, M),$$

where

$$m = \max \left\{ \log \left(1 + (1 + |x|) \sinh \frac{r}{4} \right), \log \left(1 + (1 - |x|) \frac{e^{r/2} - 1}{2} \right) \right\}$$

and

$$M = \log \left(1 + (1 + |x|) \frac{e^r - 1}{2} \right).$$

PROOF. Assertion follows from Theorem 3.1 and Proposition 2.3 (1). \square

Corollary 3.5. *Let $G = \mathbb{B}^n$, $x \in G$ and $r > 0$. Then*

$$B_k(x, m) \subset B_j(x, r) \subset B_k(x, M),$$

where

$$m = \log \left(1 + \frac{2(e^r - 1)}{1 + |x|} \right)$$

and

$$M = \min \left\{ 4 \operatorname{arcsinh} \frac{e^r - 1}{1 + |x|}, 2 \log \frac{2e^r - 1 - |x|}{1 - |x|} \right\}.$$

PROOF. Assertion follows from Corollary 3.4. \square

It is easy to verify that for $x \in \mathbb{B}^n$ we have $B_q(x, r) \subset \mathbb{B}^n$ if and only if $r < (1 - |x|)/\sqrt{2(1 + |x|^2)}$.

Theorem 3.6. *Let $G = \mathbb{B}^n$, $x \in G$ and $r \in (0, r_0)$ for $r_0 = (1 - |x|)/\sqrt{2(1 + |x|^2)}$. We define real numbers $r_1 = |x|/\sqrt{1 + |x|^2}$, $r_2 =$*

$2|x|/(1+|x|^2)$ and intervals $I_1 = [0, \min\{r_0, r_1\})$, $I_2 = [\min\{r_0, r_1\}, \min\{r_0, r_2\})$ for $|x| < \sqrt{2} - 1$, and $I_3 = [r_2, r_0)$ for $|x| < 2 - \sqrt{3}$. Then

$$B_j(x, m) \subset B_q(x, r) \subset B_j(x, M),$$

where

$$M = \log \frac{(1-|x|)(1-r^2(1+|x|^2))}{1-|x|-r(1+|x|^2)(r+\sqrt{1-r^2})}$$

and

$$m = \min\{m_1, m_2\}$$

for

$$m_1 = \begin{cases} \log \left(1 + \frac{r(1+|x|^2)(\sqrt{1-r^2}-r|x|)}{(1-|x|)(1-r^2(1+|x|^2))} \right), & r \in I_1 \text{ or} \\ & |x| < \sqrt{2} - 1 \text{ and } r \in I_2, \\ \infty, & \text{otherwise,} \end{cases}$$

$$m_2 = \begin{cases} \log \frac{(1+|x|)(1-r^2(1+|x|^2))}{1+|x|-r(r+\sqrt{1-r^2})(1+|x|^2)}, & |x| < 2 - \sqrt{3} \text{ and } r \in I_3, \\ \infty, & \text{otherwise.} \end{cases}$$

Moreover, the inclusions are sharp and $M/m \rightarrow 1$ as $r \rightarrow 0$.

PROOF. Because of symmetry of G we may assume $x = te_1$ for $t \in [0, 1)$. Since $\partial B_q(x, r)$ intersects the line $l = \{z \in \mathbb{R}^n : z = e_1 s, s \in (-\infty, \infty)\}$ twice we denote $(\partial B_q(x, r)) \cap l = \{y_1, y_2\}$. We assume that $y_1 \in (x, e_1)$ and $y_2 \in (x, -e_1)$.

We prove first that $B_q(x, r) \subset B_j(x, M)$. Our idea is to show that

$$B_q(x, r) \subset B^n(x, |x - y_1|) \subset B_j(x, M). \quad (3.7)$$

The first inclusion follows from Proposition 2.3 (4) and the observation that $|x| \leq |x|/(1-r^2(1+|x|^2))$.

The second inclusion of (3.7) follows from Lemma 2.4, because $q(x, y_1) = r$ is equivalent to

$$|y_1| = \frac{|x| + r\sqrt{1-r^2}(1+|x|^2)}{1-r^2(1+|x|^2)}$$

and thus

$$j(x, y_1) = \log \left(1 + \frac{|y_1| - |x|}{1 - |y_1|} \right) = \log \frac{(1-|x|)(1-r^2(1+|x|^2))}{1-|x|-r(1+|x|^2)(r+\sqrt{1-r^2})} = M.$$

We prove next that $B_j(x, m) \subset B_q(x, r)$. Our idea is to show that

$$B_j(x, m) \subset B^n(x, |x - y_2|) \subset B_q(x, r), \quad (3.8)$$

where the second inclusion follows from Proposition 2.3 (4) and the observation that $|x| \leq |x|/(1 - r^2(1 + |x|^2))$.

The first inequality of (3.8) follows from Lemma 2.4, if $j(x, y_2) = m$. To show this we consider three cases: $y_2 \in [0, x)$, $y_2 \in (0, -x]$ and $y_2 \in (-x, -x/|x|)$.

In the case $y_2 \in [0, x)$, $q(x, y_2) = r$ is equivalent to

$$|y_2| = \frac{|x| + r\sqrt{1 - r^2}(1 + |x|^2)}{1 - r^2(1 + |x|^2)}$$

and thus

$$j(x, y_2) = \log \left(1 + \frac{|x| - |y_2|}{1 - |x|} \right) = \log \left(1 + \frac{r(1 + |x|^2)(\sqrt{1 - r^2} - r|x|)}{(1 - |x|)(1 - r^2(1 + |x|^2))} \right) = m_1.$$

In the case $y_2 \in [0, -x)$, $q(x, y_2) = r$ is equivalent to

$$|y_2| = \frac{|x| - r\sqrt{1 - r^2}(1 + |x|^2)}{1 - r^2(1 + |x|^2)} \quad (3.9)$$

and thus

$$j(x, y_2) = \log \left(1 + \frac{|x| + |y_2|}{1 - |x|} \right) = \log \left(1 + \frac{r(1 + |x|^2)(\sqrt{1 - r^2} - r|x|)}{(1 - |x|)(1 - r^2(1 + |x|^2))} \right) = m_1.$$

In the case $y_2 \in (-x, -x/|x|)$, $q(x, y_2) = r$ is equivalent to (3.9) and thus

$$j(x, y_2) = \log \left(1 + \frac{|x| + |y_2|}{1 - |y_2|} \right) = \log \frac{(1 + |x|)(1 - r^2(1 + |x|^2))}{1 + |x| - r(r + \sqrt{1 - r^2})(1 + |x|^2)} = m_2.$$

Sharpness of m and M follow from (3.7), (3.8) and the selection of y_1 and y_2 .

We finally show that $M/m \rightarrow 1$ as $r \rightarrow 0$. By the l'Hôpital's rule we obtain

$$\lim_{r \rightarrow 0} \frac{M}{m} = \lim_{r \rightarrow 0} \frac{M}{m_1} = \lim_{r \rightarrow 0} \frac{(1 + \alpha - \beta)(1 - |x| + \gamma)}{(1 - \alpha - \beta)(1 - |x| - \gamma)} = 1,$$

where $\alpha = 2r\sqrt{1 - r^2}|x|$, $\beta = r^2(1 - |x|^2)$ and $\gamma = r(\sqrt{1 - r^2} - r)(1 + |x|^2)$. \square




Figure 4. An example of inclusions of chordal disks (black) and j -metric disks (gray) in the unit disk. The black dot is the center of the disks and the black thin circle is the unit circle.

Corollary 3.10. *Let $G = \mathbb{B}^n$, $x \in G$ and $r > 0$. Then*

$$B_q(x, m) \subset B_j(x, r) \subset B_q(x, M),$$

where

$$m = \frac{(1 - e^{-r})(1 - |x|)}{\sqrt{1 + |x|^2} \sqrt{1 + (e^{-r}(1 - |x|) - 1)^2}}$$

and

$$M = \begin{cases} \frac{(e^r - 1)(1 - |x|)}{\sqrt{1 + |x|^2} \sqrt{1 + (e^r(1 - |x|) - 1)^2}}, & r \leq \log \frac{1 + |x|}{1 - |x|}, \\ \frac{(e^r - 1)(1 + |x|)}{e^r \sqrt{1 + |x|^2} \sqrt{1 + (e^{-r}(1 - |x|) - 1)^2}}, & r > \log \frac{1 + |x|}{1 - |x|}. \end{cases}$$

Moreover, the inclusions are sharp and $M/m \rightarrow 1$ as $r \rightarrow 0$.

Remark 3.11. In Corollary 3.10, we have $B_q(x, M) \subset \mathbb{B}^n$ if $M \leq (1 - |x|)/\sqrt{2(1 + |x|^2)}$, which is equivalent to

$$r \leq \log \frac{2(1 + |x|)}{1 + 2|x| - |x|^2}.$$

PROOF OF THEOREM 1.3. The radii m_1 and m_2 follow from Theorem 3.1 and Corollary 3.4.

The radius m_3 follows from Theorem 3.6. \square

Theorem 3.12. *Let $G = \mathbb{B}^n$, $x \in G$ and $r \in (0, r_0)$ for $r_0 = (1 - |x|)/\sqrt{2(1 + |x|^2)}$. Then*

$$B_\rho(x, m) \subset B_q(x, r) \subset B_\rho(x, M),$$

where

$$m = 2 \operatorname{arcsinh} \frac{r(\sqrt{1-r^2} - r|x|)(1 + |x|^2)}{\sqrt{1-|x|^2}a\sqrt{1-a^{-2}(|x| - r\sqrt{1-r^2})(1 + |x|^2)^2}}$$

and

$$M = 2 \operatorname{arcsinh} \frac{r(\sqrt{1-r^2} + r|x|)(1 + |x|^2)}{\sqrt{1-|x|^2}a\sqrt{1-a^{-2}(|x| + r\sqrt{1-r^2})(1 + |x|^2)^2}}$$

for $a = 1 - r^2(1 + |x|^2)$.

Moreover, the inclusions are sharp and $M/m \rightarrow 1$ as $r \rightarrow 0$.

PROOF. We prove the first inclusion $B_\rho(x, m) \subset B_q(x, r)$. Let $y \in \partial B_\rho(x, m)$ with $|y| \leq |z|$ for all $z \in \partial B_\rho(x, m)$. By Lemma 2.3 (3) and (4)

$$B_\rho(x, m) \subset B^n(x, |x - y|) \subset B_q(x, r).$$

Since $q(x, y) = r$ is equivalent to

$$y = \frac{x}{|x|} \frac{|x| - r\sqrt{1-r^2}(1 + |x|^2)}{1 - r^2(1 + |x|^2)}$$

we obtain $\rho(x, y) = m$. The radius m is sharp by the selection of y .

We prove next the inclusion $B_q(x, r) \subset B_\rho(x, M)$. Let $y \in \partial B_q(x, r)$ with $|y| \geq |z|$ for all $z \in \partial B_q(x, r)$. By Lemma 2.3 (3) and (4)

$$B_q(x, r) \subset B^n(x, |x - y|) \subset B_\rho(x, M).$$

Since $q(x, y) = r$ is equivalent to

$$|y| = \frac{|x| + r\sqrt{1-r^2}(1 + |x|^2)}{1 - r^2(1 + |x|^2)}$$

we obtain $\rho(x, y) = M$. The radius M is sharp by the selection of y .

Clearly $M/m \rightarrow 1$ as $r \rightarrow 0$ and the assertion follows. \square




Figure 5. An example of inclusions of hyperbolic disks (gray) and chordal disks (black) in the unit disk. The black dot is the center of the disks and the black thin circle is the unit circle.

Corollary 3.13. Let $G = \mathbb{B}^n$, $x \in G$ and $r \in (0, r_0)$ for $r_0 = (1 - |x|)/\sqrt{2(1 + |x|^2)}$. Then

$$B_k(x, m) \subset B_q(x, r) \subset B_k(x, M),$$

where

$$m = 2 \operatorname{arcsinh} \frac{r(\sqrt{1-r^2} - r|x|)(1 + |x|^2)}{\sqrt{1-|x|^2}a\sqrt{1-a^{-2}(|x| - r\sqrt{1-r^2}(1 + |x|^2))^2}}$$

and

$$M = 4 \operatorname{arcsinh} \frac{r(\sqrt{1-r^2} + r|x|)(1 + |x|^2)}{\sqrt{1-|x|^2}a\sqrt{1-a^{-2}(|x| - r\sqrt{1-r^2}(1 + |x|^2))^2}}$$

for $a = 1 - r^2(1 + |x|^2)$.

Theorem 3.14. Let $G = \mathbb{B}^n$, $x \in G$ and $r > 0$. Then

$$B_q(x, m) \subset B_\rho(x, r) \subset B_q(x, M),$$

where

$$m = \frac{(1 - |x|^2) \sinh \frac{r}{2}}{\sqrt{1 + |x|^2} \sqrt{(1 + |x|^2) \cosh r + 2|x| \sinh r}}$$

and

$$M = \frac{(1 - |x|^2) \sinh \frac{r}{2}}{\sqrt{1 + |x|^2} \sqrt{(1 + |x|^2) \cosh r - 2|x| \sinh r}}.$$

Moreover, the inclusions are sharp and $M/m \rightarrow 1$ as $r \rightarrow 0$.

PROOF. We prove first the inclusion $B_q(x, m) \subset B_\rho(x, r)$. Let $y \in \partial B_q(x, m)$ with $|y| \geq |z|$ for all $z \in \partial B_q(x, m)$. By Lemma 2.3 (3) and (4)

$$B_q(x, m) \subset B^n(x, |x - y|) \subset B_\rho(x, r).$$

Since $q(x, y) = r$ is equivalent to

$$|y| = \frac{2|x| + (1 - |x|^2) \sinh r}{1 + |x|^2 + (1 - |x|^2) \cosh r}$$

we obtain $q(x, y) = m$. The radius m is sharp by the selection of y .

We prove next the inclusion $B_\rho(x, r) \subset B_q(x, M)$. Let $y \in \partial B_\rho(x, r)$ with $|y| \geq |z|$ for all $z \in \partial B_\rho(x, r)$. By Lemma 2.3 (3) and (4)

$$B_\rho(x, r) \subset B^n(x, |x - y|) \subset B_q(x, M).$$

Since $\rho(x, y) = r$ is equivalent to

$$y = \frac{x}{|x|} \frac{2|x| - (1 - |x|^2) \sinh r}{1 + |x|^2 + (1 - |x|^2) \cosh r}$$

we obtain $q(x, y) = M$. The radius M is sharp by the selection of y .

Clearly $M/m \rightarrow 1$ as $r \rightarrow 0$ and the assertion follows. \square

Corollary 3.15. *Let $G = \mathbb{B}^n$, $x \in G$ and $r > 0$. Then*

$$B_q(x, m) \subset B_k(x, r) \subset B_q(x, M),$$

where

$$m = \frac{(1 - |x|^2) \sinh \frac{r}{4}}{\sqrt{1 + |x|^2} \sqrt{(1 + |x|^2) \cosh \frac{r}{2} + 2|x| \sinh \frac{r}{2}}}$$

and

$$M = \frac{(1 - |x|^2) \sinh \frac{r}{2}}{\sqrt{1 + |x|^2} \sqrt{(1 + |x|^2) \cosh r - 2|x| \sinh r}}.$$

Note that in Theorem 3.14 and Corollary 3.15, $B_q(x, M) \subset \mathbb{B}^n$ if $M \leq (1 - |x|)/\sqrt{2(1 + |x|^2)}$.

ACKNOWLEDGEMENTS. The research of MATTI VUORINEN was supported by the Academy of Finland, Project 2600066611.

References

- [1] G. D. ANDERSON, M. K. VAMANAMURTHY and M. VUORINEN, Conformal Invariants, Inequalities and Quasiconformal Maps, *John Wiley & Sons, New York*, 1997.
- [2] M. CHEN, X. CHEN and T. QIAN, Quasihyperbolic distance in punctured planes, *Complex Anal. Oper. Theory*, 2012, doi 10.1007/s11785-012-0275-1.
- [3] P. HÄSTÖ, Z. IBRAGIMOV, D. MINDA, S. PONNUSAMY and S. SAHOO, Isometries of some hyperbolic-type path metrics, and the hyperbolic medial axis, In the tradition of Ahlfors—Bers. IV, Vol. 432, (by Dick Canary, Jane Gilman, Juha Heinonen and Howard Masur, eds.), *Contemp. Math., Amer Math. Soc., Providence, RI*, 2007, 63–74.
- [4] M. HUANG, S. PONNUSAMY, X. WANG and S. K. SAHOO, The Apollonian inner metric and uniform domains, *Math. Nachr.* **283**, no. 9 (2010), 1277–1290.
- [5] F. W. GEHRING and B. G. OSGOOD, Uniform domains and the quasi-hyperbolic metric, *J. Anal. Math.* **36** (1979), 50–74.
- [6] F. W. GEHRING and B. P. PALKA, Quasiconformally homogeneous domains, *J. Anal. Math.* **30** (1976), 172–199.
- [7] R. KLÉN, Local convexity properties of j -metric balls, *Ann. Acad. Sci. Fenn. Math* **33** (2008), 281–293.
- [8] R. KLÉN and M. VUORINEN, Inclusion relations of hyperbolic type metric balls, *Publ. Math. Debrecen* **81** (2012), 289–311.
- [9] V. MANOJLOVIĆ, On conformally invariant extremal problems, *Appl. Anal. Discrete Math.* **3** (2009), 97–119. .
- [10] A. OKABE, B. BOOTS and K. SUGIHARA, Spatial tessellations: Concepts and applications of Voronoi diagrams, *Wiley*, 2000.
- [11] A. RASILA and J. TALPONEN, Convexity properties of quasihyperbolic balls on Banach spaces, *Ann. Acad. Sci. Fenn. Math.* **37**, no. 1 (2012), 215–228.
- [12] P. SEITTENRANTA, Möbius-invariant metrics, *Math. Proc. Cambridge Philos. Soc.* **125** (1999), 511–533.
- [13] M. VUORINEN, Conformal invariants and quasiregular mappings, *J. Anal. Math.* **45** (1985), 69–115.
- [14] M. VUORINEN, Conformal geometry and quasiregular mappings, Vol. 1319, Lecture Notes in Math., *Springer-Verlag, Berlin – Heidelberg*, 1988.

42 R. Klén and M. Vuorinen : Inclusion relations of hyperbolic type metric...

- [15] M. VUORINEN, Metrics and quasiregular mappings, Proc. Int. Workshop on Quasiconformal Mappings and their Applications, IIT Madras, Dec 27, 2005–Jan 1, 2006, (by S. Ponnusamy, T. Sugawa and M. Vuorinen, eds.), *Quasiconformal Mappings and their Applications*, Narosa Publishing House, New Delhi, India, 2007, 291–325.

RIKU KLÉN
DEPARTMENT OF MATHEMATICS
AND STATISTICS
UNIVERSITY OF TURKU
FI-20014
FINLAND

E-mail: riku.klen@utu.fi

MATTI VUORINEN
DEPARTMENT OF MATHEMATICS
AND STATISTICS
UNIVERSITY OF TURKU
FI-20014
FINLAND

E-mail: vuorinen@utu.fi

(Received October 14, 2011; revised September 3, 2012)