

On monotonicity of some combinatorial sequences

By QING-HU HOU (Tianjin), ZHI-WEI SUN (Nanjing) and HAOMIN WEN (Philadelphia)

Abstract. We confirm Sun's conjecture that $(\sqrt[n+1]{F_{n+1}}/\sqrt[n]{F_n})_{n \geq 4}$ is strictly decreasing to the limit 1, where $(F_n)_{n \geq 0}$ is the Fibonacci sequence. We also prove that the sequence $(\sqrt[n+1]{D_{n+1}}/\sqrt[n]{D_n})_{n \geq 3}$ is strictly decreasing with limit 1, where D_n is the n -th derangement number. For m -th order harmonic numbers $H_n^{(m)} = \sum_{k=1}^n 1/k^m$ ($n = 1, 2, 3, \dots$), we show that $(\sqrt[n+1]{H_{n+1}^{(m)}}/\sqrt[n]{H_n^{(m)}})_{n \geq 3}$ is strictly increasing.

1. Introduction

A challenging conjecture of Firoozbakht states that

$$\sqrt[n]{p_n} > \sqrt[n+1]{p_{n+1}} \quad \text{for every } n = 1, 2, 3, \dots,$$

where p_n denotes the n -th prime. Note that $\lim_{n \rightarrow \infty} \sqrt[n]{p_n} = 1$ by the Prime Number Theorem. In [4] the second author conjectured further that for any integer $n > 4$ we have the inequality

$$\frac{\sqrt[n+1]{p_{n+1}}}{\sqrt[n]{p_n}} < 1 - \frac{\log \log n}{2n^2},$$

which has been verified for all $n \leq 3.5 \times 10^6$. Motivated by this and [3], Sun [4, Conjecture 2.12] conjectured that the sequence $(\sqrt[n+1]{S_{n+1}}/\sqrt[n]{S_n})_{n \geq 7}$ is strictly

Mathematics Subject Classification: Primary: 05A10; Secondary: 11B39, 11B75.

Key words and phrases: Combinatorial sequences, monotonicity, log-concavity.

The first author and the second author are supported by the National Natural Science Foundation of China with grants 11171167 and 11171140 respectively. The second author is the corresponding author.

increasing, where S_n is the sum of the first n positive squarefree numbers. Moreover, he also posed many conjectures on monotonicity of sequences of the type $(\sqrt[n+1]{a_{n+1}}/\sqrt[n]{a_n})_{n \geq N}$ with $(a_n)_{n \geq 1}$ a familiar combinatorial sequence of positive integers.

Throughout this paper, we set $\mathbb{N} = \{0, 1, 2, \dots\}$ and $\mathbb{Z}^+ = \{1, 2, 3, \dots\}$.

Let A and B be integers with $\Delta = A^2 - 4B \neq 0$. The Lucas sequence $u_n = u_n(A, B)$ ($n \in \mathbb{N}$) is defined as follows:

$$u_0 = 0, \quad u_1 = 1, \quad \text{and} \quad u_{n+1} = Au_n - Bu_{n-1} \text{ for } n = 1, 2, 3, \dots$$

It is well known that $u_n = (\alpha^n - \beta^n)/(\alpha - \beta)$ for all $n \in \mathbb{N}$, where

$$\alpha = \frac{A + \sqrt{\Delta}}{2} \quad \text{and} \quad \beta = \frac{A - \sqrt{\Delta}}{2}$$

are the two roots of the characteristic equation $x^2 - Ax + B = 0$. The sequence $F_n = u_n(1, -1)$ ($n \in \mathbb{N}$) is the famous Fibonacci sequence, see [1, p. 46] for combinatorial interpretations of Fibonacci numbers.

Our first result is as follows.

Theorem 1.1. *Let $A > 0$ and $B \neq 0$ be integers with $\Delta = A^2 - 4B > 0$, and set $u_n = u_n(A, B)$ for $n \in \mathbb{N}$. Then there exists an integer $N > 0$ such that the sequence $(\sqrt[n+1]{u_{n+1}}/\sqrt[n]{u_n})_{n \geq N}$ is strictly decreasing with limit 1. In the case $A = 1$ and $B = -1$ we may take $N = 4$.*

Remark 1.1. Under the condition of Theorem 1.1, by [2, Lemma 4] we have $u_n < u_{n+1}$ unless $A = n = 1$. Note that the second assertion in Theorem 1.1 confirms a conjecture of the second author [4, Conjecture 3.1] on the Fibonacci sequence.

For $n \in \mathbb{Z}^+$ the n -th derangement number D_n denotes the number of permutations σ of $\{1, \dots, n\}$ with $\sigma(i) = i$ for no $i = 1, \dots, n$. It has the following explicit expression (cf. [1, p. 67]):

$$D_n = \sum_{k=0}^n (-1)^k \frac{n!}{k!}.$$

Our second theorem is the following result conjectured by the second author [4, Conjecture 3.3].

Theorem 1.2. *The sequence $(\sqrt[n+1]{D_{n+1}}/\sqrt[n]{D_n})_{n \geq 3}$ is strictly decreasing with limit 1.*

Remark 1.2. It follows from Theorem 1.2 that the sequence $(\sqrt[n]{D_n})_{n \geq 2}$ is strictly increasing.

For each $m \in \mathbb{Z}^+$ those $H_n^{(m)} = \sum_{k=1}^n 1/k^m$ ($n \in \mathbb{Z}^+$) are called harmonic numbers of order m . The usual harmonic numbers (of order 1) are those rational numbers $H_n = H_n^{(1)}$ ($n = 1, 2, 3, \dots$).

Our following theorem confirms Conjecture 2.16 of Sun [4].

Theorem 1.3. *For any $m \in \mathbb{Z}^+$, the sequence $(\sqrt[n+1]{H_{n+1}^{(m)}} / \sqrt[n]{H_n^{(m)}})_{n \geq 3}$ is strictly increasing.*

We will prove Theorems 1.1–1.3 in Sections 2–4 respectively. It seems that there is no simple form for the generating function $\sum_{n=0}^{\infty} \sqrt[n]{a_n} x^n$ with $a_n = u_n, D_n, H_n^{(m)}$. Note also that the set of those sequences $(a_n)_{n \geq 1}$ of positive numbers with $(\sqrt[n+1]{a_{n+1}} / \sqrt[n]{a_n})_{n \geq 1}$ decreasing (or increasing) is closed under multiplication.

2. Proof of Theorem 1.1

PROOF OF THEOREM 1.1. Set

$$\alpha = \frac{A + \sqrt{\Delta}}{2}, \quad \beta = \frac{A - \sqrt{\Delta}}{2}, \quad \text{and} \quad \gamma = \frac{\beta}{\alpha} = \frac{A - \sqrt{\Delta}}{A + \sqrt{\Delta}}.$$

Then

$$\log u_n = \log \frac{\alpha^n(1 - \gamma^n)}{\alpha - \beta} = n \log \alpha + \log(1 - \gamma^n) - \log \sqrt{\Delta}$$

for any $n \in \mathbb{Z}^+$. Note that

$$\log \frac{\sqrt[n+1]{u_{n+1}}}{\sqrt[n]{u_n}} = \frac{\log u_{n+1}}{n+1} - \frac{\log u_n}{n} = \frac{\log(1 - \gamma^{n+1})}{n+1} - \frac{\log(1 - \gamma^n)}{n} + \frac{\log \sqrt{\Delta}}{n(n+1)}.$$

Since

$$\lim_{n \rightarrow \infty} \frac{\log(1 - \gamma^n)}{n} = \lim_{n \rightarrow \infty} \frac{-\gamma^n}{n} = 0 \quad \text{and} \quad \lim_{n \rightarrow \infty} \frac{1}{n(n+1)} = 0,$$

we deduce that

$$\lim_{n \rightarrow \infty} \log \frac{\sqrt[n+1]{u_{n+1}}}{\sqrt[n]{u_n}} = 0, \quad \text{i.e.,} \quad \lim_{n \rightarrow \infty} \frac{\sqrt[n+1]{u_{n+1}}}{\sqrt[n]{u_n}} = 1.$$

For any $n \in \mathbb{Z}^+$, clearly

$$\begin{aligned} \frac{\sqrt[n+1]{u_{n+1}}}{\sqrt[n]{u_n}} > \frac{\sqrt[n+2]{u_{n+2}}}{\sqrt[n+1]{u_{n+1}}} &\iff \frac{\log u_{n+1}}{n+1} - \frac{\log u_n}{n} > \frac{\log u_{n+2}}{n+2} - \frac{\log u_{n+1}}{n+1} \\ &\iff \Delta_n := \frac{2 \log u_{n+1}}{n+1} - \frac{\log u_n}{n} - \frac{\log u_{n+2}}{n+2} > 0. \end{aligned}$$

Observe that

$$\begin{aligned} \Delta_n &= 2 \log \alpha + \frac{2 \log(1 - \gamma^{n+1})}{n+1} - \frac{2 \log \sqrt{\Delta}}{n+1} \\ &\quad - \left(2 \log \alpha + \frac{\log(1 - \gamma^n)}{n} + \frac{\log(1 - \gamma^{n+2})}{n+2} - \frac{\log \sqrt{\Delta}}{n} - \frac{\log \sqrt{\Delta}}{n+2} \right) \\ &= \frac{\log \Delta}{n(n+1)(n+2)} + \frac{2}{n+1} \log(1 - \gamma^{n+1}) - \frac{\log(1 - \gamma^n)}{n} - \frac{\log(1 - \gamma^{n+2})}{n+2}. \end{aligned}$$

The function $f(x) = \log(1+x)$ on the interval $(-1, +\infty)$ is concave since $f''(x) = -1/(x+1)^2 < 0$. Note that $|\gamma| < 1$. If $-|\gamma| \leq x \leq 0$, then $t = -x/|\gamma| \in [0, 1]$ and hence

$$f(x) = f(t(-|\gamma|) + (1-t)0) \geq tf(-|\gamma|) + (1-t)f(0) = qx,$$

where $q = -\log(1 - |\gamma|)/|\gamma| > 0$. Note also that $\log(1+x) < x$ for $x > 0$. So we have

$$\begin{aligned} \log(1 - \gamma^{n+1}) &\geq \log(1 - |\gamma|^{n+1}) \geq -q|\gamma|^{n+1}, \\ \log(1 - \gamma^n) &\leq \log(1 + |\gamma|^n) < |\gamma|^n, \\ \log(1 - \gamma^{n+2}) &\leq \log(1 + |\gamma|^{n+2}) < |\gamma|^{n+2}. \end{aligned}$$

Therefore

$$\Delta_n > \frac{\log \Delta}{n(n+1)(n+2)} - |\gamma|^n \left(\frac{2q|\gamma|}{n+1} + \frac{1}{n} + \frac{|\gamma|^2}{n+2} \right)$$

and hence

$$\begin{aligned} n(n+1)(n+2)\Delta_n &> \log \Delta - |\gamma|^n (2q|\gamma|n(n+2) + (n+1)(n+2) + |\gamma|^2n(n+1)). \quad (1) \end{aligned}$$

Since $\lim_{n \rightarrow \infty} n^2|\gamma|^n = 0$, when $\Delta > 1$ we have $\Delta_n > 0$ for large n .

Now it remains to consider the case $\Delta = 1$. Clearly $\gamma = (A-1)/(A+1) > 0$. Recall that $\log(1-x) < -x$ for $x \in (0, 1)$. As

$$\frac{d}{dx}(\log(1-x) + x + x^2) = -\frac{1}{1-x} + 1 + 2x = \frac{x(1-2x)}{1-x} > 0 \quad \text{for } x \in (0, 0.5),$$

we have $\log(1 - x) + x + x^2 > \log 1 + 0 + 0^2 = 0$ for $x \in (0, 0.5)$. If n is large enough, then $\gamma^n < 0.5$ and hence

$$\Delta_n = \frac{2}{n+1} \log(1 - \gamma^{n+1}) - \frac{\log(1 - \gamma^n)}{n} - \frac{\log(1 - \gamma^{n+2})}{n+2} > w_n,$$

where

$$w_n := \frac{2}{n+1}(-\gamma^{n+1} - \gamma^{2n+2}) + \frac{\gamma^n}{n} + \frac{\gamma^{n+2}}{n+2}.$$

Note that

$$\lim_{n \rightarrow \infty} \frac{nw_n}{\gamma^n} = -2\gamma + 1 + \gamma^2 = (1 - \gamma)^2 > 0.$$

So, for sufficiently large n we have $\Delta_n > w_n > 0$.

Now we show that $n \geq 4$ suffices in the case $A = 1$ and $B = -1$. Note that $\Delta = 5$ and $\gamma \approx -0.382$. The sequence $(|\gamma|^n(n+1)(n+2))_{n \geq 1}$ is decreasing since

$$|\gamma| \frac{(n+2)(n+3)}{(n+1)(n+2)} < \frac{1}{2} \left(1 + \frac{2}{n+1} \right) \leq 1$$

for $n \geq 1$. It follows that $|\gamma|^n(n+1)(n+2) \leq \gamma^6 \times 7 \times 8 < 1/3$ for $n \geq 6$. In view of (1), if $n \geq 6$ then

$$\begin{aligned} n(n+1)(n+2)\Delta_n &> \log 5 - |\gamma|^n(n+1)(n+2) (2q|\gamma| + 1 + |\gamma|^2) \\ &> \log 5 - \frac{1 + 1 + \gamma^2}{3} > \log 5 - 1 > 0. \end{aligned}$$

It is easy to verify that Δ_4 and Δ_5 are positive. So $(\sqrt[n+1]{F_{n+1}}/\sqrt[n]{F_n})_{n \geq 4}$ is strictly decreasing.

In view of the above, we have completed the proof of Theorem 1.1. □

3. Proof of Theorem 1.2

PROOF OF THEOREM 1.2. Let $n \geq 3$. It is well known that $|D_n - n!/e| \leq 1/2$ (cf. [1, p. 67]). Applying the Intermediate Value Theorem in calculus, we obtain

$$\left| \log D_n - \log \left(\frac{n!}{e} \right) \right| \leq \left| D_n - \frac{n!}{e} \right| \leq 0.5.$$

Set $R_0(n) = \log D_n - \log n!$. Then $|R_0(n)| \leq 1.5$.

Since $\lim_{n \rightarrow \infty} R_0(n)/n = 0$, we have

$$\begin{aligned} \lim_{n \rightarrow \infty} \left(\frac{\log D_{n+1}}{n+1} - \frac{\log D_n}{n} \right) &= \lim_{n \rightarrow \infty} \left(\frac{\log((n+1)!)}{n+1} - \frac{\log(n!)}{n} \right) \\ &= \lim_{n \rightarrow \infty} \frac{n \log(n+1) + n \log(n!) - (n+1) \log(n!)}{n(n+1)} \\ &= \lim_{n \rightarrow \infty} \frac{n \log n + n \log(1+1/n) - \log(n!)}{n(n+1)} \\ &= \lim_{n \rightarrow \infty} \frac{\log(n^n/n!)}{n(n+1)}. \end{aligned}$$

As $n! \sim \sqrt{2\pi n}(n/e)^n$ (i.e., $\lim_{n \rightarrow \infty} n!/(\sqrt{2\pi n}(n/e)^n) = 1$) by Stirling's formula, we have $\log(n^n/n!) \sim n$ and hence

$$\lim_{n \rightarrow \infty} \left(\frac{\log D_{n+1}}{n+1} - \frac{\log D_n}{n} \right) = 0.$$

Thus $\lim_{n \rightarrow \infty} \sqrt[n+1]{D_{n+1}}/\sqrt[n]{D_n} = 1$.

From the known identity $D_n/n! = \sum_{k=0}^n (-1)^k/k!$, we have the recurrence $D_n = nD_{n-1} + (-1)^n$ for $n > 1$. Thus, if $n \geq 3$ then

$$R_0(n) - R_0(n-1) = \log \frac{D_n}{n!} - \log \frac{D_{n-1}}{(n-1)!} = \log \frac{D_n}{nD_{n-1}} = \log \left(1 + \frac{(-1)^n}{nD_{n-1}} \right).$$

Fix $n \geq 4$. If n is even, then

$$0 < R_0(n) - R_0(n-1) = \log \left(1 + \frac{1}{nD_{n-1}} \right) < \frac{1}{nD_{n-1}} = \frac{1}{D_n - 1} \leq \frac{3}{D_n + 0.5}.$$

If n is odd, then

$$0 > R_0(n) - R_0(n-1) = \log \left(1 - \frac{1}{nD_{n-1}} \right) > \frac{-2}{nD_{n-1}} = \frac{-2}{D_n + 1} \geq \frac{-3}{D_n + 0.5}$$

since $\log(1-x) + 2x > 0$ for $x \in (0, 0.5)$. So

$$|R_0(n) - R_0(n-1)| < \frac{3}{D_n + 0.5} \leq \frac{3e}{n!}$$

and hence

$$\left| \frac{R_0(n-1) - R_0(n)}{n-1} \right| < \frac{3e}{n!(n-1)} \leq \frac{3e}{n(n-1)(n+1)}.$$

Similarly, we also have

$$\left| \frac{R_0(n+1) - R_0(n)}{n+1} \right| < \frac{3e}{n!(n+1)} \leq \frac{3e}{n(n-1)(n+1)}.$$

Therefore,

$$\begin{aligned} & \left| \frac{R_0(n+1)}{n+1} - \frac{2R_0(n)}{n} + \frac{R_0(n-1)}{n-1} - \frac{2R_0(n)}{n(n-1)(n+1)} \right| \\ &= \left| \frac{R_0(n+1) - R_0(n)}{n+1} + \frac{R_0(n-1) - R_0(n)}{n-1} \right| \leq \frac{6e}{n(n-1)(n+1)} \end{aligned}$$

and hence

$$\left| \frac{R_0(n+1)}{n+1} - \frac{2R_0(n)}{n} + \frac{R_0(n-1)}{n-1} \right| \leq \frac{2|R_0(n)| + 6e}{n(n-1)(n+1)} \leq \frac{6e+3}{n(n-1)(n+1)}.$$

Thus $|R_1(n)| \leq 6e+3$, where

$$R_1(n) := n(n-1)(n+1) \left(\frac{R_0(n+1)}{n+1} - \frac{2R_0(n)}{n} + \frac{R_0(n-1)}{n-1} \right).$$

Since

$$\begin{aligned} \log((n-1)!) &= \sum_{k=1}^{n-1} \int_k^{k+1} (\log k) dx < \sum_{k=1}^{n-1} \int_k^{k+1} \log x dx \\ &= \int_1^n \log x dx = n \log n - n + 1 < \sum_{k=1}^{n-1} \int_k^{k+1} (\log(k+1)) dx = \log(n!), \end{aligned}$$

we have

$$n \log n - n < \log(n!) = \log((n-1)!) + \log n < n \log n - n + \log n + 1$$

and so $\log(n!) = n \log n - n + R_2(n)$ with $|R_2(n)| < \log n + 1$.

Observe that

$$\begin{aligned} & \frac{\log D_{n+1}}{n+1} - \frac{2}{n} \log D_n + \frac{\log D_{n-1}}{n-1} \\ &= \frac{\log((n+1)!) - 2 \log(n!) + \log((n-1)!)}{n+1} + \frac{R_1(n)}{(n-1)n(n+1)} \\ &= \frac{2 \log(n!)}{(n-1)n(n+1)} - \frac{\log n}{n-1} + \frac{\log(n+1)}{n+1} + \frac{R_1(n)}{(n-1)n(n+1)} \end{aligned}$$

$$\begin{aligned}
&= -\frac{2n}{(n-1)n(n+1)} + \frac{\log(n+1) - \log n}{n+1} + \frac{2R_2(n) + R_1(n)}{(n-1)n(n+1)} \\
&\leq -\frac{2n}{(n-1)n(n+1)} + \frac{n-1}{(n-1)n(n+1)} + \frac{2R_2(n) + R_1(n)}{(n-1)n(n+1)} \\
&= -\frac{n+1 - 2R_2(n) - R_1(n)}{(n-1)n(n+1)}.
\end{aligned}$$

If $n \geq 27$, then $n+1 - 2R_2(n) - R_1(n) > n - 2\log n - 1 - 6e - 3 > 0$, and hence we get

$$\log \frac{\sqrt[n]{D_n}}{\sqrt[n-1]{D_{n-1}}} > \log \frac{\sqrt[n+1]{D_{n+1}}}{\sqrt[n]{D_n}}.$$

By a direct check via computer, the last inequality also holds for $n = 4, \dots, 26$. Therefore, the sequence $(\sqrt[n+1]{D_{n+1}}/\sqrt[n]{D_n})_{n \geq 3}$ is strictly decreasing. This ends the proof. \square

4. Proof of Theorem 1.3

Lemma 4.1. For $x > 0$ we have

$$\log(1+x) > x - \frac{x^2}{2}. \quad (2)$$

PROOF. As

$$\frac{d}{dx} \left(\log(1+x) - x + \frac{x^2}{2} \right) = \frac{x^2}{1+x},$$

we see that $\log(1+x) - x + x^2/2 > \log 1 - 0 + 0^2/2 = 0$ for any $x > 0$. \square

Lemma 4.2. Let $m, n \in \mathbb{Z}^+$ with $n \geq 3$. If $m \geq 11$ or $n \geq 30$, then

$$H_n^{(m)} \log H_n^{(m)} > 4 \left(\frac{2}{n+2} \right)^{m-1}. \quad (3)$$

PROOF. Recall that H_n refers to $H_n^{(1)}$. If $n \geq 30$, then

$$H_n \log H_n \geq H_{30} \log H_{30} > 4,$$

and hence (3) holds for $m = 1$.

Below we assume that $m \geq 2$. As $n \geq 3$, we have

$$H_n^{(m)} \log H_n^{(m)} \geq H_3^{(m)} \log H_3^{(m)}.$$

So it suffices to show that

$$\left(\frac{n+2}{2}\right)^{m-1} H_3^{(m)} \log H_3^{(m)} > 4 \tag{4}$$

whenever $m \geq 11$ or $n \geq 30$. By Lemma 4.1,

$$\begin{aligned} \log H_3^{(m)} &= \log(1 + 2^{-m} + 3^{-m}) > 2^{-m} + 3^{-m} - \frac{(2^{-m} + 3^{-m})^2}{2} \\ &> 2^{-m} + 3^{-m} - \frac{(2^{1-m})^2}{2} = \frac{1}{2^m} + \frac{1}{3^m} - \frac{2}{4^m}. \end{aligned}$$

If $m \geq 3$, then $(4/3)^m \geq (4/3)^3 > 2$ and hence $\log H_3^{(m)} > 1/2^m$. Note also that $H_3^{(2)} \log H_3^{(2)} > 1/4$. So we always have

$$H_3^{(m)} \log H_3^{(m)} > \frac{1}{2^m}.$$

If $m \geq 11$, then $1.25^m \geq 1.25^{11} > 10$ and hence

$$\frac{1}{2^m} > \frac{4}{2.5^{m-1}} \geq \frac{4}{((n+2)/2)^{m-1}},$$

therefore (4) holds. When $n \geq 30$, we have

$$\frac{1}{2^m} \geq \frac{1}{2^{4m-6}} = \frac{4}{16^{m-1}} \geq 4 \left(\frac{2}{n+2}\right)^{m-1}$$

and hence (4) also holds. □

PROOF OF THEOREM 1.3. Let $m \geq 1$ and $n \geq 3$. Set

$$\Delta_n(m) := \log \frac{\sqrt[n+1]{H_{n+1}^{(m)}}}{\sqrt[n]{H_n^{(m)}}} - \log \frac{\sqrt[n+2]{H_{n+2}^{(m)}}}{\sqrt[n+1]{H_{n+1}^{(m)}}} = \frac{2 \log H_{n+1}^{(m)}}{n+1} - \frac{\log H_n^{(m)}}{n} - \frac{\log H_{n+2}^{(m)}}{n+2}.$$

It suffices to show that $\Delta_n(m) < 0$. This can be easily verified by computer if $m \in \{1, \dots, 10\}$ and $n \in \{3, \dots, 29\}$.

Below we assume that $m \geq 11$ or $n \geq 30$. Recall (2) and the known fact that $\log(1+x) < x$ for $x > 0$. We clearly have

$$\log \frac{H_{n+1}^{(m)}}{H_n^{(m)}} = \log \left(1 + \frac{1}{(n+1)^m H_n^{(m)}} \right) < \frac{1}{(n+1)^m H_n^{(m)}}$$

and

$$\log \frac{H_{n+2}^{(m)}}{H_n^{(m)}} > \log \left(1 + \frac{2}{(n+2)^m H_n^{(m)}} \right) > \frac{2}{(n+2)^m H_n^{(m)}} - \frac{2}{(n+2)^{2m} (H_n^{(m)})^2}.$$

It follows that

$$\begin{aligned} \Delta_n(m) &= \left(\frac{2}{n+1} - \frac{1}{n} - \frac{1}{n+2} \right) \log H_n^{(m)} + \frac{2}{n+1} \log \frac{H_{n+1}^{(m)}}{H_n^{(m)}} - \frac{1}{n+2} \log \frac{H_{n+2}^{(m)}}{H_n^{(m)}} \\ &< \frac{-2 \log H_n^{(m)}}{n(n+1)(n+2)} + \frac{2}{(n+1)^{m+1} H_n^{(m)}} \\ &\quad - \frac{2}{(n+2)^{m+1} H_n^{(m)}} + \frac{2}{(n+2)^{2m+1} (H_n^{(m)})^2}. \end{aligned}$$

Since $(n+2)^{m+1} = \sum_{k=0}^{m+1} \binom{m+1}{k} (n+1)^k$ by the binomial theorem, we obtain

$$\begin{aligned} \Delta_n(m) &\leq \frac{-2 \log H_n^{(m)}}{n(n+1)(n+2)} + \frac{2 \sum_{k=0}^m \binom{m+1}{k} (n+1)^k}{(n+1)^{m+1} (n+2)^{m+1} H_n^{(m)}} + \frac{2}{(n+2)^{m+2} H_n^{(m)}} \\ &< \frac{-2 \log H_n^{(m)}}{n(n+1)(n+2)} + \frac{2(n+1)^m \sum_{k=0}^m \binom{m+1}{k}}{(n+1)^{m+1} (n+2)^{m+1} H_n^{(m)}} + \frac{2}{(n+1)(n+2)^{m+1} H_n^{(m)}} \\ &= \frac{-2 \log H_n^{(m)}}{n(n+1)(n+2)} + \frac{2(2^{m+1} - 1) + 2}{(n+1)(n+2)^{m+1} H_n^{(m)}}. \end{aligned}$$

Thus

$$\begin{aligned} n(n+1)(n+2)\Delta_n(m) \frac{H_n^{(m)}}{2} &< -H_n^{(m)} \log H_n^{(m)} + \frac{2^{m+1}n}{(n+2)^m} \\ &< 4 \left(\frac{2}{n+2} \right)^{m-1} - H_n^{(m)} \log H_n^{(m)}. \end{aligned}$$

Applying (3) we find that $\Delta_n(m) < 0$. This completes the proof. \square

ACKNOWLEDGMENTS. The initial version of this paper was posted to arXiv in 2012 as a preprint with the ID arXiv:1208.3903. We are grateful to the two referees for their helpful comments.

References

- [1] R. P. STANLEY, Enumerative Combinatorics, Vol. 1, *Cambridge Univ. Press, Cambridge*, 1997.
- [2] Z.-W. SUN, Reduction of unknowns in diophantine representations, *Sci. China Ser. A* **35** (1992), 257–269.
- [3] Z.-W. SUN, On a sequence involving sums of primes, *Bull. Austral. Math. Soc.* **88** (2013), 197–205.
- [4] Z.-W. SUN, Conjectures involving arithmetical sequences, in: Number Theory: Arithmetic in Shangri-La, Proc. 6th China–Japan Seminar (Shanghai, August 15–17, 2011), (S. Kanemitsu, H. Li and J. Liu, eds.), *World Sci., Singapore*, 2013, 244–258.

QING-HU HOU
CENTER FOR APPLIED MATHEMATICS
TIANJIN UNIVERSITY
TIANJIN 300072
PEOPLE'S REPUBLIC OF CHINA
E-mail: qinghu.hou@gmail.com

ZHI-WEI SUN
DEPARTMENT OF MATHEMATICS
NANJING UNIVERSITY
NANJING 210093
PEOPLE'S REPUBLIC OF CHINA
E-mail: zwsun@nju.edu.cn

HAOMIN WEN
DEPARTMENT OF MATHEMATICS
THE UNIVERSITY OF PENNSYLVANIA
PHILADELPHIA, PA 19104
USA
E-mail: weh@math.upenn.edu

(Received March 25, 2013; revised March 24, 2014)