

Title: Influence of weakly \mathcal{H} -embedded subgroups on the structure of finite groups

Author(s): Mohamed Asaad, Mohamed Ramadan and Abdelrahman Heliel

Let G be a finite group, and H a subgroup of G . We say that H is an \mathcal{H} -subgroup in G if $N_G(H) \cap H^g \leq H$ for any $g \in G$. We say that H is weakly \mathcal{H} -embedded in G if G has a normal subgroup K such that $H^G = HK$ and $H \cap K$ is an \mathcal{H} -subgroup in G . For each prime p dividing the order of G , let P be a non-cyclic Sylow p -subgroup of G . We fix a p -power integer d with $1 < d < |P|$, and study the structure of G under the assumption that each subgroup of P of order d and pd is weakly \mathcal{H} -embedded in G . Some new results about the p -nilpotency and supersolvability of G are obtained.

Address:

Mohamed Asaad
Department of Mathematics
Faculty of Science
Cairo University
Giza, 12613
Egypt

Address:

Mohamed Ramadan
Department of Mathematics
Faculty of Science
Cairo University
Giza, 12613
Egypt

Address:

Abdelrahman Heliel
Department of Mathematics
Faculty of Science
Beni-Suef University
Beni-Suef 62511
Egypt