

Title: Riccati technique and oscillation constant for modified Euler type half-linear equations

Author(s): Petr Hasil, Jaroslav Jaroš and Michal Veselý

We study equations in the form

$$[r(t)t^{p-1}\Phi(x')] + \frac{s(t)}{t \log^p t} \Phi(x) = 0, \quad t \in [a, \infty),$$

where x is the unknown function, with the scalar p -Laplacian $\Phi(x) = |x|^{p-1} \operatorname{sgn} x$ for an arbitrarily given number $p > 1$. It is known that these equations are conditionally oscillatory for some coefficients. The conditional oscillation for certain non-constant coefficients r, s has been proved via the Prüfer angle. Using a new modification of the Riccati method (i.e., by a different approach), we identify easy-to-use conditions on the coefficients which assure the conditionally oscillatory behaviour as well. The obtained results cover equations whose oscillatory properties were not known and these results are new even for linear equations (i.e., for $p = 2$).

Address:

Petr Hasil
Department of Mathematics
and Statistics
Faculty of Science
Masaryk University
Kotlářská 2
611 37 Brno
Czech Republic

Address:

Jaroslav Jaroš
Department of Mathematical Analysis
and Numerical Mathematics
Faculty of Mathematics,
Physics and Informatics
Comenius University
Mlynská dolina F1
842 48 Bratislava
Slovakia

Address:

Michal Veselý
Department of Mathematics
and Statistics
Faculty of Science
Masaryk University
Kotlářská 2
611 37 Brno
Czech Republic