

Title: D'Alembert's functional equation on monoids with both an endomorphism and an anti-endomorphism

Author(s): Mohamed Ayoubi, Driss Zeglami and Aziz Mouzoun

Let M be a topological monoid, e its neutral element, let \mathbb{K} denote a division ring of characteristic $\neq 2$, and let $\psi : M \rightarrow M$ (resp. $\varphi : M \rightarrow M$) be a continuous anti-endomorphism (resp. endomorphism) of M as a semigroup. We solve the functional equation

$$g(xy) - g(x\psi(y)) = cg(x)g(y), \quad x, y \in M,$$

where $g : M \rightarrow \mathbb{K}$ is the unknown function and $c \in \mathbb{K} \setminus \{0\}$. This enables us to find, when $\psi(e) = \varphi(e) = e$, the solutions $g : M \rightarrow \mathbb{C}$ of each of the new functional equations

$$g(\varphi(x)y) \pm g(x\psi(y)) = 2g(x)g(y), \quad x, y \in M.$$

We also find the continuous, complex valued solutions of a Van Vleck's functional equation with an anti-endomorphism on M .

Address:

Mohamed Ayoubi
Department of Mathematics
E.N.S.A.M, Moulay Ismail University
B.P. 15290 Al Mansour, Meknes
Morocco

Address:

Driss Zeglami
Department of Mathematics
E.N.S.A.M, Moulay Ismail University
B.P. 15290 Al Mansour, Meknes
Morocco

Address:

Aziz Mouzoun
Department of Mathematics
E.N.S.A.M, Moulay Ismail University
B.P. 15290 Al Mansour, Meknes
Morocco