

## Rings of finite rank

By R. E. JOHNSON (Rochester, N. Y.)

Let  $R$  be a ring and  $L_r = L_r(R)$  be its lattice of right ideals. The *right rank* of  $R$ ,  $r(R)$ , is defined to be  $\max. \text{card } S^\perp$ , where  $S^\perp$  is an independent subset of  $L_r$ . The *left rank* of  $R$ ,  $l(R)$ , is defined similarly. If  $R$  is a zero ring, then  $r(R) = l(R) = k$  where  $k$  is the usual rank of an Abelian group. Our remarks are restricted in this note to rings of finite right rank.

Associated with the lattice  $L_r$  of ring  $R$  is another lattice  $L'_r$  defined as follows. For  $A, B \in L_r$ , let  $A \subset' B$  signify that  $B$  is an essential extension of  $A$ ; that is,  $A \subset B$  and  $A \cap C \neq 0$  whenever  $B \cap C \neq 0$ ,  $C \in L_r$ . Define the relation  $\sim$  in  $L_r$  by:  $A \sim B$  iff  $A \cap B \subset' A$  and  $A \cap B \subset' B$ . It is readily shown that  $\sim$  is an equivalence relation. Let  $L'_r = L_r / \sim$  and  $\varphi$  be the natural mapping of  $L_r$  onto  $L'_r$ . If the partial ordering  $\cong$  is defined in  $L'_r$  by:  $\varphi A \cong \varphi B$  iff  $A \cap B \subset' A$ , then UTUMI showed in [1] that  $L'_r$  is a complemented modular lattice. Furthermore, he showed that  $\varphi$  is a meet homomorphism of  $L_r$  onto  $L'_r$ . Evidently  $\varphi 0 = \{0\}$  and  $\varphi R = \{A \in L_r \mid A \subset' R\}$ . It may be shown that  $\varphi(A \cup B) = \varphi A \cup \varphi B$  if  $A \cap B = 0$ . From these remarks, it is clear that  $S^\perp$  in  $L_r$  iff  $(\varphi S)^\perp$  in  $L'_r$ . Consequently,  $r(R)$  is simply the dimension of lattice  $L'_r$ . Hence,  $r(R) = \text{card } S^\perp$  where  $S^\perp$  is any maximal independent subset of  $L_r$ .

Each right ideal  $A$  of ring  $R$  has a rank defined by  $r(A) = \dim(\varphi A)$  in  $L'_r$ . The *right rank* of an element  $x$  of  $R$ ,  $r(x)$ , is defined to be  $r(A)$  where  $A$  is the right ideal of  $R$  generated by  $x$ . It is evident that

$$r(xy) \cong r(x), \quad r(x+y) \cong r(x) + r(y)$$

for all  $x, y \in R$ .

It does not seem possible to say much about the rank of the elements of a general ring of finite rank. However, if we assume that  $R$  has zero singular ideal,  $R_r^\Delta = 0$ , then some of the familiar properties of rank hold. We recall that  $R_r^\Delta = \{x \in R \mid x^r \subset' R\}$ , where  $x^r$  is the right annihilator of  $x$  in  $R$ . Let us call  $R$  an  $F_r$ -ring if  $R$  has finite right rank and  $R_r^\Delta = 0$ . For an  $F_r$ -ring  $R$ , it is easily shown that  $r(x) = r(xC)$  for every  $x \in R$  and every  $C \in L_r$  such that  $C \subset' R$ .

If  $R$  is an  $F_r$ -ring, then each  $A \in L_r$  has a unique maximal essential extension  $A^* \in L_r$ , called the closure of  $A$ . The set  $L_r^*$  of all closed right ideals of  $R$  is a lattice, which is easily shown to be isomorphic to  $L'_r$ . In fact,  $\varphi A = \{B \in L_r \mid B \subset' A^*\}$  for each  $A \in L_r$ . Thus,  $r(x) = r(xR) = \dim(xR)$  in  $L_r^*$ . Since  $x^r \in L_r^*$  for each  $x \in R$ , and a maximal complement of each  $A \in L_r$  is in  $L_r^*$ , it is evident that  $r(R) = 1$  iff  $R$  is a right Ore domain.

**Theorem 1.** *If  $R$  is an  $F_r$ -ring, then  $r(x) = r(R) - r(x^r)$  for every  $x \in R$ .*

PROOF. Let  $A$  be a complement of  $x^r$  in  $L_r^*$  and  $\{A_1, \dots, A_k\}$  be an atomic basis for  $A$ . Since  $\{xA_1, \dots, xA_k\}^\perp$ , evidently  $r(x) \cong k = r(R) - r(x^r)$ . If  $\{xB_1, \dots, xB_n\}^\perp$ , where each  $B_i$  is an atom of  $L_r^*$ , and if  $B = B_1 + \dots + B_n$ , then  $B \cap x^r = 0$ . For if  $b = b_1 + \dots + b_n \in x^r$ ,  $b_i \in B_i$ , then  $xb = \sum xb_i = 0$  and  $xb_i = 0$  for each  $i$ . However,  $x^r \cap B_i = 0$  by assumption, and therefore  $b_i = 0$  for each  $i$  and  $b = 0$ . Thus,  $B$  is contained in a maximal complement of  $x^r$  and  $n \leq k$ . Consequently,  $r(x) \leq k$  and the theorem is proved.

Since  $(xy)^r \supset y^r$ , evidently

$$r(xy) \leq r(y)$$

for all  $x, y$  in an  $F_r$ -ring by Theorem 1.

**Theorem 2.** *If  $R$  is an  $F_r$ -ring and  $x, y \in R$  are such that  $xR \cap yR = 0$ , then  $r(x+y) \cong r(x)$ . If, furthermore,  $x^r + y^r \subset' R$  then  $r(x+y) = r(x) + r(y)$ .*

PROOF. If  $r(x) = k$  and  $\{xA_1, \dots, xA_k\}^\perp$ ,  $A_i$  atoms of  $L_r^*$ , then  $\{(x+y)A_1, \dots, (x+y)A_k\}^\perp$ . For if  $\sum (x+y)a_i = 0$ ,  $a_i \in A_i$ , then  $\sum xa_i = -\sum ya_i = 0$  and  $xa_i = 0$  for each  $i$ . Hence,  $a_i = 0$  for each  $i$ . Therefore,  $r(x+y) \cong r(x)$ .

To prove the second part, let  $B$  and  $C$  be relative complements of  $x^r \cap y^r$  in  $x$  and  $y^r$ , respectively, and let  $\{B_1, \dots, B_k\}$ ,  $\{C_1, \dots, C_m\}$ , and  $\{D_1, \dots, D_n\}$  be atomic bases of  $B, C$ , and  $x^r \cap y^r$ , respectively. If  $B' = B_1 + \dots + B_k$ ,  $C' = C_1 + \dots + C_m$ , and  $D' = D_1 + \dots + D_n$ , then  $B' + C' + D' \subset' R$  and  $r(x+y) = r[(x+y)(B' + C' + D')] = r[(x+y)(B' + C')]$ . Since  $(x+y)B' = yB'$  and  $(x+y)C' = xC'$ , evidently  $(x+y)B' \cap (x+y)C' = 0$ . Hence,  $r(x+y) = k + m = (k + m + n - k - n) + (k + m + n - m - n) = r(x) + r(y)$  in view of Theorem 1. This proves Theorem 2.

If  $R$  is an  $F_r$ -ring and  $U = \{u \in R \mid uR \subset' R\}$ , then  $U$  is a multiplicative semi-group by [2; 3.2]. Also, by [2; 3.3],  $u^r = u^l = 0$  for every  $u \in U$ . Since  $(ux)^r = x^r$  for all  $u \in U$  and  $x \in R$ , evidently  $r(ux) = r(x)$ . If  $r(x) = k$  and  $\{xA_1, \dots, xA_k\}^\perp$ ,  $A_i$  atoms of  $L_r^*$ , then for each  $u \in U$  we can select  $B_i \in L_r$  such that  $uR \cap A_i = uB_i$  for each  $i$ . Since  $(uB_i)^* = A_i$ , evidently  $xuB_i \neq 0$  for each  $i$ . Consequently,  $r(xu) = k$ . We have proved that

$$r(xu) = r(ux) = r(x)$$

for all  $x \in R$  and  $u \in U$ .

Let us call an  $F_r$ -ring  $R$  an  $I_r$ -ring if every  $A \in L_r^*$  contains an element  $a$  such that  $aR \subset' A$ . Thus, an  $I_r$ -ring is a generalization of a principal right ideal ring. If  $R$  is an  $I_r$ -ring then for each  $A \in L_r^*$ ,  $r(A) = r(a)$  for some  $a \in A$ . In particular,  $U \neq \Phi$  for an  $I_r$ -ring. If  $\{R_1, \dots, R_n\}$  is a set of  $I_r$ -rings, then their direct product  $R_1 \times \dots \times R_n$  is easily seen to be an  $I_r$ -ring. Also, if  $Q$  is a (right) quotient ring of an  $I_r$ -ring  $R$  (so that  $qR \cap R \neq 0$  for each nonzero  $q \in Q$ ), then  $Q$  is an  $I_r$ -ring (see [3]).

An  $F_r$ -ring is called (right) irreducible [3] iff  $\{0, R\}$  is the center of lattice  $L_r^*$ . If  $R$  is not irreducible, then the center  $C_r^*$  of  $L_r^*$  is a Boolean algebra and each atom of  $C_r^*$  is an irreducible ring. If  $\{S_1, \dots, S_n\}$  is the set of atoms of  $C_r^*$  and  $S = S_1 + \dots + S_n$  (a direct sum), then  $R$  is a quotient ring of  $S$ . Evidently  $R$  is an  $I_r$ -ring iff every  $S_i$  is an  $I_r$ -ring. Thus, the problem of describing  $I_r$ -rings reduces to that of describing irreducible  $I_r$ -rings.

In a forthcoming paper [4], an  $F_r$ -ring  $R$  is called *right potent* iff  $A^2 \neq 0$  for every atom  $A \in L_r^*$ . This is equivalent to saying that no nonzero ideal of  $L_r^*$  is nilpotent. Let us call a right potent, irreducible  $F_r$ -ring a  $P_r$ -ring.

**Theorem 3.** *If  $R$  is a  $P_r$ -ring, then each  $A \in L_r^*$  contains an element  $a$  such that  $A \dot{\cup} a^r = R$ .*

**PROOF.** The notation  $\dot{\cup}$  is used for a direct union in lattice  $L_r^*$ . Let  $r(R) = n$ . If  $n = 1$ , then  $R$  is a right Ore domain and the theorem is trivially true. So let us assume that  $n > 1$ . If  $A \in L_r^*$  is an atom, so that  $r(A) = 1$ , then  $A \cap A^r = 0$  and  $A \cap a^r = 0$  for some nonzero  $a \in A$ . Since  $a^r$  is a maximal element ( $\neq R$ ) of  $L_r^*$ , evidently  $A \dot{\cup} a^r = R$  in this case.

Assume that the integer  $k > 1$  is chosen so that the theorem is true for every element of  $L_r^*$  of rank  $< k$ , and let  $A \in L_r^*$ ,  $r(A) = k$ . Select  $B \in L_r^*$  such that  $B \subset A$  and  $r(B) = k - 1$ . By assumption, there exists some  $b \in B$  such that  $B \dot{\cup} b^r = R$ . Since  $r(b^r) = n - k + 1$  and  $B \cap b^r = 0$ , evidently  $b^r \cap A = C$  where  $C$  is an atom of  $L_r^*$ . Let us select a nonzero  $c \in C$  such that  $c^r \cap C = 0$ , and then let  $a = b + c$ . Clearly  $b^r \cap c^r = 0$  and also  $b^r + c^r \subset R$ . Hence,  $r(a) = k$  by Theorem 2. If  $x + y \in a^r \cap (B + C)$ , with  $x \in B$  and  $y \in C$ , then  $(b + c)(x + y) = 0$  and  $b(x + y) = -c(x + y) = 0$ . Therefore,  $b(x + y) = 0$ ,  $bx = 0$ , and  $x = 0$ ; and  $cy = 0$  and  $y = 0$ . We conclude that  $a^r \cap (B + C) = 0$  and consequently that  $a^r \cap A = 0$ . Hence,  $a^r \dot{\cup} A = R$ . The theorem now follows by mathematical induction.

**Corollary.** *If  $R$  is a  $P_r$ -ring, then  $R$  is an  $I_r$ -ring. In fact, for each  $A \in L_r^*$  there exist  $a \in A$  and  $b \in a^r$  such that  $a + b \in U$ .*

**PROOF.** Select  $a \in A$  so that  $A \dot{\cup} a^r = R$  and  $b \in a^r$  so that  $a^r \dot{\cup} b^r = R$ . Then  $(a + b)^r = a^r \cap b^r = 0$  and  $a + b \in U$  by [2; 3, 4].

An  $I_r$ -ring need not be a  $P_r$ -ring. Consider, for example, the ring  $R$  of all matrices of the form

$$\begin{pmatrix} a + c & 0 \\ b & c \end{pmatrix}$$

where  $c \in F$ , a field, and  $a, b \in xF[x]$ . Since the  $2 \times 2$  matrix ring  $(F(x))_2$  is a quotient ring of  $R$ , we must have  $R_r^\circ = 0$  and  $r(R) = 2$ . Every atom of  $L_r^*$  contains elements of rank 1 and the only element of  $L_r^*$  of rank 2,  $R$ , has a unity and hence contains an element of rank 2. Thus,  $R$  is an  $I_r$ -ring. However,  $R$  is not a  $P_r$ -ring since  $A = e_{21}R$  is an atom of  $L_r^*$  such that  $A^2 = 0$ .

The ring  $R$  of  $n \times n$  triangular matrices over a field is an example of a  $P_r$ -ring (and  $P_l$ -ring) which is not a principal right ideal ring. However,  $R$  is a Baer ring; i. e., every annihilating right ideal of  $R$  is generated by an idempotent. In this example,  $L_r^*$  is precisely the set of annihilating right ideals.

Let  $R$  be an  $F_r$ - and  $F_l$ -ring,  $R_r^0$  be the union in  $L_r$  of the atoms of  $L_r^*$ , and  $R_l^0$  be the corresponding union in  $L_l^*$ . The ring  $R$  is called *stable* in [5] if  $(R_r^0)^r = (R_l^0)^l = 0$ . It is proved in [5; 3, 1] that if  $R$  is stable then the lattices  $L_r^*$  and  $L_l^*$  are dual isomorphic under the correspondence  $A \rightarrow A^l$ ,  $A \in L_r^*$ . Hence,  $r(R) = l(R)$  if  $R$  is stable.

**Theorem 4.** *If  $R$  is a stable ring, then  $r(x) = l(x)$  for every  $x \in R$ .*

PROOF. By Theorem 1,  $r(x) = r(R) - r(x^r)$ . Since  $x^{rl} = (Rx)^*$  and  $r(x^r) := r(R) - l[(Rx)^*]$  by the dual isomorphism between  $L_r^*$  and  $L_l^*$ , we have  $r(x) = l[(Rx)^*] = l(x)$  as desired.

### References

- [1] Y. UTUMI, On complemented modular lattices meet-homomorphic to a modular lattice, *Kodai Math. Sem.* **4** (1952), 99–100.
- [2] R. E. JOHNSON and E. T. WONG, Quasi-injective modules and irreducible rings, *J. London Math. Soc.* **36** (1961), 260–268.
- [3] R. E. JOHNSON, Quotient rings of rings with zero singular ideal, *Pacific J. Math.* **11** (1961), 1385–1392.
- [4] R. E. JOHNSON, Potent rings, to appear.
- [5] R. E. JOHNSON, Rings with zero right and left singular ideal, to appear.

(Received January 28, 1964.)