

## A new proof on the free product structure of Hecke groups

By RONALD J. EVANS (Madison, Wis.)

In this paper, matrices are viewed as linear fractional transformations on  $H = \{\tau: \text{Im } \tau > 0\}$ . Let  $I = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$ ,  $T = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$ ,  $S_\lambda = \begin{pmatrix} 1 & \lambda \\ 0 & 1 \end{pmatrix}$ , and  $A = A_\lambda = TS_\lambda$ . Let  $Z$  denote the integers and let  $C$  denote the set of  $\lambda$  of the form  $\lambda = 2 \cos(\pi/q)$ , where  $q \in Z$ ,  $q \geq 3$ . It is well-known [2, p. 1399] that when  $\lambda \in C$ , the Hecke group  $G(\lambda) = \langle A, T \rangle$  is the free product of  $\langle A \rangle$  and  $\langle T \rangle$  (write  $G(\lambda) = \langle A \rangle * \langle T \rangle$ ). We present here a short new proof of this theorem. A short proof in the case  $\lambda = 1$  is given in [1].

**Lemma.** *Suppose that  $\lambda \in C$ ,  $\tau = x + iy \in H$ , and  $I \neq W \in \langle A \rangle$ . If  $x \geq 0$ , then  $\text{Re}(W\tau) < 0$ .*

**PROOF.** Write  $\lambda = 2 \cos \alpha$  where  $\alpha = \pi/q$ . It is easily proved by induction that for all  $n \in Z$ ,

$$A^n = \begin{pmatrix} a_n & b_n \\ c_n & d_n \end{pmatrix} = \csc \alpha \begin{pmatrix} \sin \alpha(1-n) & -\sin \alpha n \\ \sin \alpha n & \sin \alpha(n+1) \end{pmatrix}.$$

Since  $A$  has order  $q$ , we may write  $W = A^m$ , where  $1 \leq m \leq q-1$ . Write

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix} = \begin{pmatrix} a_m & b_m \\ c_m & d_m \end{pmatrix}.$$

Then

$$\text{Re}(W\tau) = \frac{(ax+b)(cx+d) + acy^2}{|c\tau+d|^2}.$$

Note that  $a, b \leq 0$  and  $c, d \geq 0$ . Hence, if  $x \geq 0$ , then  $acy^2 \leq 0$  and  $(ax+b)(cx+d) \leq 0$ . Moreover, one of these two inequalities is strict; for, if  $(ax+b)(cx+d) = 0$ , then  $bd = 0$  so that  $m = q-1$  and hence  $acy^2 < 0$ . Thus  $\text{Re}(W\tau) < 0$ . QED.

**Theorem.** If  $\lambda \in C$ , then  $G(\lambda) = \langle A \rangle * \langle T \rangle$ .

**PROOF.** Let  $\lambda \in C$ . Suppose that  $I \neq W_j \in \langle A \rangle$  ( $j=1, 2, \dots$ ). By the Lemma,  $\text{Re } W_1(i) < 0$ , so that  $\text{Re } TW_1(i) > 0$ . It follows similarly by induction that for all  $n \geq 1$ ,  $\text{Re } TW_n \dots TW_1(i) > 0$ . It follows that  $I$  and  $T$  are the only reduced words in  $\langle A, T \rangle$  which fix  $i$ , and hence  $G(\lambda) = \langle A \rangle * \langle T \rangle$ .

**References**

- [1] D. HARDY and R. WISNER, New proof of a basic theorem about the modular group, *Publ. Math. (Debrecen)* **18** (1971), 109—110.
- [2] R. C. LYNDON and J. L. ULLMAN, Groups generated by two parabolic linear fractional transformations, *Can. J. Math.* **21** (1969), 1388—1403.

RONALD J. EVANS  
UNIVERSITY OF WISCONSIN  
MADISON, WISC. 53706  
U.S.A.

(Received July 25, 1973.)