

## Remark on Ky Fan convexity

By ISTVÁN JOÓ (Budapest)

**Abstract.** In the paper is proved that the Nikaido-Isolda's theorem fails to hold if concavity is replaced by Ky Fan concavity.

Let  $X, Y$  be arbitrary sets. The function

$$f : X \times Y \rightarrow \mathbb{R}$$

is called Ky Fan concave in the variable  $x$  if

$$\begin{aligned} \forall x_1, x_2 \in X \quad \forall \lambda \in [0, 1] \quad \exists x_3 \in X \quad \forall y. \\ f(x_3, y) \geq \lambda f(x_1, y) + (1 - \lambda)f(x_2, y). \end{aligned}$$

The concavity with respect to  $y$  is defined symmetrically. In [1] the authors stated the following

**Theorem.** *There exists functions  $f_1, f_2 \in C^\infty([0, 1] \times [0, 1])$  such that  $f_1$  is Ky Fan concave in the variable  $x$ ,  $f_2$  is Ky Fan concave in the variable  $y$  and the pair  $f_1, f_2$  has no saddle point i.e. there is no point  $(x_0, y_0)$  satisfying*

$$\begin{aligned} f_1(x_0, y_0) &\geq f_1(x_1 y_0) \quad \forall x \\ f_2(x_0, y_0) &\geq f_2(x_0 y) \quad \forall y. \end{aligned}$$

This is a counterexample showing that the Nikaido-Isolda theorem fails to hold if concavity is replaced by Ky Fan concavity. The proof given in [1] was not correct; it suggested that there are polynomials  $f_1, f_2$

---

*Mathematics Subject Classification:* 90D05.

This work was supported by the Hungarian Scientific Foundation on OTKA No. T7546.

satisfying Theorem. In fact we do not know whether there are analytical functions  $f_1, f_2$  satisfying Theorem.

PROOF of Theorem. Define the functions  $k_1, k_2 : [0, 1] \times [0, 1]$  as follows. Let  $0 < \delta < \frac{1}{4}$  be fixed. For  $0 \leq x \leq \frac{1}{4}$  the function  $k_1(x), k_2(x)$  varies linearly from  $(0, 1)$  to  $(\frac{1}{2} + \delta, \frac{1}{2} - \delta)$ ; for  $\frac{1}{4} \leq x \leq \frac{1}{2}$  it goes linearly from  $(\frac{1}{2} + \delta, \frac{1}{2} - \delta)$  to  $(0, 0)$ , for  $\frac{1}{2} \leq x \leq \frac{3}{4}$  from  $(0, 0)$  to  $(\frac{1}{2} - \delta, \frac{1}{2} + \delta)$  and for  $\frac{3}{4} \leq x \leq 1$  from  $(\frac{1}{2} - \delta, \frac{1}{2} + \delta)$  to  $(1, 0)$ .

We can suppose that  $((k_1(x), k_2(x)))$  is extended linearly from  $[0, \frac{1}{4}]$  to  $(-\infty, \frac{1}{4}]$  and from  $[\frac{3}{4}, 1]$  to  $[\frac{3}{4}, \infty)$ . Consider a function

$$\varphi \in C_0^\infty(\mathbb{R}), \quad \text{supp } \varphi = [-\delta, \delta], \quad \varphi \geq 0, \quad \varphi(x) = \varphi(-x) \forall x, \quad \int_{-\infty}^{\infty} \varphi = 1.$$

The existence of a such a function is widely known.

Define

$$\hat{k}_1 = k_1 * \varphi \quad \hat{k}_2 = k_2 * \varphi.$$

Introduce the sets

$$A = \{(k_1(x), k_2(x)) : x \in [0, 1]\},$$

$$\hat{A} = \{(\hat{k}_1(x), \hat{k}_2(x)) : x \in [0, 1]\}.$$

Since the convolution by  $\varphi$  gives an average of the values  $k_i(x)$ , all points of  $\hat{A}$  belongs to the (closed) convex hull of  $A$ . Hence  $\hat{A}$  lies in the triangle of vertices  $(0, 0)$ ,  $(1, 0)$ ,  $(0, 1)$ . On the other hand,  $\int_{-\infty}^{\infty} x\varphi(x)dx = 0$  implies that  $\hat{k}_i(x) = k_i(x)$  whenever  $k_i$  varies linearly in  $[x - \delta, x + \delta]$ . Consequently  $\hat{A}$  contains the side  $[(1, 0)(0, 1)]$  of the above mentioned triangle. This means that the function

$$f_1(x, y) = (1 - y)\hat{k}_1(x) + y\hat{k}_2(x)$$

is Ky Fan-concave in  $x$ ; this follows easily from the fact that

$$\forall x_1, x_2 \in [0, 1] \quad \forall \lambda \in [0, 1] \quad \exists x_3 \in [0, 1] :$$

$$\lambda \hat{k}_1(x_1) + (1 - \lambda)\hat{k}_1(x_2) \leq \hat{k}_1(x_3),$$

$$\lambda \hat{k}_2(x_1) + (1 - \lambda)\hat{k}_2(x_2) \leq \hat{k}_2(x_3);$$

see [1], p. 138 or [2] p. 204-205 for more details. Investigate the set

$$C_1 = \{(x_0, y_0) : f_1(x_0, y_0) = \max_{x \in [0, 1]} f_1(x, y_0)\}.$$

For given  $y_0$ , those values  $x_0$  are involved for which the perpendicular projection of  $(\hat{k}_1(x_0), \hat{k}_2(x_0))$  to the line along the vector  $(1 - y_0, y_0)$  is the farthest from the origin. Keeping in mind what has been proved about the set  $\hat{A}$  we see that for  $0 \leq y_0 < \frac{1}{2}$  only the point  $(1, 0)$  is projected, for  $y_0 = \frac{1}{2}$  the whole segment  $[(1, 0), (0, -1)]$  and for  $\frac{1}{2} < y_0 \leq 1$  the only point  $(0, 1)$ . Consequently (using  $\text{supp } \varphi = [-\delta, \delta]$ )

$$C_1 = \{1\} \times \left[0, \frac{1}{2}\right) \cup \left[0, \frac{1}{4}\right] \times \left\{\frac{1}{2}\right\} \cup \left[\frac{3}{4}, 1\right] \times \left\{\frac{1}{2}\right\} \cup \{0\} \times \left(\frac{1}{2}, 1\right].$$

Considered the function

$$f_2(x, y) = -(x - y)^2;$$

it is obviously concave hence also Ky Fan-concave in  $y$ . On the other hand

$$C_2 = \{(x_0, y_0) : f_2(x_0, y_0) = \max_{y \in [0, 1]} f_2(x_0, y)\}$$

is the line segment  $y = x$ ,  $0 \leq x \leq 1$  which does not meet  $C_1$ ,

$$C_1 \cap C_2 = \emptyset$$

which proves Theorem. □

### References

- [1] M. HORVÁTH and I. JOÓ, On Ky Fan convexity, *Matematikai Lapok* **34** (1-3) (1987), 137–140.
- [2] I. JOÓ, Answer to a problem of M. Horváth and A. Sövegjártó, *Annales Univ. Sci. Budapest, Sectio Math.* **29** (1986), 203–207.

ISTVÁN JOÓ  
DEPARTMENT OF ANALYSIS  
L. EÖTVÖS UNIVERSITY  
H-1088 BUDAPEST  
MÚZEUM KRT. 6-8  
HUNGARY

*(Received October 6, 1995; revised March 11, 1996)*