

Equivalent topological properties of the space of signatures of a semilocal ring

By ALEX ROSENBERG (Ithaca, New York) and ROGER WARE (University Park, Pennsylvania)

Dedicated to the memory of Andor Kertész

1. Introduction and Notations

Let R denote a (not necessarily noetherian) connected semilocal ring, $U(R)$ its group of units, and $X=X(R)$ the topological space of signatures of R [8, Def. 2.1]. The space X is compact, Hausdorff, and totally disconnected, with a subbase of the topology being given by the subsets $W(a)=\{\sigma \text{ in } X \mid \sigma(a) = -1\}$, a in $U(R)$, of X .

By a *space* over R we will mean a pair (E, B) where E is a projective (whence free) R -module and B is a nondegenerate symmetric bilinear form on E . Isometries of spaces will be written as \cong and for any natural number m the space $E \perp E \perp \dots \perp E$ (m times) will be denoted by mE or $m(E, B)$. An element e of E is called *primitive* if it can be augmented to a basis of E . The space (E, B) is called *isotropic* if it contains a primitive element e with $B(e, e)=0$. We will write $E=\langle a_1, \dots, a_n \rangle$ to mean E has an orthogonal basis e_1, \dots, e_n such that $B(e_i, e_i)=a_i$ in $U(R)$. The Witt ring of equivalence classes of spaces will be denoted by $W(R)$ and the representative of a space (E, B) in $W(R)$ by $[E]$ or $[B]$. If E is a space then there exist a_1, \dots, a_n in $U(R)$ such that $[E]=[\langle a_1, \dots, a_n \rangle]$ in $W(R)$ [7, Thm. 1.16]. Any σ in $X(R)$ induces a ring homomorphism $\bar{\sigma}: W(R) \rightarrow \mathbf{Z}$ via $\bar{\sigma}([\langle a_1, \dots, a_n \rangle])=\sigma(a_1)+\dots+\sigma(a_n)$.

In [8, Thm. 3.20], the following Strong Approximation Property was introduced:

SAP If Y is a clopen (closed and open) subset of X , then $Y=W(a)$ for some a in $U(R)$.

Following ELMAN and LAM [4, Def. 1.5], X is said to satisfy the Weak Approximation Property if

WAP The family of subsets $W(a)$ of X , as a runs through $U(R)$, forms a basis of the topology of X .

It is easy to see (cf. [8, Thm. 3.20 and 4, p. 1159], that SAP is equivalent to: If Y_1, Y_2 are two disjoint closed subsets of X , there is an element a in $U(R)$ with $\sigma(a)=1$ for all σ in Y_1 , and $\tau(a)=-1$ for all τ in Y_2 . Moreover, WAP is equivalent

to: For any closed subset Y of X and any τ in $X - Y$, there is an element a in $U(R)$ with $\sigma(a) = 1$ for all σ in Y and $\tau(a) = -1$.

If R is a formally real field ELMAN and LAM [4, p. 1184] and PRESTEL [10, p. 319], independently, introduced the following Hasse—Minkowski Property.

HMP R satisfies HMP, if for every space (E, B) such that $|\bar{\sigma}([E])| < \text{rank } E$, for all σ in X , there is a natural number m such that $m(E, B)$ is isotropic.

It is clear that SAP implies WAP and, in [4, Thm. 3.5], it was shown that if R is a formally real field SAP and WAP are equivalent. If, in addition, R is also pythagorean, [4, Thm. 5.3] shows that HMP is equivalent to the other two conditions, and, indeed, for arbitrary formally real fields, [10, Satz 3.1 and 5, Thm. C] demonstrates the equivalence of all three conditions. The proofs depend on [3, Thms. 2.6 and 4.8] which seem to be quite deep, and, in particular, Theorem 4.8 of [3] relies on the Hauptsatz of ARASON and PFISTER.

The purpose of this note is to prove the equivalence of WAP, SAP, and HMP for connected semilocal rings with 2 in $U(R)$, using only basic facts about spaces and $W(R)$, thus resulting in considerably easier proofs, even in the field case. Although we assume, for the sake of simplicity, that our rings are connected, an extension to the non-connected case presents no difficulties because of [8, Corr. 2. 18]. In Section 4, we reprove a special case of [1, Satz 2.7] for the reader's convenience.

The first author gratefully acknowledges partial support from NSF Grant GP—40773X and the second from NSF Grant GP—37781.

2. Equivalence of WAP and SAP

We begin with the following lemma which is already implicit in, i.a., [4, 7, 8]:

Lemma 2.1. *Let R be a connected semilocal ring with 2 in $U(R)$, (E_i, B_i) $i = 1, 2$, two spaces, and $r_i = \text{rank } E_i$. If $r_1 \cong r_2$, and for all σ in X , $\bar{\sigma}([E_1]) = \bar{\sigma}([E_2])$, then there is a natural number m such that*

$$2^m E_1 \cong 2^m E_2 \perp 2^{m-1}(r_1 - r_2)\mathbf{H},$$

where \mathbf{H} denotes the hyperbolic plane $\langle 1, -1 \rangle$.

PROOF. By [8, Remark 2.2], the element $[E_1] - [E_2]$ of $W(R)$ lies in each minimal prime ideal of $W(R)$. Hence by [7, Ex. 3. 11] it is a torsion element of $W(R)$, i.e., for some natural number m we have $[2^m E_1] = [2^m E_2]$. Since 2 is in $U(R)$, the Witt Cancellation Theorem holds for $R[11]$, and so the result follows.

As usual a space (E, B) is called an n -Pfister space if $E \cong \bigotimes_1^n \langle 1, a_i \rangle$, a_i in $U(R)$.

Two n -Pfister spaces E, F are *linked* [3, Def. 4.1], if there is an $(n-1)$ -Pfister space G and elements a, b in $U(R)$, such that $E \cong \langle 1, a \rangle \otimes G$ and $F \cong \langle 1, b \rangle \otimes G$. The spaces E and F are *stably linked* if $2^m E$ and $2^m F$ are linked for some natural number m .

Theorem 2.2. (cf. [4, Thm. 3.5]). *Let R denote a connected semilocal ring with 2 in $U(R)$ and $X = X(R)$ its space of signatures. Then the following are equivalent:*

- (1) X satisfies WAP.
- (2) X satisfies SAP.
- (3) For any n -Pfister space E , there is a natural number m and an element a of $U(R)$ such that $2^m E \cong 2^{m+n-1} \langle 1, a \rangle$.
- (4) Any n -Pfister space is stably linked to $2^n \langle 1 \rangle$.
- (5) Any two n -Pfister spaces are stably linked.

PROOF. We begin by noting that, in order to prove that X satisfies SAP, it suffices to prove that given a_1, \dots, a_n in $U(R)$. There is an element a in $U(R)$ such that $\bigcap_1^n W(a_i) = W(a)$. For, since the $W(a)$'s always constitute a subbase of the topology of X , this shows that they actually form a basis. Then, if Y is an arbitrary clopen subset of X , the clopen set $X - Y$ is a union of $W(a)$'s. Since $X - Y$ is a closed subset of the compact Hausdorff space X , it is itself compact, and thus there exist a_1, \dots, a_n in $U(R)$ with $X - Y = W(a_1) \cup W(a_2) \cup \dots \cup W(a_n)$. Now from the definition, $W(-a) = X - W(a)$, and so $Y = \bigcap_1^n W(-a_i) = W(a)$ for some a in $U(R)$, once the initial statement is proved.

(1) \Rightarrow (2).*) For any a_1, \dots, a_n in $U(R)$, let $Y = \bigcap_1^n W(a_i)$. Then Y is open and compact and since X satisfies WAP, there are elements b_1, \dots, b_k in $U(R)$ with $Y = W(b_1) \cup \dots \cup W(b_k)$. By repeating some of the $W(a_i)$'s or $W(b_i)$'s, if necessary, we may suppose

$$Y = \bigcap_1^n W(a_i) = \bigcup_1^n W(b_i), \quad a_i, b_i \text{ in } U(R).$$

Let $E = \bigotimes_1^n \langle 1, -a_i \rangle$, $F = \bigotimes_1^n \langle 1, b_i \rangle$. Then

- if σ is in Y , we have $\bar{\sigma}([E]) = 2^n$, $\bar{\sigma}([F]) = 0$,
- and
- if σ is in $X - Y$, we have $\bar{\sigma}([E]) = 0$, $\bar{\sigma}([F]) = 2^n$.

Hence $E \perp F$ and $2^n \langle 1 \rangle$ satisfy the conditions of Lemma 2.1, so that for some natural number m ,

$$2^m E \perp 2^m F \cong 2^{m+n} \langle 1 \rangle \perp 2^{m+n-1} \mathbf{H}.$$

Hence $2^m E \perp 2^m F$ contains a primitive isotropic element. By [1, Satz 2.7(c)] or Proposition 4.1(i), there is an element a in $U(R)$, such that $2^m E$ represents $-a$ while $2^m F$ represents a . If R is a field, this is quite clear, without any reference.

Let $2^m E = \langle c_1, \dots, c_{2^{m+n}} \rangle$. Since for σ in Y , we have $\bar{\sigma}([2^m E]) = 2^{m+n}$, necessarily $\sigma(c_i) = 1$ for $i = 1, 2, \dots, 2^{m+n}$. Hence by [8, Lemma 2.3(ii)], $\sigma(-a) = 1$, i.e., $\sigma(a) = -1$ for all σ in Y . A similar argument based on $2^m F$ shows that for all σ in $X - Y$ we have $\sigma(a) = 1$. Thus $Y = W(a)$.

*) In case R is a field, essentially the same proof of this implication was also found by T. C. CRAVEN.

(2) \Rightarrow (3). Let $E \cong \bigotimes_1^n \langle 1, a_i \rangle$ be an n -Pfister space and $Y = W(a_1) \cup W(a_2) \cup \dots \cup \dots \cup W(a_n)$. Then $\bar{\sigma}([E]) = 0$ for σ in Y and $\bar{\sigma}([E]) = 2^n$ for σ in $X - Y$. Since X satisfies SAP, there is an element a in $U(R)$ with $Y = W(a)$ and clearly the equation $\bar{\sigma}([E]) = \bar{\sigma}([2^{n-1}\langle 1, a \rangle])$ holds for all σ in X . Lemma 2.1 then proves (3).

The implication (3) \Rightarrow (4) is clear.

(4) \Rightarrow (5). If E and F denote two n -Pfister spaces, (4) guarantees the existence of elements a, b, c, d in $U(R)$, natural numbers m, m' , an $(m+n-1)$ -Pfister space G , and an $(m'+n-1)$ -Pfister space H , such that

$$2^m E \cong \langle 1, a \rangle \otimes G, \quad 2^{m+n} \langle 1 \rangle \cong \langle 1, b \rangle \otimes G,$$

$$2^{m'} F \cong \langle 1, c \rangle \otimes H, \quad 2^{m'+n} \langle 1 \rangle \cong \langle 1, d \rangle \otimes H.$$

Since for all σ in X , we have $\bar{\sigma}(2^{m+n} \langle 1 \rangle) = 2^{m+n}$, Lemma 2.1 proves the existence of natural numbers k, k' with

$$2^k G \cong 2^{m+n+k-1} \langle 1 \rangle, \quad 2^{k'} H \cong 2^{m'+n+k'-1} \langle 1 \rangle.$$

But then it is clear that $2^{m+k+m'+k'} E$ and $2^{m+k+m'+k'} F$ are linked, proving (5).

(5) \Rightarrow (2). As already remarked, it suffices to show that for a_1, \dots, a_n in $U(R)$, there is an a in $U(R)$ with $Y = \bigcap_1^n W(a_i) = W(a)$. Let $E = \bigotimes_1^n \langle 1, -a_i \rangle$ and $F = 2^n \langle 1 \rangle$. By (5), there is a natural number m , an $(n+m-1)$ -Pfister space G , and elements b, c in $U(R)$ with

$$2^m E \cong \langle 1, b \rangle \otimes G, \quad 2^m F \cong \langle 1, c \rangle \otimes G \cong 2^{m+n} \langle 1 \rangle.$$

Thus for all σ in X , we have $\bar{\sigma}([G]) = 2^{m+n-1}$. But

for σ in Y , we have $\bar{\sigma}([2^m E]) = 2^{m+n}$, and

for σ in $X - Y$, we have $\bar{\sigma}([2^m E]) = 0$.

It follows, therefore, that $\sigma(b) = 1$ for σ in Y , and $\sigma(b) = -1$ for σ in $X - Y$. Hence $Y = W(-b)$, and the proof of Theorem 2.2 is complete since the implication (2) \Rightarrow (1) is clear.

Remark 2.3. In [7, Def. 3.12] the notion of an abstract Witt ring for any abelian p -primary group G was introduced. Conditions (2)–(5) are still equivalent for a “small” [8, Rem. 3.17] abstract Witt ring, if G is a group of exponent 2. The n -Pfister spaces are replaced by elements $\prod_1^n (1 + \bar{g}_i)$ of the abstract Witt ring, the group $U(R)$ is replaced by G , the sets $W(a)$ by the sets $W(\bar{g})$ as defined in [8, Thm. 3.18 (ii)], and the isometries in (3), (4), and (5) by equalities in the small abstract Witt ring. However, Craven [2, Section 3] has shown that the implication (1) \Rightarrow (2) cannot hold in such abstract Witt rings, so that it is not surprising that this part of the proof uses facts about units of R represented by spaces.

3. Equivalence of HMP and SAP

Theorem 3.1. (cf. [5, Thm. C and 10, Satz 3.1]). *Let R be a connected semilocal ring with 2 in $U(R)$, then SAP is equivalent to HMP.*

PROOF. Let E be a space of rank n over R , and suppose $|\bar{\sigma}([E])| < n$ for all σ in X . Then the possible values of $\bar{\sigma}([E])$ are $-n+2, -n+4, \dots, n-4, n-2$. Let $Y_k = \{\sigma \text{ in } X \mid \bar{\sigma}([E]) = -n+2k\}$, $k=1, 2, \dots, n-1$. Clearly the Y_k are a partition of X and, of course, some Y_k may be empty. Now, the function $f_E: X \rightarrow \mathbf{Z}$ defined by $f_E(\sigma) = \bar{\sigma}([E])$ is continuous if \mathbf{Z} is given the discrete topology [8, Lemma 3.3 (iv)], so that each Y_k is a clopen subset of X . Since X satisfies SAP, there are elements b_1, b_2, \dots, b_{n-2} in $U(R)$ such that $W(b_1) = Y_1, W(b_2) = Y_1 \cup Y_2, \dots, W(b_{n-2}) = Y_1 \cup Y_2 \cup \dots \cup Y_{n-2}$. Now let $F \cong \langle b_1, \dots, b_{n-2} \rangle$ and σ be an element of Y_k , $k=1, 2, \dots, n-2$. Since Y_k is contained in $W(b_k), W(b_{k+1}), \dots, W(b_{n-2})$, but is disjoint from $W(b_1), W(b_2), \dots, W(b_{k-1})$, we have $\bar{\sigma}([F]) = (k-1) - (n-2-k+1) = -n+2k = \bar{\sigma}([E])$. If σ lies in Y_{n-1} , then since Y_{n-1} is disjoint from all $W(b_i)$, we still have $\bar{\sigma}([F]) = n-2 = \bar{\sigma}([E])$. Since the Y_i 's cover X , Lemma 2.1 guarantees the existence of a natural number m such that $2^m E \cong 2^m F \perp 2^m H$. Hence $2^m E$ is isotropic, and R satisfies HMP.

HMP \Rightarrow SAP. As at the beginning of the proof of Theorem 2.2, it suffices to prove that for any two elements a, b of $U(R)$, there is an element c in $U(R)$ with $W(a) \cap W(b) = W(c)$. Thus, for a, b in $U(R)$, let $E \cong \langle -1, a, b, ab \rangle$. It is easily verified that for all σ in X , we have $\bar{\sigma}([E]) = \pm 2$, so that by HMP, there is a natural number m such that $2^m E$ is isotropic. By multiplying by 2, if necessary, we assume $m \geq 1$. Thus $2^m \langle -1 \rangle \perp 2^m \langle a, b, ab \rangle$ is isotropic and so again by [1, Satz 2.7 (c)] or Proposition 4.1 (i), there is an element t in $U(R)$ such that $-t$ is represented by $2^m \langle -1 \rangle$ and t is represented by $2^m \langle a, b, ab \rangle$. Thus for some r_i in R , $t = \sum r_i^2$, so that by [8, Lemma 2.3 (ii)], $\sigma(t) = 1$ for all σ in X . Now t is represented by $2^m \langle a \rangle \perp \langle b \rangle \otimes \otimes 2^m \langle 1, a \rangle$ and by [1, Satz 2.7 (a)] or Prop. 4.1 (ii), there are units s and v in R such that $t = s + v$, s is represented by $2^m \langle a \rangle$, and v is represented by $\langle b \rangle \otimes 2^m \langle 1, a \rangle$. Setting $c = b^{-1}v$ we then have $t = s + bc$, with c represented by $2^m \langle 1, a \rangle$. Once more [8, Lemma 2.3 (ii)] shows $W(s) = W(a)$ and $W(c) \subset W(a)$. Now let σ lie in $W(c) \subset W(a)$ and suppose $\sigma(b) = 1$. Then $\sigma(bc) = -1 = \sigma(a)$, which again by [8, Lemma 2.3 (ii)] would force $\sigma(t) = -1$, a contradiction. Hence $W(c) \subset W(a) \cap W(b)$. On the other hand, if σ lies in $W(a) \cap W(b)$, then $\sigma(s) = -1$ and so, since $\sigma(t) = 1$, we again see by [8, Lemma 2.3 (ii)] that $\sigma(bc) = 1$, but this means $\sigma(c) = -1$, or $W(a) \cap W(b) \subset W(c)$, i.e., $W(a) \cap W(b) = W(c)$ as desired.

4. Representations of units

The following Proposition is a special case of [1, Satz 2.7], and is only given here for the reader's convenience.

Proposition 4.1. Let R be a connected semilocal ring with 2 in $U(R)$ and (E, B) a space over R . Let $E = E_1 \perp E_2$.

(i) *If the rank of E is ≥ 3 and e is a primitive isotropic element of E , there exists a primitive isotropic element e' of E such that $e' = e'_1 + e'_2$, e'_i in E_i , and $B(e'_i, e'_i)$ lies in $U(R)$.*

(ii) If, in addition, $\text{rank } E_i \cong 2$, $i=1, 2$, and e is an element of E with $B(e, e)$ in $U(R)$, then there exists an element e' in E with $e' = e'_1 + e'_2$, e'_i in E_i , such that $B(e, e) = B(e', e')$ and $B(e'_i, e'_i)$ lies in $U(R)$.

PROOF. We first prove (i) and (ii) in case R is a field, necessarily of characteristic different from 2. Since $E = E_1 \perp E_2$, we have $e = e_1 + e_2$, with e_i in E_i , and if both $B(e_1, e_1)$ and $B(e_2, e_2)$ are different from zero we may take $e' = e$, $e'_i = e_i$. Hence, by renumbering, if necessary, we may suppose $B(e_1, e_1) = 0$, i.e., E_1 is isotropic. Since E is nondegenerate, E_1 and E_2 are also, and so E_1 is universal. Since E_2 is nondegenerate and the characteristic of R is not 2, there is a vector e'_2 in E_2 with $B(e'_2, e'_2) \neq 0$. To complete (i), we simply choose e'_1 in E_1 with $B(e'_1, e'_1) = -B(e'_2, e'_2)$.

In case (ii), let $B(e_2, e_2) = B(e, e) = a \neq 0$. Suppose first that R is not the field of 3 elements. Since R contains at least three nonzero elements, R contains a nonzero element $b \neq \pm a$, and since E_1 is universal there is a vector \tilde{e}_1 in E_1 with $B(\tilde{e}_1, \tilde{e}_1) = b$. Now there is a nonzero element c in R , $c \neq \pm 1$, such that $b = ca$. Since $c = \left(\frac{c+1}{2}\right)^2 - \left(\frac{c-1}{2}\right)^2$, we have $c = u^2 - v^2$ with u, v nonzero elements of R . Thus

$b = (u^2 - v^2)a$, or $a = \frac{b}{u^2} + \frac{v^2}{u^2}a$, and setting $e'_1 = \frac{\tilde{e}_1}{u}$, $e'_2 = \frac{v}{u}e_2$, $e' = e'_1 + e'_2$, completes the proof if R is a field of more than three elements. If R is the field of three elements, then $2a + 2a = a$. But over R any space of rank 2 is universal [9, 62.1, p. 157] and thus there exist e'_i in E_i with $B(e'_i, e'_i) = 2a$.

Still supposing R a field, a classical theorem due to Witt [9, 42.17, p. 98], ensures that there is an isometry T of E such that $T(e) = e'$. Now there is a vector f in E orthogonal to e and such that $B(f, f) \neq 0$. For, if $\dim E = n$, then $\dim (\text{Re})^\perp = n - 1$, and if $(\text{Re})^\perp$ were totally isotropic then $n - 1 \leq \frac{n}{2}$, which forces $n \leq 2$. Thus, if

necessary, multiplying T on the right by the reflection $x \rightarrow x - \frac{2B(x, f)}{B(f, f)}f$, we may assume that $\det T = +1$.

To settle the case where R is a semilocal ring, let $O^+(E)$ denote the group of those isometries of E whose images on $E/m_j E$ have determinant $+1$ for all maximal ideals m_j , $j=1, \dots, k$, of R . By the above, there exist vectors \tilde{e}'_j in $E/m_j E$ with the desired projections on $E_i/m_j E_i$, $i=1, 2$, and elements T_j in $O^+(E/m_j E)$, such that $T_j(e + m_j E) = \tilde{e}'_j$. Now KNEBUSCH has proved that the natural map $O^+(E) \rightarrow \prod_1^k O^+(E/m_j E)$ is surjective [6, Satz 0.4]. Hence if T is an element in $O^+(E)$ mapping onto (T_1, T_2, \dots, T_k) in $\prod_1^k O^+(E/m_j E)$, it is clear that $e' = T(e)$ and the projections of e' into E_1 and E_2 , have the desired properties.

References

- [1] R. BAEZA and M. KNEBUSCH, Annulatoren von Pfisterformen über semilokalen Ringen, *Math. Z.*, **140** (1974), 41—62.
- [2] T. C. CRAVEN, The Boolean space of orderings of a field, *Trans. Amer. Math. Soc.* **209** (1975), 225—235.

- [3] R. ELMAN and T. Y. LAM, Pfister forms and K-theory of fields, *J. of Alg.* **23** (1972), 181—213.
- [4] R. ELMAN and T. Y. LAM, Quadratic forms over formally real fields and pythagorean fields, *Amer. J. Math.* **94** (1972), 1155—1194.
- [5] R. ELMAN, T. Y. LAM, and A. PRESTEL, On some Hasse principles over formally real fields, *Mat. Z.* **134** (1973), 291—301.
- [6] M. KNEBUSCH, Isometrien über semilokalen Ringen, *Mat. Z.* **108** (1969), 255—268.
- [7] M. KNEBUSCH, A. ROSENBERG and R. WARE, Structure of Witt rings and quotients of abelian group rings, *Amer. J. Math.* **94** (1972), 119—155.
- [8] M. KNEBUSCH, A. ROSENBERG and R. WARE, Signatures on semilocal rings, *J. of Alg.* **26** (1973), 208—250.
- [9] O. T. O'MEARA, Introduction to quadratic forms, *Berlin—Heidelberg—Göttingen*, 1963.
- [10] A. Prestel, Quadratische Semi-Ordnungen und quadratische Formen, *Mat. Z.* **133** (1973), 319—342.
- [11] A. ROY, Cancellation of quadratic forms over commutative rings, *J. of Alg.* **10** (1968), 286—298.

ALEX ROSENBERG
 DEPARTMENT OF MATHEMATICS
 CORNELL UNIVERSITY
 ITHACA, N.Y, 14853
 U.S.A.

ROGER WARE
 DEPARTMENT OF MATHEMATICS
 PENNSYLVANIA STATE UNIVERSITY
 UNIVERSITY PARK, PA, 16802
 U.S.A.

(Received January 22, 1975.)