

# A strong topology for the union of topological spaces

By Á. SZÁZ (Debrecen)

## 0. Introduction

If  $(X_i)_{i \in I}$  is a family of topological spaces then  $X = \bigcup_{i \in I} X_i$  is usually considered to be equipped with the finest topology for which the identity mappings of all the spaces  $X_i$  into  $X$  are continuous.

In this paper, we shall study a finer topology on  $X$ , namely we are considering  $X$  to be equipped with the coarsest topology for which the identity mappings of all the spaces  $X_i$  into  $X$  are open.

In connection with this topology, we prove here a few immediate results which are needed as a basis for some topological considerations in the multiplier extensions of admissible vector modules [8].

We are grateful to Z. BALOGH for many helpful discussions concerning this material. In particular, for Remark 2.2 and Example 3.5, and for pointing out a substantial simplification of the proof of Theorem 2.11.

## 1. Final topologies

*Notation 1.1.* Let  $(X_i)_{i \in I}$  be a family of topological spaces and  $Y$  be a set. Moreover, for each  $i \in I$ , let  $f_i$  be a mapping from  $X_i$  into  $Y$ , and suppose that  $Y = \bigcup_{i \in I} f_i(X_i)$ .

**Theorem 1.2.** *There exists a coarsest topology on  $Y$  for which all the mappings  $f_i$  are open. Moreover, this topology has the family  $\bigcup_{i \in I} f_i(\mathcal{T}_i)$ , where  $\mathcal{T}_i$  denotes the topology of  $X_i$ , as a subbase.*

PROOF. Obvious.

*Remark 1.3.* This theorem does not require that  $Y = \bigcup_{i \in I} f_i(X_i)$ .

**Theorem 1.4.** *The coarsest topology on  $Y$  for which all the mappings  $f_i$  are open is finer than the finest topology on  $Y$  for which all the mappings  $f_i$  are continuous.*

PROOF. Denote  $\mathcal{T}$  (resp.  $\mathcal{T}^*$ ) the coarsest (resp. finest) topology on  $Y$  for which all the mappings  $f_i$  are open (resp. continuous). If  $V \in \mathcal{T}^*$ , then we have

$$V = \bigcup_{i \in I} V \cap f_i(X_i) = \bigcup_{i \in I} f_i(f_i^{-1}(V)),$$

whence it is clear that  $V \in \mathcal{T}$ . Consequently,  $\mathcal{T}^* \subset \mathcal{T}$ .

**Corollary 1.5.** Equip  $Y$  with the coarsest topology for which all the mappings  $f_i$  are open. If  $\varphi$  is a mapping from  $Y$  into a topological space  $Z$  such that  $\varphi \circ f_i$  is continuous for all  $i \in I$ , then  $\varphi$  is continuous.

PROOF. This follows immediately from Theorem 1.4.

## 2. Union spaces

*Notation 2.1.* Let  $(X_i)_{i \in I}$  be a family of topological spaces,  $X = \bigcup_{i \in I} X_i$ , and equip  $X$  with the coarsest topology for which the identity mappings of all the spaces  $X_i$  into  $X$  are open.

*Remark 2.2.* By Theorem 1.2, the topology of  $X$  may also be described as the supremum of the topologies  $\mathcal{T}_i \cup \{X\}$ , where  $\mathcal{T}_i$  denotes the topology of  $X_i$ .

**Theorem 2.3.** For each  $i \in I$ ,  $X_i$  is an open subset of  $X$ , and the restriction to  $X_i$  of the topology of  $X$  is finer than the original topology of  $X_i$ .

PROOF. This is an immediate consequence of the definition of the topology of  $X$ .

**Theorem 2.4.** For some  $i \in I$ ,  $X_i$  is a subspace of  $X$  if and only if the identity mapping of  $X_i$  into  $X$  is continuous.

PROOF. This follows at once from Theorem 2.3.

**Theorem 2.5.** Suppose that there exists  $i_0 \in I$  such that  $X_{i_0} \subset X_i$  for all  $i \in I$ . Then the identity mapping of  $X_{i_0}$  into  $X$  is continuous if and only if the identity mapping of  $X_{i_0}$  into  $X_i$  is continuous for all  $i \in I$ .

PROOF. This is quite obvious from Theorem 1.2.

**Theorem 2.6.** Suppose that  $I$  is directed, and  $X_i \subset X_j$  if  $i \preceq j$ . Then the following conditions are equivalent:

- (i) the identity mappings of all the spaces  $X_i$  into  $X$  are continuous;
- (ii) the identity mapping of  $X_i$  into  $X_j$  is continuous and open if  $i \preceq j$ .

PROOF. Suppose that (i) holds. If  $i, j \in I$  such that  $i \preceq j$ , and  $U$  is an open subset of  $X_i$ , then  $U$  is also open in  $X$ , and thus  $U = U \cap X_j$  is also open in  $X_j$ . On the other hand, if  $V$  is open in  $X_j$ , then  $V$  is also open in  $X$ , and thus  $V \cap X_i$  is open in  $X_i$ . This proves (ii).

Suppose now that (ii) holds. If  $i, j \in I$  and  $V$  is open in  $X_j$ , then choosing  $k \in I$  such that  $i \preceq k$  and  $j \preceq k$ , we can infer that  $V$  is open in  $X_k$ , and hence that  $V \cap X_i$  is open in  $X_i$ . Hence, by Theorem 1.2, (i) follows.

*Remark 2.7.* The condition that the identity mapping of  $X_i$  into  $X_j$  is open if  $i \preceq j$  fails to hold in most of the applications.

*Remark 2.8.* If the identity mappings of all the spaces  $X_i$  into  $X$  are continuous, then by Theorem 1.4, the topology of  $X$  coincides with the finest topology on  $X$  for which the identity mappings of all the spaces  $X_i$  into  $X$  are continuous.

**Theorem 2.9.** *Suppose  $I$  is directed and  $X_i \subset X_j$  if  $i \cong j$ . Moreover, suppose that there exists a cofinal subset  $K$  of  $I$  such that  $X_k$  is Hausdorff for all  $k \in K$ . Then  $X$  is also Hausdorff.*

**PROOF.** If  $x_1, x_2 \in X$ , then by the assumptions there exists  $k \in K$  such that  $x_1, x_2 \in X_k$ . Thus, if  $x_1 \neq x_2$ , there are disjoint open subsets  $V_1$  and  $V_2$  of  $X_k$  such that  $x_1 \in V_1$  and  $x_2 \in V_2$ . Now, since  $V_1$  and  $V_2$  are also open in  $X$ , the proof is complete.

**Theorem 2.10.** *Let  $(x_\alpha)$  be a net in  $X$  and  $x \in X$ . Then the following conditions are equivalent:*

- (i)  $x \in \lim_{\alpha} x_\alpha$  in  $X$ ;
- (ii) for each  $i \in I$  such that  $x \in X_i$ , there exists  $\alpha_0$  such that  $\{x_\alpha\}_{\alpha \cong \alpha_0} \subset X_i$  and  $x \in \lim_{\alpha \cong \alpha_0} x_\alpha$  in  $X_i$ .

**PROOF.** Suppose first that we have (i). Let  $i \in I$  such that  $x \in X_i$ , and let  $V$  be an open subset of  $X_i$  such that  $x \in V$ . Then, since  $V$  is also open in  $X$ , there exists  $\alpha_0$  such that  $x_\alpha \in V$  for all  $\alpha \cong \alpha_0$ . Hence, it is clear that  $\{x_\alpha\}_{\alpha \cong \alpha_0} \subset X_i$  and  $x \in \lim_{\alpha \cong \alpha_0} x_\alpha$  in  $X_i$ .

Suppose now that (ii) holds. Let  $V$  be an open subset of  $X$  such that  $x \in V$ . Then, by Theorem 1.2, there is a finite subset  $\{i_k\}_{k=1}^n$  of  $I$ , and for each  $k=1, 2, \dots, n$  an open subset  $V_k$  of  $X_{i_k}$  such that  $x \in \bigcap_{k=1}^n V_k \subset V$ . Hence, by (ii), for each  $k=1, 2, \dots, n$ , there exists  $\alpha_k$  such that  $x_\alpha \in V_k$  for all  $\alpha \cong \alpha_k$ . Choosing  $\alpha_0$  such that  $\alpha_k \cong \alpha_0$  for all  $k=1, 2, \dots, n$ , we have  $x_\alpha \in V$  for all  $\alpha \cong \alpha_0$ . This proves (i).

**Theorem 2.11.** *Suppose that  $I$  is directed, and  $X_i \subset X_j$  if  $i \cong j$ . If  $C$  is a compact subset of  $X$ , then there exists  $i \in I$  such that  $C \subset X_i$ , and moreover, if  $C \subset X_j$ , then  $C$  is also compact in  $X_j$ .*

**PROOF.** Let  $C$  be a compact subset of  $X$ . Since  $\{X_i\}_{i \in I}$  is an open cover of  $C$ , there exists a finite subset  $\{i_k\}_{k=1}^n$  of  $I$  such that  $C \subset \bigcup_{k=1}^n X_{i_k}$ . Choosing  $i \in I$  such that  $i_k \cong i$  for all  $k=1, 2, \dots, n$ , we have  $C \subset X_i$ . Finally, if  $C \subset X_j$ , then by Theorem 1.2, it is clear that  $C$  is also compact in  $X_j$ .

**Corollary 2.12.** *Suppose that  $I$  is directed, and moreover, suppose that  $X_i \subset X_j$  and the identity mapping of  $X_i$  into  $X_j$  is continuous and open if  $i \cong j$ . Then  $C$  is a compact subset of  $X$  if and only if  $C$  is a compact subset of some  $X_i$ .*

**PROOF.** This follows immediately from Theorems 2.6 and 2.11.

*Remark 2.13.* Some of the results of this section can slightly be generalized according to the first section.

### 3. The case of topological vector spaces

*Notation 3.1.* Let  $X$  be a vector space over  $\mathbf{K}$  ( $=\mathbf{R}$  or  $\mathbf{C}$ ), and  $(X_i)_{i \in I}$  be a family of subspaces of  $X$  such that  $X = \bigcup_{i \in I} X_i$ . Suppose that each  $X_i$  is equipped with a vector topology, and consider  $X$  to be equipped with the coarsest topology for which the identity mappings of all the spaces  $X_i$  into  $X$  are open.

**Theorem 3.2.** (i) *If  $x, y \in X$  such that  $x + y \in X_i$  implies  $x, y \in X_i$ , then the addition in  $X$  is continuous at the point  $(x, y)$ .*

(ii) *If  $0 \neq \lambda \in \mathbf{K}$  and  $x \in X$ , then the scalar multiplication in  $X$  is continuous at the point  $(\lambda, x)$ .*

**PROOF.** Let  $x, y \in X$  as in (i), and suppose that  $W$  is an open subset of  $X$  such that  $x + y \in W$ . By Theorem 1.2, we may suppose that  $W$  is an open subset of some  $X_i$ . Then, by the assumption, we also have  $x, y \in X_i$ . Hence, since the addition in  $X_i$  is continuous, we can infer that there are open subsets  $U$  and  $V$  of  $X_i$  such that  $x \in U$ ,  $y \in V$  and  $U + V \subset W$ . Now, since  $U$  and  $V$  are also open in  $X$ , the proof of (i) is complete.

The assertion (ii) can be proved quite similarly, namely  $\lambda x \in X_i$  implies  $x \in X_i$  if  $\lambda \neq 0$ , and the scalar multiplication in  $X_i$  is continuous.

**Theorem 3.3.** *Suppose that  $I$  is a directed set, and moreover, suppose that  $X_i \subset X_j$  and the identity mapping of  $X_i$  into  $X_j$  is open if  $i \leq j$ . Then  $X$  is also a topological vector space.*

**PROOF.** The proof is similar to that of Theorem 3.2.

*Example 3.4.* The Euclidean space  $\mathbf{R}^2$  with its usual topology is a topological vector space over  $\mathbf{R}$  and  $\mathbf{R} \times \{0\}$  is a closed subspace of  $\mathbf{R}^2$ . Let  $\mathcal{T}$  be the coarsest topology on  $\mathbf{R}^2$  for which the identity mappings of the spaces  $\mathbf{R} \times \{0\}$  and  $\mathbf{R}^2$  into  $\mathbf{R}^2$  are open. Then  $(\mathbf{R}^2, \mathcal{T})$  is not a topological vector space.

To prove this, we show that the translation

$$(x, y) \rightarrow (x, y) + (0, -1)$$

is not continuous at the point  $(1, 1)$  for  $\mathcal{T}$ . For this, let  $(y_n)$  be a sequence in  $\mathbf{R} \setminus \{1\}$  such that  $\lim_n y_n = 1$ . Then, by Theorem 2.10, it is clear that  $\lim_n (1, y_n) = (1, 1)$  in  $(\mathbf{R}^2, \mathcal{T})$ , but the sequence  $((1, y_n) + (0, -1)) = ((1, y_n - 1))$  fails to converge in  $(\mathbf{R}^2, \mathcal{T})$ .

*Example 3.5.* The space  $l^2 = l^2(\{1, 2, \dots\})$  [5] with its usual topology is a topological vector space over  $\mathbf{C}$ , and

$$l_c^2 = \{(x_n) \in l^2 : x_n \neq 0 \text{ only for finitely many } n\}$$

is a subspace of  $l^2$  such that  $l_c^2$  and  $l^2 \setminus l_c^2$  are also dense in  $l^2$ . Let  $\mathcal{T}$  be the coarsest topology on  $l^2$  for which the identity mappings of the spaces  $l_c^2$  and  $l^2$  into  $l^2$  are open. Then  $(l^2, \mathcal{T})$  is not uniformizable.

To prove this, we show that  $(l^2, \mathcal{T})$  is not completely regular. For this, suppose indirectly that  $(l^2, \mathcal{T})$  is completely regular. Then, in particular, for

$0 \in I_c^2 \in \mathcal{T}$ , there exists a continuous function  $\varphi: (I^2, \mathcal{T}) \rightarrow [0, 1]$  such that  $\varphi(0) = 1$  and  $\varphi(x) = 0$  for all  $x \in I^2 \setminus I_c^2$ . Since  $0 \in \varphi^{-1}\left(\left[\frac{1}{2}, 1\right]\right) \in \mathcal{T}$  and  $I_c^2$  is considered as a topological subspace of  $I^2$ , by Theorem 1.2, there exists an open subset  $U$  of  $I^2$  such that  $0 \in U \cap I_c^2 \subset \varphi^{-1}\left(\left[\frac{1}{2}, 1\right]\right)$ . Since  $I^2 \setminus I_c^2$  is dense in  $I^2$ , we can choose an  $x_0 \in U \setminus I_c^2$ . Since  $x_0 \in \varphi^{-1}\left(\left[0, \frac{1}{2}\right]\right) \in \mathcal{T}$  and  $x_0 \notin I_c^2$ , again by Theorem 1.2, there exists an open subset  $V$  of  $I^2$  such that  $x_0 \in V \subset \varphi^{-1}\left(\left[0, \frac{1}{2}\right]\right)$ . Since  $U \cap V$  is a nonvoid open subset of  $I^2$  and  $I_c^2$  is also dense in  $I^2$ ,  $(U \cap V) \cap I_c^2 \neq \emptyset$ . Thus, we have  $\varphi^{-1}\left(\left[0, \frac{1}{2}\right]\right) \cap \varphi^{-1}\left(\left[\frac{1}{2}, 1\right]\right) \neq \emptyset$ , and this is a contradiction. Consequently,  $(I^2, \mathcal{T})$  can not be completely regular.

### References

- [1] J. M. ANTHONY, Topologies for quotient fields of commutative integral domains, *Pacific J. Math.* **36** (1971), 585—601.
- [2] N. BOURBAKI, *Topologie Générale*, Paris 1953.
- [3] J. DUGUNDJI, *Topology*, Boston 1966.
- [4] W. W. FAIRCHILD and C. I. TULCEA, *Topology*, Philadelphia 1971.
- [5] E. HEWITT and K. STROMBERG, *Real and Abstract Analysis*, Berlin 1969.
- [6] J. HORVÁTH, *Topological Vector Spaces and Distributions*, London 1966.
- [7] J. L. KELLEY, *General Topology*, New York 1955.
- [8] Á. SZÁZ, Hausdorff topologies for the multiplier extensions of admissible vector modules, *Publ. Math. (Debrecen)*, to appear.

(Received February 10, 1979)