

# Rees algebras and their varieties

By IVAN CHAJDA (Přerov) and JAROMÍR DUDA (Brno)

Following [3, 8], a subalgebra  $\mathfrak{B}$  of an algebra  $\mathfrak{A}$  is called a *Rees subalgebra* whenever there exists a congruence  $\theta$  on  $\mathfrak{A}$  such that  $\langle x, y \rangle \in \theta$  if and only if either  $x=y$  or both  $x, y$  are elements of  $\mathfrak{B}$ . In this case,  $\theta$  is called a *Rees congruence* on  $\mathfrak{A}$  induced by  $\mathfrak{B}$ . Rees congruences were introduced by D. REES [6] for semigroups; Rees congruences on lattices were used by G. SZÁSZ [7] for a construction of special kind.

The present paper deals with algebras having Rees subalgebras only. They are called *Rees algebras*. Rees algebras are closely related to *Hamiltonian algebras*, see [5], and thus also with algebras having the *Congruence Extension Property*, see [4]. For basic concepts and notations used in this paper see [2]. We will write  $A, B$ , etc. for the universes of the algebras  $\mathfrak{A}, \mathfrak{B}$ , etc.

## 1. Basic concepts

In the sense of [5], an algebra  $\mathfrak{A}$  is called *Hamiltonian* if and only if every subalgebra  $\mathfrak{B}$  of  $\mathfrak{A}$  is a block of the congruence  $\theta(B \times B)$ , where  $\theta(B \times B)$  denotes the smallest congruence on  $\mathfrak{A}$  collapsing  $B$ . In some particular cases, this congruence may be of the form  $\theta(B \times B) = B \times B \cup \omega_A$ , where  $\omega_A$  denotes the diagonal of  $A$ . However, this form of  $\theta(B \times B)$  is identical to that of [3]. In this way we introduce

*Definition 1.* An algebra  $\mathfrak{A}$  is called a *Rees algebra* if  $B \times B \cup \omega_A$  is a congruence on  $\mathfrak{A}$  for every subalgebra  $\mathfrak{B}$  of  $\mathfrak{A}$ . A variety  $\mathcal{V}$  is a *Rees variety* if each  $\mathfrak{A} \in \mathcal{V}$  is a Rees algebra.

Thus a Rees algebra is a special case of a Hamiltonian algebra. As it was proved by E. W. KISS [4], every variety of Hamiltonian algebras has the Congruence Extension Property (briefly CEP). For this reason recall

*Definition 2.* An algebra  $A$  has CEP if every congruence on an arbitrary subalgebra of  $\mathfrak{A}$  is a restriction of some congruence on  $\mathfrak{A}$ .

For Rees algebras, it will be useful to modify the definition of CEP in the following sense:

*Definition 3.* An algebra  $\mathfrak{A}$  satisfies *strong CEP* whenever  $\theta \cup \omega_A \in \text{Con } \mathfrak{A}$  for any congruence  $\theta$  on a subalgebra  $\mathfrak{B}$  of  $\mathfrak{A}$ .

A class of algebras  $\mathcal{C}$  has *CEP* (*strong CEP*) if each algebra in  $\mathcal{C}$  has CEP (strong CEP).

Apparently, strong CEP implies CEP and any algebra having strong CEP is a Rees one. A natural question arises: does there exist a Rees algebra which has not strong CEP? The answer is positive:

*Example 1.* Let  $\mathfrak{A}$  be a four element groupoid given by its multiplicative table:

$\cdot$	$a$	$b$	$c$	$z$
$a$	$c$	$c$	$a$	$a$
$b$	$c$	$c$	$a$	$c$
$c$	$c$	$c$	$b$	$b$
$z$	$a$	$c$	$b$	$z$

Clearly a subset  $\mathfrak{B} = \{a, b, c\}$  forms a subgroupoid of  $\mathfrak{A}$  and  $\mathfrak{A}$  has no other proper subgroupoid. Clearly  $\mathfrak{A}$  is a Rees groupoid since  $B \times B \cup \omega_A$  is a congruence on  $\mathfrak{A}$ . However,  $\mathfrak{A}$  has no strong CEP since the equivalence  $\theta$  with classes  $\{a, b\}, \{c\}$  is a congruence on  $\mathfrak{B}$  although  $\theta \cup \omega_A$  is not a congruence on  $\mathfrak{A}$  because

$$\langle a, b \rangle \in \theta, \quad \langle z, z \rangle \in \omega_A \quad \text{but} \quad \langle a, c \rangle = \langle a \cdot z, b \cdot z \rangle \notin \theta \cup \omega_A.$$

## 2. Rees algebras

**Theorem 1.** *For an algebra  $\mathfrak{A}$ , the following conditions are equivalent:*

- (1)  $\mathfrak{A}$  is a Rees algebra;
- (2) every subalgebra of  $\mathfrak{A}$  generated by two elements is Rees;
- (3) for every unary algebraic function  $\varphi$  over  $\mathfrak{A}$  and any two elements  $a, b$  of  $\mathfrak{A}$  we have either (i)  $\varphi(a) = \varphi(b)$ , or (ii)  $\varphi(a) = s(a, b)$ ,  $\varphi(b) = t(a, b)$  for some binary polynomials  $s$  and  $t$  of  $\mathfrak{A}$ .

**PROOF.** (1) $\Rightarrow$ (2) is trivial. Prove (2) $\Rightarrow$ (3): Let  $\varphi$  be a unary algebraic function over  $\mathfrak{A}$  and  $a, b \in \mathfrak{A}$ . Let  $\mathfrak{B}$  be a subalgebra of  $\mathfrak{A}$  generated by elements  $a, b$ . Then, using (2),  $B \times B \cup \omega_A$  is a congruence on  $\mathfrak{A}$  containing the pair  $\langle a, b \rangle$ . Consequently, also  $\langle \varphi(a), \varphi(b) \rangle \in B \times B \cup \omega_A$ , whence (3) is evident.

(3) $\Rightarrow$ (1): Let  $\mathfrak{B}$  be an arbitrary subalgebra of  $\mathfrak{A}$ . We have to show that the binary relation  $B \times B \cup \omega_A$  is a congruence on  $\mathfrak{A}$ . Evidently, it suffices to verify the Substitution Property only: choose  $\langle a, b \rangle \in B \times B \cup \omega_A$  and a unary algebraic function  $\varphi$  over  $\mathfrak{A}$ . Applying the hypothesis (3), we have either (i)  $\varphi(a) = \varphi(b)$  or (ii)  $\varphi(a) = s(a, b)$  and  $\varphi(b) = t(a, b)$  for some binary polynomials  $s, t$  of  $\mathfrak{A}$ . Clearly the first case means that  $\langle \varphi(a), \varphi(b) \rangle \in \omega_A$ . Suppose  $\varphi(a) \neq \varphi(b)$ . Then  $\langle a, b \rangle \in B \times B$  and thus by (ii),

$$\langle \varphi(a), \varphi(b) \rangle = \langle s(a, b), t(a, b) \rangle \in B \times B.$$

Summarizing, we get

$$\langle \varphi(a), \varphi(b) \rangle \in B \times B \cup \omega_A$$

which clearly implies the Substitution Property.

*Example 2.* (1) For a group  $\mathfrak{G}$ , the following two conditions are equivalent:

- (a)  $\mathfrak{G}$  is a Rees group;
- (b)  $\mathfrak{G} \cong Z_p$ , the cyclic group of prime order.

**PROOF.** (a) $\Rightarrow$ (b): Let  $\mathfrak{H}$  denote a subgroup of  $\mathfrak{G}$ . By the hypothesis,  $H \times H \cup \omega_A$  is a congruence on  $\mathfrak{G}$ . Now, the regularity of groups (see e.g. [1]) implies  $\mathfrak{H} = \mathfrak{G}$  or  $\mathfrak{H} = \{0\}$ , whence the conclusion  $\mathfrak{G} \cong Z_p$ ,  $p$  prime, follows. The converse implication (b) $\Rightarrow$ (a) is trivial since  $Z_p$ ,  $p$  prime, has no proper subgroup.

(2) For a semilattice  $\mathfrak{S}$ , the following two conditions are equivalent:

- (a)  $\mathfrak{S}$  is a Rees semilattice;
- (b) the length of  $\mathfrak{S}$  is at most 1.

**PROOF.** (b) $\Rightarrow$ (a) is trivial. Prove (a) $\Rightarrow$ (b). Suppose that the length of  $\mathfrak{S}$  is at least 2. Then  $\mathfrak{S}$  contains a three element chain  $a < b < c$ . Consequently,  $\{a, c\}$  is a subsemilattice of  $\mathfrak{S}$  and, by the hypothesis,  $\{a, c\}$  is a block of some congruence on  $\mathfrak{S}$ . However, any congruence block is convex, which is a contradiction.

(3) $\Rightarrow$ (2) holds also for lattices.

(4) Every unary algebra is a Rees algebra.

**PROOF.** This is a trivial consequence of Theorem 1 (3).

**Proposition 1.** *Being a Rees algebra is hereditary for subalgebras and homomorphic images.*

**PROOF.** The first assertion is evident. Further, let  $h: \mathfrak{A} \rightarrow \mathfrak{A}'$  be a homomorphism of  $\mathfrak{A}$  onto  $\mathfrak{A}'$  and  $\mathfrak{B}'$  be a subalgebra of  $\mathfrak{A}'$ . Put  $\mathfrak{B} = h^{-1}(\mathfrak{B}')$ . Then, by the hypothesis,  $B \times B \cup \omega_A$  is a congruence on  $\mathfrak{A}$ . It is routine to verify the formula

$$(h \times h)(B \times B \cup \omega_A) = B' \times B' \cup \omega_{A'}$$

which implies that  $B' \times B' \cup \omega_{A'}$  is a subalgebra of the square  $\mathfrak{A}' \times \mathfrak{A}'$ . Thus  $B' \times B' \cup \omega_{A'}$  is a congruence on  $\mathfrak{A}'$  which finishes the proof.

*Remark 1.* The class of all Rees algebras of the same type is not closed under forming direct products: consider the two-element semilattice (or lattice)  $\mathfrak{C}_2$ . Then the direct product  $\mathfrak{C}_2 \times \mathfrak{C}_2$  is not a Rees semilattice (or lattice, respectively).

### 3. Characterizations of Rees varieties

Although Rees algebras of the same type need not form a variety, there exist varieties of algebras whose all members are Rees algebras as we can see in Example 2 (4). It motivates our aim to characterize such varieties.

Let  $\mathcal{V}$  be a variety. An  $n$ -ary polynomial  $p$  of  $\mathcal{V}$  is called *essentially  $k$ -ary* ( $0 \leq k \leq n$ ) on the variety  $\mathcal{V}$ , if the polynomial  $p$  on the countably generated free algebra of  $\mathcal{V}$  depends on exactly  $k$  variables, see [1]. We say that  $\mathcal{V}$  is *at most unary* if every polynomial  $p$  of  $\mathcal{V}$  is either essentially unary or essentially nullary.

Further, denote by  $F_n(x_1, \dots, x_n)$  the free algebra of  $\mathcal{V}$  generated by the set of free generators  $\{x_1, \dots, x_n\}$ .

Recently, E. W. Kiss [4] has proved that any variety of Hamiltonian algebras has CEP. In this section we discuss the relationship between Rees varieties and strong CEP; moreover, we present different characterizations of Rees varieties.

**Theorem 2.** *For a variety  $\mathcal{V}$ , the following conditions are equivalent:*

- (1)  $\mathcal{V}$  is a Rees variety;
- (2)  $\mathcal{V}$  is at most unary;
- (3) every algebraic function over  $\mathfrak{A} \in \mathcal{V}$  is either a constant or a polynomial;
- (4) for every unary algebraic function  $\varphi$  over  $\mathfrak{A} \in \mathcal{V}$  and any two elements  $a, b$  of  $\mathfrak{A}$ , either (i)  $\varphi(a) = \varphi(b)$  or (ii) there exists a unary polynomial  $u$  of  $\mathcal{V}$  such that  $\varphi(a) = u(a)$ ,  $\varphi(b) = u(b)$ ;
- (5)  $\mathcal{V}$  has strong CEP.

**PROOF.** (1) $\Rightarrow$ (2): Let  $\mathcal{V}$  be a Rees variety. Consider the free algebra  $F_{2+n}(x, y, z_1, \dots, z_n)$ . Suppose  $f$  is an  $(n+1)$ -ary polynomial of  $\mathcal{V}$  depending on the first variable. Then  $\varphi(v) = f(v, z_1, \dots, z_n)$  is a unary algebraic function over  $F_{2+n}(x, y, z_1, \dots, z_n)$  and, by Theorem 1 (3), we have either (i)  $f(x, z_1, \dots, z_n) = f(y, z_1, \dots, z_n)$  or (ii)  $f(x, z_1, \dots, z_n) = s(x, y)$  and  $f(y, z_1, \dots, z_n) = t(x, y)$  for some binary polynomials  $s$  and  $t$  of  $\mathcal{V}$ . The case (i) is impossible since  $f$  depends on the first variable. The second case (ii) implies  $f(x, z_1, \dots, z_n) = t(x, x)$ , i.e.  $f$  is at most unary.

The implications (2) $\Rightarrow$ (3) $\Rightarrow$ (4) are evident. Prove (4) $\Rightarrow$ (5). Let  $\mathfrak{B}$  be an arbitrary subalgebra of an algebra  $\mathfrak{A} \in \mathcal{V}$ . Further, let  $\theta$  be a congruence on  $\mathfrak{B}$ . We have to show that the trivial extension  $\theta \cup \omega_{\mathfrak{A}}$  of  $\theta$  is a congruence on  $\mathfrak{A}$ . Let  $\varphi$  be a unary algebraic function over  $\mathfrak{A}$  and  $\langle a, b \rangle \in \theta \cup \omega_{\mathfrak{A}}$ . Suppose  $\varphi(a) \neq \varphi(b)$ . Then  $\langle a, b \rangle \in \theta$  and, by the hypothesis (4),

$$\varphi(a) = u(a), \quad \varphi(b) = u(b)$$

hold for some unary polynomial  $u$  of  $\mathcal{V}$ . Thus

$$\langle \varphi(a), \varphi(b) \rangle = \langle u(a), u(b) \rangle \in \theta$$

which completes the proof.

The implication (5) $\Rightarrow$ (1) is trivial.

*Remark 2.* Examples 2 (1), (2) and (3) show that condition (4) of Theorem 2 is weaker than part (3) of Theorem 1, i.e. a Rees algebra alone need not be unary. Further, we have proved that a Rees variety is equivalent to a variety having strong CEP although this need not be true for a single algebra, see Example 1. The following propositions point out which single algebras has strong CEP.

**Proposition 2.** *If  $\mathfrak{A} \times \mathfrak{A}$  is a Rees algebra then  $\mathfrak{A}$  has strong CEP (and thus CEP).*

**PROOF.** Let  $\theta$  be an arbitrary congruence on a subalgebra  $\mathfrak{B}$  of  $\mathfrak{A}$ . Evidently,  $\theta$  is a subalgebra of the square  $\mathfrak{B} \times \mathfrak{B}$  and so it is a subalgebra of the Rees algebra  $\mathfrak{A} \times \mathfrak{A}$ . The diagonal  $\omega_{\mathfrak{A}}$  has the same property and, moreover,  $\theta \cap \omega_{\mathfrak{A}} \neq \emptyset$ . Then, using the former result of [8; Proposition 2.1, p. 230],  $\theta \cup \omega_{\mathfrak{A}}$  is also a subalgebra of  $\mathfrak{A} \times \mathfrak{A}$  which completes the proof.

**Proposition 3.** *Any idempotent Rees algebra has strong CEP.*

PROOF. Let  $\mathfrak{B}$  be a subalgebra of an idempotent Rees algebra  $\mathfrak{A}$ . We have to show that  $\theta \cup \omega_{\mathfrak{A}} \in \text{Con } \mathfrak{A}$  for any congruence  $\theta$  on  $\mathfrak{B}$ . Take a unary algebraic function  $\varphi$  over  $\mathfrak{A}$  and suppose  $\langle a, b \rangle \in \theta \cup \omega_{\mathfrak{A}}$ . If  $\varphi(a) \neq \varphi(b)$  then  $\langle a, b \rangle \in \theta$  and, by Theorem 1 (3),

$$\varphi(a) = s(a, b), \quad \varphi(b) = t(a, b)$$

for some binary polynomials  $s, t$  of  $\mathfrak{A}$ . Since  $\mathfrak{A}$  is idempotent, we have

$$\langle s(a, b), a \rangle = \langle s(a, b), s(a, a) \rangle \in \theta,$$

$$\langle t(a, b), a \rangle = \langle t(a, b), t(a, a) \rangle \in \theta$$

and thus

$$\langle \varphi(a), \varphi(b) \rangle = \langle s(a, b), t(a, b) \rangle \in \theta.$$

Hence  $\langle \varphi(a), \varphi(b) \rangle \in \theta \cup \omega_{\mathfrak{A}}$ , which was to be proved.

The next theorem characterizes Rees varieties in terms of subalgebras:

**Theorem 3.** *For a variety  $\mathcal{V}$ , the following three conditions are equivalent:*

- (1)  $\mathcal{V}$  is a Rees variety;
- (2) subalgebras of each  $\mathfrak{A} \in \mathcal{V}$  are closed under set union;
- (3)  $F_n(x_1, \dots, x_n) = \bigcup_{i=1}^n F_1(x_i)$ .

PROOF. By Theorem 2 (2),  $\mathcal{V}$  is at most unary, thus evidently (1) $\Rightarrow$ (2). The implication (2) $\Rightarrow$ (3) is trivial. The condition (3) implies that  $\mathcal{V}$  is at most unary. Thus, by Theorem 2, also (3) $\Rightarrow$ (1) is true.

*Remark 3.* Example 2 (2) shows that the characterization (2) of Theorem 3 does not hold for a single algebra. Nevertheless, the following local version of Theorem 3 (2) can easily be verified:

**Proposition 4.** *If subalgebras of  $\mathfrak{A} \times \mathfrak{A}$  are closed under set union then  $\mathfrak{A}$  is a Rees algebra.*

PROOF. Let  $\mathfrak{B}$  be a subalgebra of  $\mathfrak{A}$ . Then  $\mathfrak{B} \times \mathfrak{B}$  as well as  $\omega_{\mathfrak{A}}$  are subalgebras of  $\mathfrak{A} \times \mathfrak{A}$ . By the hypothesis, also  $\mathfrak{B} \times \mathfrak{B} \cup \omega_{\mathfrak{A}}$  is a subalgebra of  $\mathfrak{A} \times \mathfrak{A}$ , i.e.  $\mathfrak{B} \times \mathfrak{B} \cup \omega_{\mathfrak{A}}$  is a congruence on  $\mathfrak{A}$  which was to be proved.

## References

- [1] B. CSÁKÁNY, Congruences and subalgebras, *Annales Univ. Sci. Budapest, sectio Math.* **18** (1975), 37—44.
- [2] G. GRÄTZER, Universal Algebra, 2-nd edition, *Springer* 1979.
- [3] P. KIRSCHENHOFER, R. F. TICHY, Polynomalgebren und Diskrepanz, *Sitzungsberichten der Österreich, Akad. Wiss. Mathem.-natur. Kl.*, **189** (1980), 321—339.
- [4] E. W. KISS, Each Hamiltonian variety has the Congruence Extension Property, *Alg. Univ.* **12** (1981), 395—398.
- [5] L. KLUKOVITS, Hamiltonian varieties of universal algebras, *Acta Sci. Math. (Szeged)*, **37** (1975), 11—15.
- [6] D. REES, On semigroups, *Proc. Cambridge Phil. Soc.* **36** (1940), 387—400.
- [7] G. SZÁSZ, Rees factor lattices, *Publ. Math. (Debrecen)* **15** (1968), 259—266.
- [8] R. F. TICHY, The Rees congruence in universal algebras, *Publications de l'Institut Math.* **29** (43), (1981), 229—239.

AUTHORS' ADDRESSES: IVAN CHAJDA  
TRÍDA LIDOVÝCH MILICÍ 22  
750 00 PŘEROV, CZECHOSLOVAKIA

JAROMÍR DUDA  
KROFTOVA 21  
616 00 BRNO, CZECHOSLOVAKIA

(Received May 13, 1983)