

## On an arithmetical property of $\sqrt{2}$

By I. JOÓ and M. JOÓ (Budapest)

*Deditated to Professor Paul Erdős for his 75th birthday*

In [2] it is proved that for every  $q \in (1, 2)$  excepted a set of first category there exists an expansion  $1 = \sum_i q^{-n_i}$  of 1 such that  $\limsup_{i \rightarrow \infty} (n_{i+1} - n_i) = \infty$ . Let  $\{y\}$  denote the fractional part of the real number  $y$ . We shall prove the following

**Theorem.**

- (i) For almost every  $x \in \mathbf{R}$ ,
- (ii) For every  $x \in \mathbf{R}$  excepted a set of first category we have

$$(1) \quad \inf_n \left( \{2^n x\} + \left\{ 2^n(1-x)/\sqrt{2} \right\} \right) = 0.$$

*Remarks.*

1. (1) means that in the dyadic expansions of  $x$  and of  $\frac{1-x}{\sqrt{2}}$  there are "arbitrarily long gaps at the same places" i.e. if

$$(2) \quad x = \sum_{k=0}^N \bar{a}_k 2^k + \sum_{k=1}^{\infty} a_k / 2^k, \quad \frac{1-x}{\sqrt{2}} = \sum_{k=0}^M \bar{b}_k 2^k + \sum_{k=1}^M b_k / 2^k,$$

$a_n, \bar{a}_n, b_n, \bar{b}_n \in \{0, 1\}$  then for every  $m \in \mathbf{N}$  there is an  $n_0$  such that

$$a_{n_0+1} = \dots = a_{n_0+m} = 0 = b_{n_0+1} = \dots = b_{n_0+m}.$$

2. Taking an  $x \in (0, 1)$  such that (1) holds we obtain from (2) that

$$1 = \sum 1/(\sqrt{2})^{n_i}$$

with  $\limsup_{i \rightarrow \infty} (n_{i+1} - n_i) = \infty$ .

PROOF. Let

$$A_\varepsilon := \left\{ x \in \mathbf{R} : \inf_n \left( \{2^n x\} + \{2^n(1-x)/\sqrt{2}\} \right) < \varepsilon \right\}, \quad \varepsilon > 0.$$

We prove that  $A_\varepsilon$  contains some dense open set and  $\text{mes}(\mathbf{R} \setminus A_\varepsilon) = 0$  for any fixed  $\varepsilon > 0$ . Denote by  $Z$  the set of integers. It is well known (see [1], p. 58) that the mod 1 distribution of the set  $\{r/\sqrt{2} : r \in Z\}$  is uniform in the following sense: for every  $0 < \varepsilon < 1$  there exists a natural number  $X(\varepsilon)$  such that any subsegment of  $[0,1]$  of length at least  $\varepsilon/4$  contains mod 1 a number  $r/\sqrt{2}$  of any sequence  $t/\sqrt{2}, \dots, (t+s)/\sqrt{2}$  if  $s > X(\varepsilon)$ ;  $t \in Z$ . Consider some  $r \in Z$  satisfying

$$(3) \quad ||| r/\sqrt{2} - \varepsilon/4 ||| < \varepsilon/8,$$

where  $|||y|||$  denotes the distance of the real number  $y$  from the nearest integer. Let  $n$  be a natural number, then we claim that  $I := [1 - \frac{1}{2^n}, 1 - \frac{1}{2^n} + (\varepsilon\sqrt{2}/8 \cdot 2^n)] \subset A_\varepsilon$ . Indeed, if  $y \in I$  then  $\{2^n y\} \leq \varepsilon\sqrt{2}/8 < \varepsilon/4$  and because of  $\frac{r}{\sqrt{2}} - \frac{\varepsilon}{8} \leq 2^n(1-y)/\sqrt{2} \leq r/\sqrt{2}$  we get  $\{2^n(1-y)/\sqrt{2}\} < \frac{3\varepsilon}{8}$ . We see that for any natural  $n$ , any segment of length  $\geq \frac{X(\varepsilon)+3}{2^n}$  contains a subsegment of length  $\varepsilon\sqrt{2}/8 \cdot 2^n$  belonging to  $A_\varepsilon$ . Indeed, we call  $r \in Z$  good if (3) holds. We know that the distance between two consecutive good numbers is not greater than  $X(\varepsilon) + 2$ , hence the distance of the corresponding left endpoints of the intervals  $I$  is  $\geq \frac{X(\varepsilon)+2}{2^n}$ , and so any segment of length  $\frac{X(\varepsilon)+2}{2^n} + \varepsilon\frac{\sqrt{2}}{8} \frac{1}{2^n} < \frac{X(\varepsilon)+3}{2^n}$  contains indeed a segment of length  $\varepsilon\frac{\sqrt{2}}{8} \frac{1}{2^n}$  from  $A_\varepsilon$ . Hence  $A_\varepsilon$  contains a dense open set i.e. the complement of  $\bigcap_{\varepsilon>0} A_\varepsilon$  is of first category, further the complement of  $A_\varepsilon$  has no density point consequently  $\text{mes}(\mathbf{R} \setminus A_\varepsilon) = 0$  i.e. the complement of  $\bigcap_{\varepsilon>0} A_\varepsilon$  has measure zero. The theorem is proved.  $\square$

The method of the proof works also for  $\sqrt[n]{2}$  ( $n > 2$ ).

*Problem.* What can we say about  $\sqrt{3}$ ?

## References

- [1] J. W. S. CASSELS, An introduction to diophantine approximation, *Cambridge University Press*, (1957).

- [2] I. JOÓ, On Riesz bases, *Annales Univ. Sci., Budapest, Sectio Math.* **31** (1988), 141-153.

I. JOÓ  
EÖTVÖS LORÁND UNIVERSITY  
CHAIR FOR ANALYSIS  
MÚZEUM KRT. 6-8.  
1088 BUDAPEST  
HUNGARY

M. JOÓ  
309 PAYNE ST  
AUBURN, AL 36830  
U.S.A.

*(Received April 25, 1989)*