

Year: 2022

Vol.: 100

Fasc.: 1-2

Title: A generalization of Skiba's problem

Author(s): Shuqin Dong, Tingting Qiu, Zhichao Gao, Long Miao and Jingyu He

Let H/K be any chief factor of a finite group G , p a prime, and j a nonnegative integer. We say that $G \in \mathfrak{S}_{p^j}^*$ if every chief factor H/K is a p -group or its p -part satisfies $|H/K|_p \leq p^j$. In this paper, we generalize Skiba's problem to the class $\mathfrak{S}_{p^j}^*$ of groups containing some non-solvable groups. We prove that $G \in \mathfrak{S}_{p^j}^*$ if and only if every maximal subgroup of a Sylow p -subgroup of G has a subnormal supplement in $\mathfrak{S}_{p^j}^*$, where $p > j$ or $p = j = 2$.

Address:

Shuqin Dong
School of Mathematical Sciences
Huaiyin Normal University
223300 Huaian
P. R. China

Address:

Tingting Qiu
Yancheng Biological Engineering
Higher Vocational Technology School
224000 Yancheng
P. R. China

Address:

Zhichao Gao
School of Mathematical Sciences
Yangzhou University
225002 Yangzhou
P. R. China

Address:

Long Miao
Basic Course Department
Jiangsu Vocational College
of Information Technology
214153 Wuxi
P. R. China

Address:

Jingyu He
Basic Course Department
Jiangsu Vocational College
of Information Technology
214153 Wuxi
P. R. China