

Year: 2022

Vol.: 101

Fasc.: 1-2

Title: On exponential Diophantine equations concerning Pythagorean triples

Author(s): Nobuhiro Terai and Yasutsugu Fujita

In 1956, Jeśmanowicz conjectured that the equation $(u^2 - v^2)^x + (2uv)^y = (u^2 + v^2)^z$ has only the positive integer solution $(x, y, z) = (2, 2, 2)$, where u and v are positive integers with $u > v$, $\gcd(u, v) = 1$ and $u \not\equiv v \pmod{2}$. Related to Jeśmanowicz' Conjecture, we propose the conjecture that the equation $x^2 + (2uv)^m = (u^2 + v^2)^n$ has exactly two positive integer solutions $(x, m, n) = (u - v, 1, 1)$, $(u^2 - v^2, 2, 2)$ except for the cases $(u, v) = (244, 231)$ and $3u^2 - 8uv + 3v^2 = -1$. We show that this conjecture is true for several cases. The proof is based on the deep results concerning (i) Generalized Lebesgue–Nagell equations, (ii) Generalized Fermat's equations, (iii) Primitive divisors of Lucas numbers, (iv) Linear forms in two logarithms.

Address:

Nobuhiro Terai
Division of Mathematical Sciences
Department of Integrated
Science and Technology
Faculty of Science and Technology
Oita University
700 Dannoharu
Oita 870–1192
Japan

Address:

Yasutsugu Fujita
Department of Mathematics
College of Industrial Technology
Nihon University
2-11-1 Shin-ei
Narashino, Chiba
Japan