

Title: Certain bilinear operators on power-weighted Morrey spaces

Author(s): Jing Zhou and Fayou Zhao

In this paper, we consider the boundedness properties of two classes of bilinear operators on Morrey spaces with power weights. The first operator is the bilinear maximal operator $T^*(f, g)(x) = \sup_j |T_j(f, g)(x)|$, where $T_j(f, g)$ is a bilinear operator with the kernel K_j satisfying the uniform estimate

$$|K_j(x, y_1, y_2)| \lesssim \frac{1}{(|x - y_1| + |x - y_2|)^{2n}},$$

where $x, y_1, y_2 \in \mathbb{R}^n$ with $x \neq y_k$ for some $k \in \{1, 2\}$. The second operator is $\mathcal{T}(f, g)$, which, being a bilinear operator, satisfies

$$|\mathcal{T}(f, g)(x)| \leq \int_{\mathbb{R}^n} \frac{|f(x - ty)g(x - y)|}{|y|^n} dy$$

for $x \in \mathbb{R}^n$ and $0 < |t| \leq 1$ such that $0 \notin \text{supp}(f(x - t \cdot)) \cap \text{supp}(g(x + \cdot))$. We obtain that these two operators are bounded operators from the product weighted Morrey spaces $L^{q, \lambda_1}(\mathbb{R}^n, |x|^\beta dx) \times L^{r, \lambda_2}(\mathbb{R}^n, |x|^\tau dx)$ to the weighted Morrey spaces $L^{p, \lambda}(\mathbb{R}^n, |x|^\alpha dx)$ with the assumption of the boundedness on Lebesgue spaces. As applications, we yield that many well-known bilinear operators are bounded on power-weighted Morrey spaces.

Address:

Jing Zhou
Department of Mathematics
Shanghai University
200444 Shanghai
P. R. China

Address:

Fayou Zhao
Department of Mathematics
Shanghai University
200444 Shanghai
P. R. China