

Title: The isoperimetric problem in Randers plane

Author(s): Arti Sahu Gangopadhyay, Ranadip Gangopadhyay, Hemangi Madhusudan Shah and Banktेशwar Tiwari

In 1947, Busemann observed that a Minkowski circle need not be a solution of the isoperimetric problem in a Minkowski plane. Li and Mo recently showed that the Euclidean circles centred at the origin in a unit ball with the Funk metric are solutions of the isoperimetric problem [9]. In this paper, we construct a class of Randers planes in which *any* Euclidean circle, centered at the origin in \mathbb{R}^2 , turns out to be a local minimum of the isoperimetric problem with respect to the various well-known volume forms in Finsler geometry. As a consequence, it turns out that the Euclidean circles centred at the origin are solutions of the isoperimetric problem in a Randers type Minkowski plane.

Address:

Arti Sahu Gangopadhyay
Centre for Interdisciplinary
Mathematical Sciences
Institute of Science
Banaras Hindu University
Varanasi-221005
India

Address:

Ranadip Gangopadhyay
Department of Mathematical Sciences
Indian Institute of Science Education
and Research
Mohali, Sector-81, Knowledge City
SAS Nagar Pin-140306
India

Address:

Hemangi Madhusudan Shah
Harish-Chandra Research Institute
A CI of Homi Bhaba National Institute
Chhatnag Road, Jhansi
Prayagraj-211019
India

Address:

Banktेशwar Tiwari
Centre for Interdisciplinary
Mathematical Sciences
Institute of Science
Banaras Hindu University
Varanasi-221005
India