

On the generalized Ramanujan–Nagell equation
 $D_1x^2 - D_2 = p^n$

By YASUTSUGU FUJITA (Narashino) and MAOHUA LE (Zhanjiang)

Abstract. Let D be a nonsquare positive integer, and let D_1, D_2 be positive integers such that $D_1D_2 = D$ and $\gcd(D_1, D_2) = 1$. Further, let p be an odd prime with $p \nmid D$. Denote by $N(D_1, D_2, p)$ the number of positive integer solutions (x, n) to the generalized Ramanujan–Nagell equation $D_1x^2 - D_2 = p^n$. In this paper, we prove $N(D_1, D_2, p) \leq 4$. Moreover, we generalize the case where $D_1 = 1$ to give the *exceptional* triple (D_1, D_2, p) , which satisfies

$$p^\nu = D_1a^2 + \eta, \quad D_2 = \frac{1}{D_1} \left(\frac{p^m - \eta}{2a} \right)^2 - p^m, \quad a, \nu, m \in \mathbb{N}, \quad \nu < m, \quad \eta \in \{1, -1\}.$$

In this case, it holds $N(D_1, D_2, p) \geq 3$. We prove that if (D_1, D_2, p) is non-exceptional, and the squarefree part of D_1 is at most 1001, then $N(D_1, D_2, p) \leq 3$.

YASUTSUGU FUJITA
DEPARTMENT OF MATHEMATICS
COLLEGE OF INDUSTRIAL TECHNOLOGY
NIHON UNIVERSITY
2-11-1 SHIN-EI
NARASHINO, CHIBA
JAPAN

MAOHUA LE
INSTITUTE OF MATHEMATICS
LINGNAN NORMAL COLLEGE
ZHANJIANG, GUANGDONG, 524048
CHINA

Mathematics Subject Classification: 11D45, 11D61, 33C90, 11J86.

Key words and phrases: exponential Diophantine equation, generalized Ramanujan–Nagell equation, hypergeometric function, Baker’s method.