

Methods for the calculation of values of a norm form

By A. BÉRCZES (Debrecen) and J. KÖDMÖN (Nyíregyháza)

Abstract. In this paper we analyse an important property of norm form functions, i.e. how hard is the computation of the function value if the arguments are given. We will present three algorithms for the calculation of the value of a norm form. Their complexity will be discussed and the running time of their implementations in MAPLE will be compared.

1. Introduction

The calculation of the values of a multivariate polynomial usually is a difficult task. Let $P(\underline{X})$ be a k -variable polynomial of total degree n . We can write

$$P(\underline{X}) = \sum_{j=1}^m a_j \underline{X}^{e_j},$$

where $\underline{e} = (e_1, \dots, e_k) \in \mathbb{Z}^k$, $0 \leq e_1 + \dots + e_k \leq n$ and $\underline{X}^{\underline{e}} := X_1^{e_1} X_2^{e_2} \dots X_k^{e_k}$.

Let $P(\underline{x})$ denote the value of the polynomial $P(\underline{X})$ at the point $\underline{x} = (x_1, \dots, x_k)$. The number of the terms of P is at most $\binom{n+1+k-1}{k} = \binom{n+k}{k} \approx \frac{2^{n+k}-2}{n+k-1} < 2^{n+k}$. For the naïve calculation of one term $O(n^2 \log^2 \mathbb{X})$ bit operations are needed, where $\mathbb{X} = \max\{|x_i|\}$ and the constant in O depends

Mathematics Subject Classification: 11Y40, 11Y16.

Key words and phrases: norm forms, complexity, algorithms.

Research was supported by the Grants F034981, T042985 and T038225, FKFP grant 3272-13066/201 and by the Netherlands Organization for Scientific Research.

only on $\max\{|a_j|\}$. Thus, the complexity of the determination of the substitution value is $O(2^{n+k}n^2 \log^2 \mathbb{X})$, which is exponential in $n+k$. For the complexities of basic algorithmic ingredients as integer arithmetic, matrix multiplication, etc. see [13].

There are also more efficient multiplication methods. The best known algorithm is due to SCHÖNHAGE and STRASSEN (see [12]), and it runs in $O(l \log l \log \log l)$ bit operations on l -bit numbers. However, this algorithm is worth using only for numbers having more than thousand digits. For smaller numbers the method of KARATSUBA–OFMAN (see [8]) may be useful.

If the vector $\underline{x} = (x_1, \dots, x_k)$ has rational integer components we can use an intelligent powering method such as the Right-Left Binary algorithm (see Algorithm 1.2.1 in [3]) which has a cost of $O(\log n \log^2 \mathbb{X})$ binary operations. In this case the complexity of the substitution is $O(2^{n+k+1} \log n \log^2 \mathbb{X})$.

So, determining the value $P(\underline{x})$ is in general a really difficult task.

There are, however, special polynomials where the calculation of the value is much easier. Let us take the following example

$$P(\underline{X}) = \begin{vmatrix} L_{1,1}(\underline{X}) & \cdots & L_{1,n}(\underline{X}) \\ \cdots & \cdots & \cdots \\ L_{n,1}(\underline{X}) & \cdots & L_{n,n}(\underline{X}) \end{vmatrix},$$

where $L_{i,j}(\underline{X}) = a_{ij1}X_1 + \cdots + a_{ijk}X_k$, $a_{ijk} \in \mathbb{Z}$; $1 \leq i, j \leq n$, $1 \leq k \leq n$ are linear forms. After expanding the determinant the result is a k -variable homogeneous polynomial of degree n . However, the value of this can be calculated in polynomial time in n since after determining the value of the linear forms $L_{i,j}(\underline{X})$ the calculation of the determinant has only polynomial complexity in n .

When the norm of a linear form with algebraic integer coefficients is to be computed, we also can do it by computing a determinant similar to the one above. The polynomial given this way is actually a norm form. Norm forms are playing an important role in the theory of diophantine equations (see [1], [5] and [10]). For computational results concerning norm form equations see [6] and the references given there.

2. Norm forms

Let θ be an algebraic integer of degree n , and denote by $T(X) \in \mathbb{Z}[X]$ its minimal polynomial over \mathbb{Q} . Put $K := \mathbb{Q}(\theta)$, and denote by $\sigma_i : K \rightarrow \mathbb{C}$ the distinct embeddings of K into the field of complex numbers. Further, for any $\alpha \in K$ denote by $\alpha^{(i)} := \sigma_i(\alpha)$ ($i = 1, \dots, n$) the field conjugates of α .

Let $\alpha_1, \alpha_2, \dots, \alpha_u \in K$ be \mathbb{Q} -linearly independent algebraic integers. Consider the linear form

$$L(\underline{X}) = \alpha_1 X_1 + \dots + \alpha_u X_u, \tag{1}$$

where $u \leq n$ and put

$$L^{(i)}(\underline{X}) = \alpha_1^{(i)} X_1 + \dots + \alpha_u^{(i)} X_u. \tag{2}$$

The polynomial

$$\text{Norm}_{\mathbb{K}/\mathbb{Q}}(L(\underline{X})) = \prod_{i=1}^n L^{(i)}(\underline{X})$$

is called a norm form. It is easily seen that $\text{Norm}_{\mathbb{K}/\mathbb{Q}}(L(\underline{X}))$ is a homogeneous polynomial of degree n , with integer coefficients.

Consider now the special case $\alpha_1 = 1, \alpha_2 = \theta, \dots, \alpha_u = \theta^{u-1}$. Each norm form can be transformed by a linear transformation with rational coefficients to a norm form of the above special type (see e.g. [1]), i.e. to a norm form

$$\mathcal{N}(\underline{X}) = \text{Norm}_{\mathbb{K}/\mathbb{Q}}(X_1 + \theta X_2 + \dots + \theta^{u-1} X_u). \tag{3}$$

In the rest of the paper we will restrict our investigations to the case when the norm form is of the shape (3). Further, due to some technical issues, in Sections 4 and 5 we also suppose that $1, \theta, \dots, \theta^{n-1}$ is a power integral basis for \mathbb{Z}_K , where \mathbb{Z}_K denotes the ring of integers of K .

In the following sections we consider and compare three different methods to compute values of $\mathcal{N}(\underline{X})$ at points $\underline{x} \in \mathbb{Z}^u$.

3. Calculation of $\mathcal{N}(\underline{x})$ by the definition

In the first method we use a floating point approximation $\tilde{\alpha}_j^{(i)}$ of each algebraic integer $\alpha_j^{(i)}$. Knowing that $\mathcal{N}(\underline{x})$ is an integer for $\underline{x} \in \mathbb{Z}^u$ it is enough to approximate $\alpha_j^{(i)}$ by $\tilde{\alpha}_j^{(i)}$ with such a precision that $|\mathcal{N}(\underline{x}) - \tilde{\mathcal{N}}(\underline{x})| < \frac{1}{2}$, where $\tilde{\mathcal{N}}(\underline{x}) := \prod_{i=1}^n (\sum_{j=1}^u \tilde{\alpha}_j^{(i)} x_j)$. Let us give an upper bound on the error of the calculation.

Lemma 1. *Using the above notation put $\lceil \alpha_j \rceil = \max \{ |\alpha_j^{(i)}| : 1 \leq i \leq n \}$, $A = \sum_{j=1}^u \lceil \alpha_j \rceil \geq 2$, $\Delta = \max \{ |\alpha_j^{(i)} - \tilde{\alpha}_j^{(i)}| : 1 \leq i \leq n, 1 \leq j \leq u \}$ and $\mathbb{X} = \max \{ |x_1|, \dots, |x_u| \}$. Then we have:*

$$\Delta_{\tilde{\mathcal{N}}(\underline{x})} = |\mathcal{N}(\underline{x}) - \tilde{\mathcal{N}}(\underline{x})| \leq nu\Delta(A\mathbb{X})^n,$$

provided that $\Delta \leq \frac{A}{nu\mathbb{X}}$.

PROOF. Let $\beta^{(i)} = \sum_{j=1}^u \alpha_j^{(i)} x_j$, $\tilde{\beta}^{(i)} = \sum_{j=1}^u \tilde{\alpha}_j^{(i)} x_j$ and $\tilde{\beta}^{(0)} = 1$. Then we have:

$$\begin{aligned} \Delta_{\tilde{\mathcal{N}}(\underline{x})} &= |\mathcal{N}(\underline{x}) - \tilde{\mathcal{N}}(\underline{x})| = \left| \prod_{i=1}^n \beta^{(i)} - \prod_{i=1}^n \tilde{\beta}^{(i)} \right| \\ &= \left| \sum_{j=0}^{n-1} \left(\prod_{i=0}^j \tilde{\beta}^{(i)} \prod_{i=j+1}^n \beta^{(i)} - \prod_{i=1}^{j+1} \tilde{\beta}^{(i)} \prod_{i=j+2}^n \beta^{(i)} \right) \right| \\ &= \left| \sum_{j=0}^{n-1} (\beta^{(j+1)} - \tilde{\beta}^{(j+1)}) \prod_{i=0}^j \tilde{\beta}^{(i)} \prod_{i=j+2}^n \beta^{(i)} \right|. \end{aligned}$$

Now using that

$$|\beta^{(i)}| \leq \mathbb{X} \sum_{j=1}^u \lceil \alpha_j \rceil = A\mathbb{X},$$

$$|\tilde{\beta}^{(i)}| \leq \mathbb{X} \sum_{j=1}^u (\lceil \alpha_j \rceil + \Delta) = \mathbb{X}(A + u\Delta)$$

$$|\beta^{(i)} - \tilde{\beta}^{(i)}| = \left| \sum_{j=1}^u (\alpha_j^{(i)} - \tilde{\alpha}_j^{(i)}) x_j \right| \leq \mathbb{X}u\Delta$$

and

$$\Delta \leq \frac{A}{nu\mathbb{X}}$$

we obtain that

$$\begin{aligned} \Delta_{\tilde{\mathcal{N}}(\underline{x})} &\leq u\Delta\mathbb{X}^n \sum_{j=0}^{n-1} (A + u\Delta)^j A^{n-j-1} = u\Delta\mathbb{X}^n A^{n-1} \sum_{j=0}^{n-1} \left(1 + \frac{u\Delta}{A}\right)^j \\ &= u\Delta\mathbb{X}^n A^{n-1} \frac{\left(1 + \frac{u\Delta}{A}\right)^n - 1}{\frac{u\Delta}{A}} = (A\mathbb{X})^n \left(\left(1 + \frac{u\Delta}{A}\right)^n - 1 \right) \\ &= (A\mathbb{X})^n \sum_{i=1}^n \binom{n}{i} \left(\frac{u\Delta}{A}\right)^i \leq (A\mathbb{X})^n \sum_{i=1}^n \frac{(nu\Delta)^i}{A^i i!} \\ &\leq (A\mathbb{X})^n \frac{nu\Delta}{2A} \sum_{i=1}^{\infty} \frac{(nu\Delta)^i}{A^i i!} \leq (A\mathbb{X})^n \frac{nu\Delta}{2A} \exp\left(\frac{nu\Delta}{A}\right) \\ &< \frac{enu\Delta}{2A} (A\mathbb{X})^n \leq nu\Delta (A\mathbb{X})^n. \end{aligned}$$

This concludes the proof of Lemma 1. □

The definition of the norm function implies that $\mathcal{N}(\underline{x}) \in \mathbb{Z}$ for $\underline{x} \in \mathbb{Z}^u$, thus the upper bound on the error of the calculation has to satisfy the inequality $\Delta_{\tilde{\mathcal{N}}(\underline{x})} < \frac{1}{2}$. From this it follows that we must choose the precision of the calculation as:

$$\Delta < \frac{1}{2nu(A\mathbb{X})^n}.$$

After determining the necessary precision of the approximations $\tilde{\alpha}_i^{(j)}$ to $\alpha_i^{(j)}$, we study the complexity of the calculation of $\mathcal{N}(\underline{x})$ using directly the definition of the norm form and floating point approximations to its coefficients. We have the following theorem:

Theorem 1. *The complexity of determination of the function value $\mathcal{N}(\underline{x})$ according to (3) is $O(n^6 + n^4 \log^2 \mathbb{X})$, where the constant in O depends only on A .*

PROOF. Let $m := n \log \mathbb{X} + \log(2nuA^n)$ and choose approximations $\tilde{\alpha}_i^{(j)}$ to each $\alpha_i^{(j)}$ having the following properties

- both the real and the imaginary part of $\tilde{\alpha}_i^{(j)}$ has a fractional part of at most $m + 1$ digits in its binary representation
- $|\Re(\alpha_i^{(j)}) - \Re(\tilde{\alpha}_i^{(j)})| < 2^{-m-1}$ and
- $|\Im(\alpha_i^{(j)}) - \Im(\tilde{\alpha}_i^{(j)})| < 2^{-m-1}$.

Then we have $\Delta < 2^{-m}$. Further, it is easily seen that

$$m \leq n \log \mathbb{X} + C_1 n u.$$

For any complex number z denote by $l(z)$ the maximum of the binary length of the real and imaginary part of the number. Since $u \leq n$ we get

$$l\left(\tilde{\alpha}_i^{(j)}\right) \leq \log \overline{|\alpha_i|} + n \log \mathbb{X} + C_1 n^2 = n \log \mathbb{X} + C_1 n^2 + C_2.$$

Now let us determine the complexity of the calculation of the values $L_j(\underline{x}) = \sum_{i=1}^u \tilde{\alpha}_i^{(j)} x_i$.

First of all we estimate the number of binary operations needed to calculate the appropriate approximations to the algebraic integers $\alpha_i^{(j)}$. According to Theorem 19.2 of [11] this can be done in

$$O(n^3 \log n + n^4 \log \mathbb{X} + C_1 n^5 + C_2 n^3) = O(n^5 + n^4 \log \mathbb{X}) \tag{4}$$

bit operations.

Now we turn to estimate the complexity of the remaining part of the calculation. The number of the binary operations needed for one multiplication is at most

$$C_3(n \log^2 \mathbb{X} + C_1 n^2 \log \mathbb{X} + C_2 \log \mathbb{X}),$$

hence u ($\leq n$) pieces of such multiplications demand less than

$$C_3 n^2 \log^2 \mathbb{X} + C_4 n^3 \log \mathbb{X} + C_5 n \log \mathbb{X}$$

binary operations.

Let $l := \max_j \{l(L_j)\} := \max_j \{l(L_j(\underline{x}))\} \leq \max_{i,j} \{l(\tilde{\alpha}_i^{(j)})\} + \log \mathbb{X} + n \leq (n + 1) \log \mathbb{X} + (C_1 + 1)n^2 + C_2$ be the maximal length of one $L_j(\underline{x})$.

Let us determine the complexity of the calculation of $\prod_{j=1}^n L_j(\underline{x})$. The number of the binary operations needed for it is obviously the sum of the elements of the sequence

$$l(L_1)l(L_2), [l(L_1) + l(L_2)]l(L_3), \dots, [l(L_1) + \dots + l(L_{n-1})]l(L_n),$$

times a constant. This can be estimated from above by the sum of the elements of the sequence $l^2, 2l^2, \dots, (n - 1)l^2$ times a constant. So, the complexity of the calculation of the complete product is:

$$C_6 \frac{n(n - 1)}{2} l^2 = C_6 \frac{n(n - 1)}{2} ((n + 1) \log \mathbb{X} + C_1 n^2 + C_2)^2 \tag{5}$$

$$\approx C_7 n^6 + C_8 n^4 \log^2 \mathbb{X}.$$

Further, the calculation of n pieces of L_j demands at most

$$C_3 n^3 \log^2 \mathbb{X} + C_4 n^4 \log \mathbb{X} + C_2 n^2 \log \mathbb{X} \tag{6}$$

binary operations.

Now (4), (5) and (6) imply that the complexity of the whole calculation is:

$$O(n^6 + n^4 \log^2 \mathbb{X}).$$

This concludes the proof of Theorem 1. □

4. Calculation of $\mathcal{N}(\underline{x})$ by matrix representation

If $\Omega = \{\omega_1, \dots, \omega_n\}$ is a \mathbb{Q} -basis of an algebraic number field K and if $\alpha \in K$, then multiplication by α is an endomorphism of the \mathbb{Q} -vector space K , and we can represent α by the matrix M_α of this endomorphism in the basis Ω . This matrix has in general rational entries. This representation is unique, and the map $\alpha \mapsto M_\alpha$ is a homomorphism from K to the algebra of $n \times n$ matrices over \mathbb{Q} . If $\Omega := \{1, \theta, \theta^2, \dots, \theta^{n-1}\}$ is a power integral basis of K and $\alpha = a_1 \omega_1 + \dots + a_n \omega_n$ where $a_i \in \mathbb{Z}$ for $1 \leq i \leq n$, then M_α has integral entries. It is well known (see e.g. [1]) that the norm of α is the determinant of the corresponding matrix M_α . Further details concerning matrix representation of algebraic numbers can be found in [3] and [1].

Theorem 2. *Let K be an algebraic number field with a power integral basis $1, \theta, \dots, \theta^{n-1}$, $\underline{x} \in \mathbb{Z}^u$ and define the norm form $\mathcal{N}(\underline{X})$ by (3). Then $\mathcal{N}(\underline{x})$ can be determined with integer arithmetic.*

In the proof the matrix representation of the linear form $L(\underline{X})$ will be used.

PROOF. Here an algorithm is given for the computation of $\mathcal{N}(\underline{x})$.

As earlier, we consider the number field $K = \mathbb{Q}(\theta)$ and we also recall the following notation. Let $u \leq n$ and put $\alpha_1 = 1, \alpha_2 = \theta, \dots, \alpha_u = \theta^{u-1}$. Consider the following linear form used earlier:

$$L(\underline{X}) = \alpha_1 X_1 + \dots + \alpha_u X_u = \sum_{i=1}^u \theta^{i-1} X_i.$$

For each $\underline{x} \in \mathbb{Z}^u$ the value $L(\underline{x})$ will be an element of the field K , thus we can compute its matrix representation in the basis Ω . However, we can first compute a matrix representation $\Lambda(\underline{X})$ of $L(\underline{X})$ on the basis Ω , which will have linear forms in \underline{X} as its entries, and which will have the property that $\Lambda(\underline{x})$ gives the matrix representation of the element $L(\underline{x})$ for each $\underline{x} \in \mathbb{Z}^u$. Now we compute this matrix representation of $L(\underline{X})$.

In order to determine the elements of this matrix consider the following products:

$$\theta^k L(\underline{X}) = \sum_{i=0}^{u-1} \theta^{i+k} X_i, \text{ where } 0 \leq k \leq n-1. \tag{7}$$

In (7) we shall substitute each θ^s with its standard representation in the base Ω . Put

$$\theta^s = \sum_{j=0}^{n-1} r_{s,j} \theta^j, \text{ where } 0 \leq s \leq 2n-2. \tag{8}$$

The standard representation of θ^s for $0 \leq s \leq n-1$ is trivial and we have $r_{s,j} = 1$ if $j = s$ and $r_{s,j} = 0$ otherwise. The standard representation of the powers θ^{n+k} ($0 \leq k \leq n-2$) can be determined from the coefficients of the minimal polynomial $T(X)$ of θ using the Newton recursion formulae. The method will be described later in Lemma 2.

Then by (7) and (8) we have

$$\theta^k L(\underline{X}) = \sum_{i=0}^{u-1} \left(\sum_{j=0}^{n-1} r_{i+k,j} \theta^j \right) X_i.$$

Changing the order of summation we get

$$\theta^k L(\underline{X}) = \sum_{j=0}^{n-1} \left(\sum_{i=0}^{u-1} r_{i+k,j} X_i \right) \theta^j = \sum_{j=0}^{n-1} F_{kj}(\underline{X}) \theta^j, \text{ where } 0 \leq k \leq n-1.$$

Thus the matrix representation of the linear form $L(\underline{X})$ is the matrix

$$\Lambda(\underline{X}) = \begin{pmatrix} F_{0,0}(\underline{X}) & \cdots & F_{0,n-1}(\underline{X}) \\ \cdots & \cdots & \cdots \\ F_{n-1,0}(\underline{X}) & \cdots & F_{n-1,n-1}(\underline{X}) \end{pmatrix},$$

where $F_{k,j}(\underline{X}) := \sum_{i=0}^{u-1} r_{i+k,j} X_i$; ($0 \leq k, j \leq n - 1$) are linear forms in u variables. Since $\mathcal{N}(\underline{x}) = \det(\Lambda(\underline{x}))$, the norm of $L(\underline{x})$ can be calculated by determining the determinant $\det(\Lambda(\underline{x}))$. So the calculation of the desired function value is possible with integer arithmetic and Theorem 2 is proved. \square

Lemma 2. *Let $T(X) = \sum_{s=0}^n t_s X^s \in \mathbb{Z}[X]$ be a monic irreducible polynomial over \mathbb{Q} of degree n . Denote by θ one of the roots of $T(X)$. Then $\theta^{n+i} = \sum_{j=0}^{n-1} r_{n+i,j} \theta^j$, $i \geq 0$, where $r_{n,j} = -t_j$ for $j = 0, \dots, n - 1$ and*

$$r_{n+i+1,j} = \begin{cases} r_{n+i,j-1} - t_j r_{n+i,n-1} & \text{if } j \geq 1 \\ -t_0 r_{n+i,n-1} & \text{if } j = 0. \end{cases}$$

PROOF. See Chapter 4.2.2 of [3]. \square

Now we summarize the main steps of the above construction, which enables us to determine the value $\mathcal{N}(\underline{x})$ from the given numbers $x_1, \dots, x_u \in \mathbb{Z}$ and the coefficients $t_0, \dots, t_n \in \mathbb{Z}$ of the polynomial $T(X)$:

1. The determination of the values $r_{i,j}$ via Newton’s formulae

In this step we calculate the values $r_{i,j}$ using Lemma 2.

This calculation can be done in advance since only the coefficients t_s of the polynomial $T(X)$ are needed.

The matrix representation of the linear form $L(\underline{X})$ will be the matrix $\Lambda(\underline{X})$ and its entries $F_{kj}(\underline{X})$ are u -variable linear forms having $r_{i+k,j}$ as their coefficients.

2. The determination of the values of the linear forms $F_{kj}(\underline{X})$

In this step the values of the n^2 forms $F_{kj}(\underline{X})$ are calculated at the vector $\underline{x} = (x_1, \dots, x_u) \in \mathbb{Z}^u$.

3. The calculation of $\det(\Lambda)$

Here, we calculate the determinant of the matrix $\Lambda(\underline{x})$ which has rational integer entries.

Theorem 3. *The complexity of the computation of $\mathcal{N}(\underline{x})$ with integer arithmetic, using the algorithm described above is $O(n^7 + n^6 \log \mathbb{X} + n^5 \log^2 \mathbb{X})$, where the constant in O depends only on the coefficients of $T(X)$.*

PROOF. We estimate the complexity of the three steps above. The whole complexity of the algorithm is given by the sum of these.

For the calculation of each $r_{i,j}$ only the coefficients t_i of the polynomial $T(X) = \sum_{i=0}^n t_i X^i$ are needed. Let $\mathbb{X} = \max\{|x_i|\}$ and $t = \max\{|t_i|\}$. The binary lengths of \mathbb{X} and t are $\log \mathbb{X}$ and $\log t$, respectively. For the calculation of the entries of the matrix of size $n \times (n-1)$ consisting of $r_{i,j}$ with the Newton's recursion formulae altogether $n^2 - 2n$ multiplications are needed, thus its complexity is

$$O(n^2). \quad (9)$$

The calculation of the values of the linear polynomials $F_{kj}(\underline{x})$ is possible with at most u multiplications. Since the length of $r_{i,j}$ is $C_1 n \log t = C_2 n$ and $u \leq n$, the complexity of the determination of the n^2 values is

$$O(n^4 \log \mathbb{X}). \quad (10)$$

The binary length of the numbers obtained is at most $C_3(C_2 n + \log \mathbb{X}) = C_4 n + C_3 \log \mathbb{X}$.

For the calculation of $\det(\Lambda(\underline{x}))$ the matrix $\Lambda(\underline{x})$ is reduced to triangular form by Gaussian-elimination and the product of the elements of the principal diagonal is taken. For this, however, operations in \mathbb{Q} are to be done. This problem can be solved by the multiplication with the common denominator of each of the eliminated lines. At the end the value of the determinant obtained this way should be divided by these multipliers, but the result will surely be a rational integer. Let l denote the maximum length of the elements of $\Lambda(\underline{x})$. Thus the length of the numbers obtained by the elimination of the j -th column will be jl . Since the worst case is when the denominators of the obtained numbers are coprime, the determination of the common denominator requires $O(jl(jl-1+1)) = O(j^2 l^2)$ binary operations. The steps of the elimination can be done altogether by $O(n^3)$ operations, so to transform our matrix into triangular form using integer arithmetic is possible in at most $O(n^5 l^2)$ binary operation. The maximum

length of the elements of the principal diagonal is nl , therefore $O(\frac{n^3-n^2}{2}l^2)$ binary operations are needed for the calculation of their product.

Since $l = C_4n + C_3 \log \mathbb{X}$, for the calculation of the determinant at most

$$O\left(n^5 + \frac{n^3 - n^2}{2}\right) (C_4n + C_3 \log \mathbb{X})^2 = O(n^7 + n^6 \log \mathbb{X} + n^5 \log^2 \mathbb{X}) \quad (11)$$

operations are needed.

The sum of (9), (10) and (11) is $O(n^7 + n^6 \log \mathbb{X} + n^5 \log^2 \mathbb{X})$ and Theorem 3 is proved. □

Remark 1. Since the calculation of step 1 can be done in advance, in practice only steps 2 and 3 need to be applied with the knowledge of the numbers x_1, \dots, x_u in order to calculate the value $\mathcal{N}(\underline{x})$. The complexity of steps 2 and 3 is essentially the one proved in Theorem 3.

5. Calculation of $\mathcal{N}(\underline{x})$ with modular arithmetic

Now another version of the algorithm described in the previous section will be presented, where the necessary operations are done by modular arithmetic.

Before describing the algorithm we outline the essence of the use of modular arithmetic.

In the present section $\text{mod } m$ denotes the remainder function such that $-\lceil \frac{m-1}{2} \rceil \leq x \text{ mod } m \leq \lfloor \frac{m}{2} \rfloor$, for every $x \in \mathbb{Z}$. Here $\lceil \cdot \rceil$ denotes the integer part function.

Let $m_1, \dots, m_v > 0$ be pairwise coprime integers and $M := m_1 m_2 \cdots m_v$. Then we assign to each $x \in \mathbb{Z}$ with $-\lceil \frac{M-1}{2} \rceil \leq x \leq \lfloor \frac{M}{2} \rfloor$ a v -tuple in the following way:

$$\varphi(x) = x^{(M)} = (x \text{ mod } m_1, \dots, x \text{ mod } m_v).$$

The map $\varphi : x \leftrightarrow x^{(M)}$ is a ring homomorphism from \mathbb{Z} to $\mathbb{Z}/(m_1) \times \cdots \times \mathbb{Z}/(m_v) \cong \mathbb{Z}/(m_1 \cdots m_v)$. The vector $x^{(M)}$ is called the modular representation of the rational integer x .

Let now \circ denote addition, subtraction or multiplication. If we obtain that

$$-\left\lfloor \frac{M-1}{2} \right\rfloor \leq x, y, x \circ y \leq \left\lfloor \frac{M}{2} \right\rfloor$$

then the result of the operation $x \circ y$ can be computed with modular arithmetic in the following steps:

1. Determination of $x^{(M)}$ and $y^{(M)}$ with Euclidean division.
2. Calculation of $x^{(M)} \circ y^{(M)}$ in the residue class rings.
3. Determination of $x \circ y = \varphi^{-1}(x^{(M)} \circ y^{(M)})$ with CRA (Chinese Remainder Algorithm).

The advantage of the procedure outlined above is that in step 2 the operations can be done with relatively small integers. Its disadvantage is that we have to determine in advance an upper bound for the possible values of $x \circ y$. Furthermore, the operation of division can only be done in a complicated way by determining modular inverses and the comparison of numbers is not possible at all.

In Chapter 4.3 of [9] there is an effective Chinese Remainder Algorithm, which applies recursion and certain parameters can be calculated in advance reducing the complexity of the algorithm.

Now let us present the modular arithmetic version of the algorithm described in the previous section.

Again let $\mathbb{X} := \max\{|x_i|\}$, $T(X) := \prod_{i=0}^n t_i X^i$ and $t := \max\{|t_i|\}$. Then $|\mathcal{N}(\underline{x})| \leq B := n^{2n} t^{n^2} \mathbb{X}^n$. We shall choose moduli $m_i := p_i^{k_i}$, $k_i \in \mathbb{Z}_{\geq 0}$ for $1 \leq i \leq v$, with distinct primes p_1, \dots, p_v such that $M := p_1^{k_1} p_2^{k_2} \dots p_v^{k_v} > 2B$. With such a choice we can guarantee

$$-\left\lfloor \frac{M-1}{2} \right\rfloor \leq \mathcal{N}(\underline{x}) \leq \left\lfloor \frac{M}{2} \right\rfloor.$$

Now we summarize in four steps the algorithm which enables us to determine the function value $\mathcal{N}(\underline{x})$ at $x_1, \dots, x_u \in \mathbb{Z}$ with the knowledge of the coefficients $t_0, \dots, t_n \in \mathbb{Z}$ of the polynomial T using modular arithmetic. Prime power moduli $p_1^{k_1}, \dots, p_v^{k_v}$ are calculated in advance having the properties $p_i \leq C$ for $i = 1, \dots, v$ with some constant C and $M := p_1^{k_1} p_2^{k_2} \dots p_v^{k_v} > 2B$. In order to be able to guarantee $M > 2B$ it is enough

to suppose $\prod_{p \leq C} p > 2B$, and since $\prod_{p \leq C} p > \sqrt{C}^{\pi(C) - \pi(\sqrt{C})}$ it is sufficient to have $\sqrt{C}^{\pi(C) - \pi(\sqrt{C})} > 2B$. Thus we suppose $C = C' \sqrt{\log(2B)}$.

The four main steps of our algorithm are the following:

1. Determination of the values $r_{i,j}$ with Newton's formulae

In this step the modular arithmetic is not yet used since on the one hand the maximum length of the numbers $r_{i,j}$ is $Cn \log t$, on the other hand these calculations can be performed in advance as only the coefficients t_i of the polynomial $T(X)$ are needed. Hence, the numbers $r_{i,j}$ are determined according to the description in the previous section.

2. Determination of the values of the linear polynomials $F_{kj}(\underline{x})$

Here the modular arithmetic is already applied. In this step the values of the linear polynomials $F_{kj}(\underline{x}) \pmod{p_1^{k_1}}, \dots, \pmod{p_v^{k_v}}$ are calculated at the vector $\underline{x} = (x_1, \dots, x_u) \in \mathbb{Z}^u$. This way in fact we determine the matrices $\Lambda^{(1)}(\underline{x}), \dots, \Lambda^{(v)}(\underline{x})$ which all have rational integer entries.

3. Calculation of $\det(\Lambda^{(M)}(\underline{x})) = \mathcal{N}^{(M)}(\underline{x})$

Now we determine the determinants of the matrices $\Lambda^{(1)}(\underline{x}), \dots, \Lambda^{(v)}(\underline{x})$. This will be the modular representation $(\mathcal{N}^{(1)}(\underline{x}), \dots, \mathcal{N}^{(v)}(\underline{x}))$ of the function value $\mathcal{N}(\underline{x})$.

4. Determination of the function value $\mathcal{N}(\underline{x})$

The value of $\mathcal{N}(\underline{x})$ is computed using the Chinese Remainder Algorithm from the modular representation $(\mathcal{N}^{(1)}(\underline{x}), \dots, \mathcal{N}^{(v)}(\underline{x}))$.

Theorem 4. *The complexity of the determination of $\mathcal{N}(\underline{x})$ with modular arithmetic is $O(n^7 + n^6 \log \mathbb{X} + n^2 \log^{3/2} \mathbb{X})$, where the constant in O depends only on the coefficients of $T(X)$.*

PROOF. We estimate the complexity of the four steps described above. The whole complexity of the algorithm is given by the sum of these. Let $t := \max\{|t_i|\}$ and $P := \max\{p_i^{k_i}\}$.

Since the numbers $r_{i,j}$ are computed in the same way as in the algorithm presented in the previous section, also the complexity of this step is

$$O(n^2). \tag{12}$$

In order to compute the values of the linear polynomials $F_{kj}(\underline{x})$ we have to determine first the remainders $x_i \pmod{p_1^{k_1}}, \dots, x_i \pmod{p_v^{k_v}}$, $i = 1, \dots, u$

and $r_{i,j} \bmod p_1^{k_1}, \dots, r_{i,j} \bmod p_v^{k_v}$, $i = n, \dots, n - 2$ and $j = 0, \dots, n - 1$. This needs $O(vn \log P \log \mathbb{X} + vn^3 \log P)$ binary operations. From now on the length of every number we encounter during the computation can be at most $\log P$. Since $u \leq n$, the complexity of the determination of the n^2 values is $O(vn^3 \log^2 P)$. Thus the whole complexity of the second step is

$$O(vn \log P \log \mathbb{X} + vn^3 \log^2 P). \tag{13}$$

In the third step the determinants of the matrices $\Lambda^{(M)}(\underline{x}) \in \mathbb{Z}^{n \times n}$ are calculated. The complexity of the calculation of the computation of each determinant is $O(n^5 \log^2 P)$. Therefore, the complexity of the third step is

$$O(vn^5 \log^2 P). \tag{14}$$

In the fourth step we compute the function value $\mathcal{N}(\underline{x})$ from its modular representation $(\mathcal{N}^{(1)}(\underline{x}), \dots, \mathcal{N}^{(v)}(\underline{x}))$ using the Chinese Remainder Algorithm. Now Algorithm 4.1 of [9] will be used. The input of this will be $(\mathcal{N}^{(1)}(\underline{x}), \dots, \mathcal{N}^{(v)}(\underline{x}))$, $p_1^{k_1}, \dots, p_v^{k_v}$, and q_2, \dots, q_v and s_2, \dots, s_v , where

$$q_i = \prod_{j=1}^{i-1} p_j^{k_j}, \quad s_i \equiv \prod_{j=1}^{i-1} s_j^{(i)} \bmod p_i^{k_i} \quad (i = 2, \dots, v),$$

where

$$s_j^{(i)} p_j^{k_j} + s_i^{(j)} p_i^{k_i} = 1.$$

The numbers $s_j^{(i)}$ and $s_i^{(j)}$ can be determined by $O(\log^2 P)$ binary operations by the extended Euclidean algorithm (see Theorem 3.8 in [9]). The determination of the numbers s_i demands $O((v - 1)^2 \log^2 P)$ binary operations, and for the calculation of the values q_i at most $O(\frac{1}{2}(v^2 - 3v + 2) \log^2 P)$ binary operations are needed. After these precomputations the CRA needs $O(v \log^2 P)$ binary operations. Therefore the overall complexity of the operations of the fourth step is

$$O(v^2 \log^2 P). \tag{15}$$

By (12), (13), (14) and (15) we can estimate the complexity of the algorithm by

$$O(v^2 n^5 \log^2 P + vn \log P \log \mathbb{X}). \tag{16}$$

Since $v \leq \frac{cC}{\log C}$ and $\log P \leq \log C$ we have $v \log P \leq cC$. Thus (16) takes the form

$$O(C^2 n^5 + Cn \log \mathbb{X}). \tag{17}$$

Further, since $C = C' \sqrt{\log(2B)}$ and $B = n^{2n} t^{n^2} \mathbb{X}^n$ we see that

$$C^2 \leq c_1 n^2 + c_2 n \log \mathbb{X}$$

and

$$C \leq c_3 n + c_4 n^{1/2} \log^{1/2} \mathbb{X} < c_5 n \log^{1/2} \mathbb{X}.$$

This shows that the complexity of the whole algorithm is at most

$$O(n^7 + n^6 \log \mathbb{X} + n^2 \log^{3/2} \mathbb{X})$$

and this was to be proved. □

Remark 2. The calculation of step 1 can be done in advance and the numbers s_i and q_i of step 4 can also be determined in advance.

6. Comparison of the algorithms

For the calculation of the function value $\mathcal{N}(\underline{x})$ three algorithms were discussed. The complexity of the algorithm using directly the definition of the norm form is $O(n^6 + n^4 \log^2 \mathbb{X})$. Its remarkable disadvantage is that we have to do operations with real numbers with precision $\frac{1}{2nu(A\mathbb{X})^n}$.

The complexity of the algorithm using the matrix representation of the linear form $L(\underline{X})$ is $O(n^7 + n^6 \log \mathbb{X} + n^5 \log^2 \mathbb{X})$. However, its important advantage is that it operates on integers.

The complexity of the algorithm using the matrix representation with modular arithmetic is $O(n^7 + n^6 \log \mathbb{X} + n^2 \log^{3/2} \mathbb{X})$.

The above three algorithms have been implemented in MAPLE *V Release 4* (see [2]). Let us refer to these procedures as *ALG1*, *ALG2* and *ALG3*, respectively. These algorithms were also compared with an algorithm *ALG4* which uses the built-in *Norm()* procedure of MAPLE. All the four algorithms were run with randomly generated polynomials and data. Every algorithm was run with the same parameters 50 times and

the running time was measured then later their average was taken. The following table shows the characteristic results of the test:

	$n = 5$	$n = 5$	$n = 5$	$n = 5$	$n = 5$	$n = 5$
	$u = 4$	$u = 4$	$u = 4$	$u = 4$	$u = 4$	$u = 4$
	$Et = 10$	$Et = 10$	$Et = 10$	$Et = 10$	$Et = 10$	$Et = 10$
	$Ex = 50$	$Ex = 70$	$Ex = 90$	$Ex = 110$	$Ex = 130$	$Ex = 150$
<i>ALG1</i>	0.2427	0.4892	0.5942	0.6128	1.233	1.244
<i>ALG2</i>	0.0130	0.0115	0.0126	0.0215	0.0181	0.0206
<i>ALG3</i>	0.0697	0.1032	0.1727	0.3099	0.2329	0.2897
<i>ALG4</i>	0.0221	0.0220	0.0262	0.0395	0.0506	0.0542

In the table n denotes the degree of the minimal polynomial, u denotes the number of the variables of the linear form, Et denotes the size of the coefficients of the minimal polynomial and Ex denotes the size of the substitution values.

The table shows that the fastest algorithm is *ALG2*, which uses integer arithmetic and applies the matrix representation of the norm form. *ALG1* runs in the first test with a precision of 400 digits and in the sixth with 900 digits precision, this is why its running time is the worst.

The above table also shows, how the increase of the substitution values effects the running time of our algorithms. In this respect the most sensitive is algorithm *ALG1* and the increase of the substitution values has the least effect on the running time of algorithm *ALG2*. However, the real slowing factor is, the increase of the degree. The following table shows the effect of this on the running time of the four algorithms:

	$n = 4$	$n = 5$	$n = 6$	$n = 7$	$n = 8$	$n = 9$
	$u = 4$	$u = 4$	$u = 5$	$u = 6$	$u = 7$	$u = 8$
	$Et = 10$	$Et = 10$	$Et = 10$	$Et = 10$	$Et = 10$	$Et = 10$
	$Ex = 100$	$Ex = 100$	$Ex = 100$	$Ex = 100$	$Ex = 100$	$Ex = 100$
<i>ALG1</i>	0.3996	0.6247	1.533	4.035	6.751	10.13
<i>ALG2</i>	0.0085	0.0160	0.0527	0.1161	0.2153	0.3811
<i>ALG3</i>	0.0757	0.1570	0.5232	1.137	2.245	4.316
<i>ALG4</i>	0.0245	0.0306	0.0581	0.0847	0.1383	0.2104

Here, in the sixth test *ALG1* calculates with a precision of 1500 digits, so its running time increases a significantly. Up to the third test the

running time of *ALG2* is the best. In the additional tests *ALG4* will become the fastest. It should be noted that algorithm *ALG4* uses the original *Norm()* procedure of MAPLE which was not implemented in MAPLE's own language but in *C*. This partly explains the differences in speed.

Algorithm *ALG3* using modular arithmetic is not fast enough because it uses the *chrem()* Chinese Remainder Algorithm of MAPLE in which the speedup due to precomputations (see Remark 2) cannot be utilized.

ACKNOWLEDGEMENTS. The authors would like to thank Professor A. PETHŐ for his generous help and the referee for his valuable remarks.

References

- [1] Z. I. BOREVICH and I. R. SHAFAREVICH, Number Theory, 2nd edn., *Academic Press, New York*, 1967.
- [2] B. W. CHAR, K. O. GEDDES, G. H. GONNET, B. L. LEONG, M. B. MONAGAN and S. M. WATT, Maple V Language reference manual, *Springer Verlag*, 1990.
- [3] H. COHEN, A course in computational algebraic number theory, *Springer Verlag*, 1993.
- [4] J.-H. EVERTSE and K. GYŐRY, Finiteness criteria for decomposable form equations, *Acta Arith.* **50** (1988), 357–379.
- [5] J.-H. EVERTSE, K. GYŐRY, C. L. STEWART and R. TIJDEMAN, *S*-unit equations and their applications, in: New Advances in Transcendence Theory, (A. Baker, ed.), *Cambridge Univ. Press*, 1988, 110–174.
- [6] I. GAÁL, An efficient algorithm for the explicit resolution of norm form equations, *Publ. Math. Debrecen* **56** (2000), 375–390.
- [7] P. HENRICI, Elements of numerical analysis, *Wiley and Sons, Inc.*, 1964.
- [8] A. KARATSUBA and YU. OFMAN, Multiplication of Many-Digital Numbers by Automatic Computers, *Doklady Akad. Nauk SSSR* **145** (1962), 293–294.
- [9] A. PETHŐ, Algebraische Algorithmen, *Vieweg*, 1999.
- [10] W. M. SCHMIDT, Diophantine Approximation, Lecture Notes in Mathematics, Vol. 785, *Springer Verlag, Berlin*, 1980.
- [11] A. SCHÖNHAGE, The fundamental theorem of algebra in terms of computational complexity (Preliminary report), unpublished manuscript.
- [12] A. SCHÖNHAGE and V. STRASSEN, Schnelle Multiplikation grosser Zahlen, *Computing* **7** (1971), 281–292.

[13] F. WINKLER, Polynomial algorithms in computer algebra, *Springer Verlag*, 1996.

A. BÉRCZES
INSTITUTE OF MATHEMATICS
UNIVERSITY OF DEBRECEN
NUMBER THEORY RESEARCH GROUP
HUNGARIAN ACADEMY OF SCIENCES AND
UNIVERSITY OF DEBRECEN
H-4010 DEBRECEN, P.O. BOX 12
HUNGARY

E-mail: berczesa@math.klte.hu

J. KÖDMÖN
FACULTY OF HEALTH COLLEGE
UNIVERSITY OF DEBRECEN
H-4400 NYÍREGYHÁZA, SÓSTÓI 2.
HUNGARY

E-mail: kodmonj@de-efk.hu

(Received January 21, 2003; revised June 25, 2003)