

## Some non-commutative products of distributions

By BRIAN FISHER (Leicester)

**Abstract.** Let  $g$  be a distribution and let  $g_n = (g * \delta_n)(x)$ , where  $\delta_n(x)$  is a certain sequence converging to the Dirac delta-function. The product  $f.g$  of two distributions  $f$  and  $g$  is defined to be the limit of the sequence  $\{fg_n\}$ , provided its limit  $h$  exists in the sense that

$$\lim_{n \rightarrow \infty} \langle f(x)g_n(x), \varphi(x) \rangle = \langle h(x), \varphi(x) \rangle$$

for all functions  $\varphi$  in  $\mathcal{D}$ . It is proved that

$$(\operatorname{sgn} x |x|^\lambda \ln^p |x|) \cdot (|x|^\mu \ln^q |x|) = \operatorname{sgn} x |x|^{\lambda+\mu} \ln^{p+q} |x|,$$

$$(|x|^\lambda \ln^p |x|) \cdot (\operatorname{sgn} x |x|^\mu \ln^q |x|) = \operatorname{sgn} x |x|^{\lambda+\mu} \ln^{p+q} |x|$$

for  $-2 < \lambda + \mu \leq -1$  and  $p, q = 0, 1, 2, \dots$

In the following, we let  $\mathcal{D}$  be the space of infinitely differentiable functions with compact support and let  $\mathcal{D}'$  be the space of distributions defined on  $\mathcal{D}$ . Now let  $\rho(x)$  be a function in  $\mathcal{D}$  having the following properties:

- (i)  $\rho(x) = 0$  for  $|x| \geq 1$ ,
- (ii)  $\rho(x) \geq 0$ ,
- (iii)  $\rho(x) = \rho(-x)$ ,
- (iv)  $\int_{-1}^1 \rho(x) dx = 1$ .

---

*Mathematics Subject Classification:* 46F10.

*Key words and phrases:* distribution, delta-function, product of distributions.

Putting  $\delta_n(x) = n\rho(nx)$  for  $n = 1, 2, \dots$ , it follows that  $\{\delta_n(x)\}$  is a regular sequence of infinitely differentiable functions converging to the Dirac delta-function  $\delta(x)$ .

If now  $f$  is an arbitrary distribution in  $\mathcal{D}'$ , we define

$$f_n(x) = (f * \delta_n)(x) = \langle f(t), \delta_n(x - t) \rangle$$

for  $n = 1, 2, \dots$ . It follows that  $\{f_n(x)\}$  is a regular sequence of infinitely differentiable functions converging to the distribution  $f(x)$ .

A first extension of the product of a distribution and an infinitely differentiable function is the following, see for example [2].

*Definition 1.* Let  $f$  and  $g$  be distributions in  $\mathcal{D}'$  for which on the interval  $(a, b)$ ,  $f$  is the  $k$ -th derivative of a locally summable function  $F$  in  $L^p(a, b)$  and  $g^{(k)}$  is a locally summable function in  $L^q(a, b)$  with  $1/p + 1/q = 1$ . Then the product  $fg = gf$  of  $f$  and  $g$  is defined on the interval  $(a, b)$  by

$$fg = \sum_{i=0}^k \binom{k}{i} (-1)^i [Fg^{(i)}]^{(k-i)}.$$

It follows easily that the products  $(|x|^\lambda \ln^p |x|)(|x|^\mu \ln^q |x|)$  and exist by Definition 1 and

$$(|x|^\lambda \ln^p |x|)(|x|^\mu \ln^q |x|) = |x|^{\lambda+\mu} \ln^{p+q} |x|, \quad (1)$$

$$(\operatorname{sgn} x |x|^\lambda \ln^p |x|)(|x|^\mu \ln^q |x|) = \operatorname{sgn} x |x|^{\lambda+\mu} \ln^{p+q} |x|, \quad (2)$$

$$(|x|^\lambda \ln^p |x|)(\operatorname{sgn} x |x|^\mu \ln^q |x|) = \operatorname{sgn} x |x|^{\lambda+\mu} \ln^{p+q} |x|, \quad (3)$$

$$(\operatorname{sgn} x |x|^\lambda \ln^p |x|)(\operatorname{sgn} x |x|^\mu \ln^q |x|) = |x|^{\lambda+\mu} \ln^{p+q} |x| \quad (4)$$

for  $\lambda + \mu > -1$  and  $p, q = 0, 1, 2, \dots$

The following definition for the non-commutative product of two distributions was given in [3] and generalizes Definition 1.

*Definition 2.* Let  $f$  and  $g$  be distributions in  $\mathcal{D}'$  and let  $g_n(x) = (g * \delta_n)(x)$ . We say that the product  $f.g$  of  $f$  and  $g$  exists and is equal to the distribution  $h$  on the interval  $(a, b)$  if

$$\lim_{n \rightarrow \infty} \langle f(x)g_n(x), \varphi(x) \rangle = \langle h(x), \varphi(x) \rangle$$

for all functions  $\varphi$  in  $\mathcal{D}$  with support contained in the interval  $(a, b)$ .

It was proved that if the product  $fg$  exists by Definition 1, then it exists by Definition 2 and  $fg = f.g$ .

The following theorem is easily proved.

**Theorem 1.** *Let  $f$  and  $g$  be distributions in  $\mathcal{D}'$  and suppose that the products  $f.g$  and  $f.g'$  (or  $f'.g$ ) exists. Then the product  $f'.g$  (or  $f.g'$ ) exists and*

$$(f.g)' = f'.g + f.g'. \tag{5}$$

The next theorem was proved in [4].

**Theorem 2.** *The product  $(x^r \ln^p |x|).(x^{-r-1} \ln^q |x|)$  exists and*

$$(x^r \ln^p |x|).(x^{-r-1} \ln^q |x|) = x^{-1} \ln^{p+q} |x| \tag{6}$$

for  $r = 0, \pm 1, \pm 2, \dots$  and  $p, q = 0, 1, 2, \dots$

We now prove the following theorem which generalizes equations (2), (3) and (6).

**Theorem 3.** *The products  $(\operatorname{sgn} x |x|^\lambda \ln^p |x|).( |x|^\mu \ln^q |x|)$  and  $( |x|^\lambda \ln^p |x|).( \operatorname{sgn} x |x|^\mu \ln^q |x|)$  exist and*

$$(\operatorname{sgn} x |x|^\lambda \ln^p |x|).( |x|^\mu \ln^q |x|) = \operatorname{sgn} x |x|^{\lambda+\mu} \ln^{p+q} |x|, \tag{7}$$

$$( |x|^\lambda \ln^p |x|).( \operatorname{sgn} x |x|^\mu \ln^q |x|) = \operatorname{sgn} x |x|^{\lambda+\mu} \ln^{p+q} |x| \tag{8}$$

for  $-2 < \lambda + \mu \leq -1$  and  $p, q = 0, 1, 2, \dots$

PROOF. We first of all prove equation (7) when  $\lambda > -1$ . Putting

$$( |x|^\mu \ln^q |x|)_n = ( |x|^\mu \ln^q |x|) * \delta_n(x),$$

we have

$$\int_{-a}^a (\operatorname{sgn} x |x|^\lambda \ln^p |x|)( |x|^\mu \ln^q |x|)_n dx = 0, \tag{9}$$

since the integrand is odd.

Further, if  $\psi$  is an arbitrary continuous function, we have

$$\int_{-a}^a (\operatorname{sgn} x |x|^\lambda \ln^p |x|)( |x|^\mu \ln^q |x|)_n x \psi(x) dx$$

$$= \int_{-a}^a |x|^{\lambda+1} \ln^p |x| (|x|^\mu \ln^q |x|)_n \psi(x) dx$$

and it follows that

$$\begin{aligned} & \lim_{n \rightarrow \infty} \int_{-a}^a (\operatorname{sgn} x |x|^\lambda \ln^p |x|) (|x|^\mu \ln^q |x|)_n x \psi(x) dx \\ &= \int_{-a}^a |x|^{\lambda+\mu+1} \ln^{p+q} |x| \psi(x) dx \\ &= \int_{-a}^a \operatorname{sgn} x |x|^{\lambda+\mu} \ln^{p+q} |x| x \psi(x) dx, \end{aligned} \tag{10}$$

since on using equation (1), the sequence  $\{|x|^{\lambda+1} \ln^p |x| (|x|^\mu \ln^q |x|)_n\}$  converges in the distributional sense to the locally summable function  $|x|^{\lambda+\mu+1} \ln^{p+q} |x|$ .

Now let  $\varphi$  be an arbitrary function in  $\mathcal{D}$  and choose  $a$  so that  $\operatorname{supp} \varphi \subset [-a, a]$ . By the mean value theorem, we have

$$\varphi(x) = \varphi(0) + x\varphi'(\xi x),$$

where  $0 < \xi < 1$ . Then

$$\begin{aligned} & \langle (\operatorname{sgn} x |x|^\lambda \ln^p |x|) (|x|^\mu \ln^q |x|)_n, \varphi(x) \rangle \\ &= \int_{-\infty}^{\infty} \operatorname{sgn} x |x|^\lambda \ln^p |x| (|x|^\mu \ln^q |x|)_n \varphi(x) dx \\ &= \varphi(0) \int_{-a}^a \operatorname{sgn} x |x|^\lambda \ln^p |x| (|x|^\mu \ln^q |x|)_n dx \\ &\quad + \int_{-a}^a \operatorname{sgn} x |x|^\lambda \ln^p |x| (|x|^\mu \ln^q |x|)_n x \varphi'(\xi x) dx. \end{aligned}$$

Using equations (9) and (10), it follows that

$$\begin{aligned} & \lim_{n \rightarrow \infty} \langle (\operatorname{sgn} x |x|^\lambda \ln^p |x|) (|x|^\mu \ln^q |x|)_n, \varphi(x) \rangle \\ &= \int_{-a}^a \operatorname{sgn} x |x|^{\lambda+\mu} \ln^{p+q} |x| x \varphi'(x) dx \\ &= \int_{-\infty}^{\infty} \operatorname{sgn} x |x|^{\lambda+\mu} \ln^{p+q} |x| [\varphi(x) - \varphi(0)] dx \\ &= \langle \operatorname{sgn} x |x|^{\lambda+\mu} \ln^{p+q} |x|, \varphi(x) \rangle \end{aligned}$$

for arbitrary  $\varphi$  in  $\mathcal{D}$ , proving equation (7) for  $\lambda > -1$ ,  $-2 < \lambda + \mu \leq -1$  and  $p, q = 0, 1, 2, \dots$

Equation (8) follows similarly for  $\lambda > -1$ ,  $-2 < \lambda + \mu \leq -1$  and  $p, q = 0, 1, 2, \dots$

When  $-2 < \lambda, \lambda + \mu \leq -1$ , we have from equation (1)

$$|x|^{\lambda+1}(|x|^\mu \ln^q |x|) = |x|^{\lambda+\mu+1} \ln^q |x| \quad (11)$$

for  $q = 0, 1, 2, \dots$ . Differentiating equation (11) we get

$$\begin{aligned} & (\lambda + 1)(\operatorname{sgn} x |x|^\lambda) \cdot (|x|^\mu \ln^q |x|) \\ & + \mu |x|^{\lambda+1} \cdot (\operatorname{sgn} x |x|^{\mu-1} \ln^q |x|) + q |x|^{\lambda+1} \cdot (\operatorname{sgn} x |x|^{\mu-1} \ln^{q-1} |x|) \quad (12) \\ & = (\lambda + \mu + 1) \operatorname{sgn} x |x|^{\lambda+\mu} \ln^q |x| + q \operatorname{sgn} x |x|^{\lambda+\mu} \ln^{q-1} |x|. \end{aligned}$$

Using Theorem 1 and equation (7), which has been proved for  $\lambda > -1$ , it follows that

$$(\operatorname{sgn} x |x|^\lambda) \cdot (|x|^\mu \ln^q |x|) = \operatorname{sgn} x |x|^{\lambda+\mu} \ln^q |x|. \quad (13)$$

Equation (7) therefore holds for  $-2 < \lambda, \lambda + \mu \leq -1$ ,  $p = 0$  and  $q = 0, 1, 2, \dots$

It follows similarly that

$$|x|^\lambda \cdot (\operatorname{sgn} x |x|^\mu \ln^q |x|) = \operatorname{sgn} x |x|^{\lambda+\mu} \ln^q |x|. \quad (14)$$

Equation (8) therefore holds for  $-2 < \lambda, \lambda + \mu \leq -1$ ,  $p = 0$  and  $q = 0, 1, 2, \dots$

Now suppose that equations (7) and (8) hold for some  $k$  with  $-k < \lambda$ ,  $-2 < \lambda + \mu \leq -1$  and  $p, q = 0, 1, 2, \dots$ . This is certainly true when  $k = 1$ . Also suppose that equations (7) and (8) hold for some  $p$  with  $-k - 1 \leq \lambda$ ,  $-2 < \lambda + \mu \leq -1$  and  $q = 0, 1, 2, \dots$ . This is also true when  $k = 1$  and  $p = 0$ . Then with  $-k - 1 < \lambda$  and  $-2 \leq \lambda + \mu < -1$ , it follows from equation (1) that

$$(|x|^{\lambda+1} \ln^{p+1} |x|)(|x|^\mu \ln^q |x|) = |x|^{\lambda+\mu+1} \ln^{p+q+1} |x|. \quad (15)$$

Differentiating equation (15), we get

$$\begin{aligned}
& (\lambda + 1)(\operatorname{sgn} x|x|^\lambda \ln^{p+1} |x|) \cdot (|x|^\mu \ln^q |x|) \\
& \quad + (p + 1)(\operatorname{sgn} x|x|^\lambda \ln^p |x|) \cdot (|x|^\mu \ln^q |x|) \\
& \quad + \mu(|x|^{\lambda+1} \ln^{p+1} |x|) \cdot (\operatorname{sgn} x|x|^{\mu-1} \ln^q |x|) \\
& \quad + q(|x|^{\lambda+1} \ln^{p+1} |x|) \cdot (\operatorname{sgn} x|x|^{\mu-1} \ln^{q-1} |x|) \\
& = (\lambda + \mu + 1) \operatorname{sgn} x|x|^{\lambda+\mu} \ln^{p+q+1} |x| + (p + q + 1) \operatorname{sgn} x|x|^{\lambda+\mu} \ln^{p+q} |x|.
\end{aligned}$$

Using our assumptions and Theorem 1, it follows that

$$(\operatorname{sgn} x|x|^\lambda \ln^{p+1} |x|) \cdot (|x|^\mu \ln^q |x|) = \operatorname{sgn} x|x|^{\lambda+\mu} \ln^{p+q} |x|$$

giving equation (7) for  $p + 1$  and  $-k - 1 < \lambda$ .

Similarly, differentiation of the equation

$$(\operatorname{sgn} x|x|^{\lambda+1} \ln^{p+1} |x|)(\operatorname{sgn} x|x|^\mu \ln^q |x|) = |x|^{\lambda+\mu+1} \ln^{p+q+1} |x|,$$

using our assumptions and Theorem 1, it follows that

$$(|x|^\lambda \ln^{p+1} |x|) \cdot (\operatorname{sgn} x|x|^\mu \ln^q |x|) = \operatorname{sgn} x|x|^{\lambda+\mu} \ln^{p+q} |x|$$

giving equation (8) for  $p + 1$  and  $-k - 1 < \lambda$ .

Equations (7) and (8) now follow by induction for all  $\lambda, \mu$ , with  $-2 < \lambda + \mu \leq 1$  and  $p, q = 0, 1, 2, \dots$ , completing the proof of the theorem.  $\square$

We finally consider what happens if  $\lambda + \mu \leq 2$ . With  $\lambda > -1$ , the sequence  $\{(|x|^\mu \ln^q |x|)_n\}$  will converge to the distribution  $|x|^\mu \ln^q |x|$  and the integral

$$\int_{-n}^n \operatorname{sgn} x|x|^{\lambda+\mu} \ln |x|^{p+q} x \psi(x) dx$$

in equation (10) will in general be divergent. This means that

$$\lim_{n \rightarrow \infty} \langle (\operatorname{sgn} x|x|^\lambda \ln^p |x|)(|x|^\mu \ln^q |x|)_n, \varphi(x) \rangle$$

cannot exist for all functions  $\varphi$ . The product  $(\operatorname{sgn} x|x|^\lambda \ln^p |x|) \cdot (|x|^\mu \ln^q |x|)$  will therefore not exist in this case.

Similarly, the product  $(|x|^\lambda \ln^p |x|) \cdot (\operatorname{sgn} x|x|^\mu \ln^q |x|)$  will not exist if  $\lambda + \mu < -2$ .

ACKNOWLEDGEMENT. The author would like to thank the referee for his suggestion in the improvement of this paper.

### References

- [1] J. G. VAN DER CORPUT, Introduction to the neutrix calculus, *J. Analyse Math.* **7** (1959–60), 291–398.
- [2] B. FISHER, The product of distributions, *Quart. J. Math. Oxford (2)*, **22** (1971), 291–298.
- [3] B. FISHER, On defining the product of distributions, *Math. Nachr.* **99** (1980), 230–240.
- [4] B. FISHER and S. YAKUBOVICH, Results on the non-commutative neutrix product of distributions, *Radovi Mat.* **11**(1) (2001), 85–93.
- [5] I. M. GEL'FAND and G. E. SHILOV, Generalized Functions, Vol. I, *Academic Press*, 1964.

BRIAN FISHER  
DEPARTMENT OF MATHEMATICS  
UNIVERSITY OF LEICESTER  
LEICESTER, LE1 7RH  
ENGLAND

*E-mail:* fbr@le.ac.uk

*(Received March 14, 2002; revised July 30, 2003)*