

On the Lebesgue property in uniform spaces.

To the memory of Professor Tibor Szele.

By KIYOSHI ISÉKI in Kobé.

This paper is mainly concerned with what might be called the finite Lebesgue property. A. A. MONTEIRO and M. M. PEIXOTO ([7], [8]) have introduced the notion of Lebesgue number in metric space and have discussed the relations between the Lebesgue number and uniform continuity of continuous functions on it.¹⁾

As far as possible, we use the notations and notions of P. SAMUEL [9]. All spaces considered will be assumed to be separated uniform spaces.

The following definitions and notations are used throughout the paper.

Let E be a separated uniform space. A *covering* of E is a family of open sets whose union is E . A covering is called *binary* if it consists of two open sets or *finite* if it consists of a finite family of open sets.

Let $\mathfrak{S}$ be a filter of the diagonal of $E \times E$ (a filter of surroundings of E). If $A \subset E$ and $V \in \mathfrak{S}$, we denote by $V(A)$ the image of the set $(E \times A) \cap V$ by the projection of $E \times E$ onto the first factor E .

We shall say that a (finite) covering $\mathfrak{F} = \{O_\alpha\}$ of E has (finite) *Lebesgue property* if there is a surrounding V of $\mathfrak{S}$ such that, for every x of E , we can find an open set O_α satisfying $V(x) \subset O_\alpha$.

We shall say that a uniform space E has the (finite) *Lebesgue property*, if any (finite) covering of E has the (finite) Lebesgue property.

Lemma. *If any binary covering of the uniform space E has the Lebesgue property, then E has the finite Lebesgue property.*

PROOF. We shall prove first that E is normal.

Let F_1, F_2 be two disjoint closed sets, and $O_i = E - F_i$ ($i = 1, 2$). Then $\{O_1, O_2\}$ is a binary covering of E . Therefore, since $\{O_1, O_2\}$ has the Lebesgue property, there is a surrounding $V \in \mathfrak{S}$ such that $V(x) \subset O_1$, or $V(x) \subset O_2$ for any x of E . For the surrounding V , we can find a symmetric surrounding W such that $W \circ W \subset V$. We shall show that $W(F_1) \cap W(F_2) = \emptyset$. To prove

¹⁾ Generalizations of these results are treated in my papers [3], [4] and [5].

it, suppose that $W(F_1) \cap W(F_2) \neq \emptyset$, and let x be a point of this set. Then there exist two points a_1, a_2 such that $(x, a_1) \in W$, $(x, a_2) \in W$ and $a_i \in F_i$ ($i=1, 2$). Therefore $(a_1, a_2) \in W \circ W \subset V$. On the other hand, since $a_1 \in F_1$, $a_1 \notin O_1$ and we have $V(a_1) \subset O_2$. Since $a_2 \in F_2$, $a_2 \notin O_2$ and we have $a_2 \notin V(a_1)$. This shows that $(a_1, a_2) \notin V$, which gives a contradiction. Thus we have proved the normality of E .

Now we shall prove that E has the finite Lebesgue property. Let $\{O_i\}$ ($i=1, 2, \dots, n$) be a finite covering of E . Since E is normal, $\{O_i\}$ is shrinkable and we can find a covering $\{G_i\}$ such that $G_i \subset \bar{G}_i \subset O_i$ ($i=1, 2, \dots, n$). Let $H_i = E - \bar{G}_i$, then, for each i , $\{O_i, H_i\}$ is a binary covering of E . Hence, by the assumptions of our Lemma, it has the Lebesgue property. There exists a surrounding $V_i \in \mathfrak{S}$ such that $V_i(x) \subset H_i$ or $V_i(x) \subset O_i$ for any x of E . Let $V = \bigcap_{i=1}^n V_i$, then, if $V(x) \subset H_i$ ($i=1, 2, \dots, n$), we have

$$V(x) \subset \bigcap_{i=1}^n H_i = \bigcap_{i=1}^n (E - \bar{G}_i) = E - \bigcap_{i=1}^n \bar{G}_i = \text{empty},$$

which contradicts the fact that $V(x)$ is not empty. Hence there is an index i such that $V(x) \subset O_i$, and thus the proof is completed.

Theorem 1. *A uniform space E has the finite Lebesgue property, if and only if E is normal and every bounded continuous function on E is uniformly continuous.*

PROOF. Suppose that E has the finite Lebesgue property. By our Lemma E is normal. Let $f(x)$ be a bounded continuous function and ε a given positive number. Since $f(x)$ is bounded, the range of $f(x)$ is contained in a finite interval $[-\alpha, \alpha]$. Let $\{I_i\}$ ($i=1, 2, \dots, n$) be a finite covering of $[-\alpha, \alpha]$ consisting of open intervals with length less than ε . Putting $O_i = f^{-1}(I_i)$ ($i=1, 2, \dots, n$), $\{O_i\}$ is a finite covering of E , hence it has the Lebesgue property. Therefore we can find a surrounding $V \in \mathfrak{S}$ such that $V(a) \subset O_i$ for some index i depending of a . Hence from $x, y \in V(a)$,

$$|f(x) - f(y)| \leq |f(x) - f(a)| + |f(a) - f(y)| < 2\varepsilon.$$

This shows that any bounded continuous function is uniformly continuous.

The proof of the "if" part of Theorem 1 is contained in my paper [4], therefore we shall omit it here.

We shall consider the Lebesgue property of a uniform space (not necessarily finite). An important contribution is due to S. KASAHARA [6]. He proved the following

THEOREM. *If a uniform space E has the Lebesgue property, then E is paracompact.*

Let $\{O_\alpha\}$ be a given covering of E , then we can find a surrounding V for $\{O_\alpha\}$.

Let W be a surrounding such that $W \circ W \circ W \subset V$, then it is easily seen that the covering $\{W(x) | x \in E\}$ is a star refinement of the covering $\{O_\alpha\}$. Thus E is fully normal in the sense of J. W. TUKEY. Thus E is paracompact. This is the idea of the proof by S. KASAHARA.

On the other hand, J. DE GROOT and H. DE VRIES [1] (or K. ISÉKI [2]) proved the following.

THEOREM. *A locally metrizable, paracompact T_3 -space is metrizable.*

This theorem follows from a result of the present author [2].

Therefore, we can easily prove the following

Theorem 2. *A uniform space E having the Lebesgue property is metrizable, if and only if E is locally metrizable.*

By a theorem of A. A. MONTEIRO and M. M. PEIXOTO [8], we can conclude

Theorem 3. *If a uniform space has the Lebesgue property, then every continuous function is uniformly continuous.*

Bibliography.

- [1] J. DE GROOT and H. DE VRIES, A note on non-Archimedean metrizations, *Nederl. Akad. Wetensch. Indag. Math.* **17** (1955), 222—224.
- [2] K. ISÉKI, A note on the general metrization problem, *Proc. Japan Acad.* **30** (1954), 855—856.
- [3] K. ISÉKI, On the property of Lebesgue in uniform spaces, *Proc. Japan Acad.* **31** (1955), 220—221.
- [4] K. ISÉKI, On the property of Lebesgue in uniform spaces, II., *Proc. Japan Acad.* **31** (1955), 270—271.
- [5] K. ISÉKI, On the property of Lebesgue in uniform spaces, III., *Proc. Japan Acad.* **31** (1955), 441—442.
- [6] S. KASAHARA, On the Lebesgue property in uniform spaces, to be published soon.
- [7] A. A. MONTEIRO and M. M. PEIXOTO, Note on uniform continuity, *Proc. of International Congress of Mathematicians, Cambridge Mass.* **1** (1950), 385.
- [8] A. A. MONTEIRO and M. M. PEIXOTO, Le nombre de Lebesgue et la continuité, *Portugal. Math.* **10** (1951), 105—113.
- [9] P. SAMUEL, Ultrafilters and compactification of uniform spaces, *Trans. Amer. Math. Soc.* **64** (1948), 100—132.

(Received September 20, 1955.)