

# The minimum number of spanning paths in a strong tournament

By J. W. MOON (Cape Town)

**1. Introduction.** A tournament  $T_n$  consists of a finite set of nodes  $1, 2, \dots, n$  such that each pair of distinct nodes  $i$  and  $j$  is joined by exactly one of the arcs  $\vec{ij}$  or  $\vec{ji}$ . If the arc  $\vec{ij}$  is in  $T_n$  we say that  $i$  beats  $j$  or  $j$  loses to  $i$  and write  $i \rightarrow j$ . If each node  $a$  of a subtournament  $A$  beats each node  $b$  of a subtournament  $B$  we write  $A \rightarrow B$  and let  $A+B$  denote the tournament determined by the nodes of  $A$  and  $B$ .

A path is a sequence  $P = \{p_1, p_2, \dots, p_k\}$  of distinct nodes of  $T_n$  such that  $p_i \rightarrow p_{i+1}$  if  $1 \leq i < k$ ; a spanning path contains every node of  $T_n$ . Let  $h(T_n)$  denote the number of spanning paths in the tournament  $T_n$ . RÉDEI [1] showed that  $h(T_n)$  is odd for every  $T_n$  and SZELE [2] showed that if  $H(n)$  denotes the maximum of  $h(T_n)$  over all tournaments with  $n$  nodes then

$$n!/2^{n-1} \leq H(n) \leq (n+1)!/2^{(3/4)n-3} \quad \text{for } n \geq 1.$$

A tournament  $T_n$  is strong if it cannot be expressed as  $T_n = A+B$  for some nonempty subtournaments  $A$  and  $B$ . If a tournament  $T_n$  is not strong, or weak, it has a unique expression of the type  $T_n = A+B+\dots+K$  where the nonempty subtournaments  $A, B, \dots, K$  all are strong and  $h(T_n) = h(A) \cdot h(B) \dots h(K)$ . Thus in considering lower bounds for  $h(T_n)$  we may as well assume  $T_n$  is strong. Our object here is to prove the following result.

**Theorem.** If  $h(n)$  denotes the minimum number of spanning paths a strong tournament  $T_n$  can have, then  $h(1)=1$  and

$$(1) \quad \alpha^{n-1} \leq h(n) \leq \begin{cases} \beta^{n-1} & n \equiv 1 \pmod{3} \\ 9 \cdot 5^{-4/3} \beta^{n-1} & \text{if } n \equiv 2 \pmod{3} \\ 3 \cdot 5^{-2/3} \beta^{n-1} & n \equiv 3 \pmod{3} \end{cases}$$

for  $n \geq 3$ , where  $\alpha = 6^{1/4} = 1.565$  and  $\beta = 5^{1/3} = 1.710$ .

**2. An upper bound for  $h(n)$ .** If  $n \geq 3$  let  $R_n$  denote the strong tournament in which  $i \rightarrow j$  if and only if  $i < j$  except that  $n \rightarrow 1$ . Any spanning path  $P$  of  $R_n$  that involves the arc  $\vec{n1}$  partitions the nodes  $(2, 3, \dots, n-1)$  into two subsets, those that come before  $n$  and those that come after  $1$ ; it follows from the definition of  $R_n$  that the nodes in these two subsets must occur in natural order in the path  $P$ . Conversely, each partition of the nodes  $(2, 3, \dots, n-1)$  into two subsets determines a unique spanning path that involves the arc  $\vec{n1}$ . There is only one spanning path of  $R_n$  that doesn't

involve the arc  $\vec{n1}$ , namely,  $\{1, 2, \dots, n\}$ . Consequently,  $h(R_n) = 1 + 2^{n-2}$  for  $n \geq 3$ .

More generally, let  $\{s_1, s_2, \dots, s_t\}$  denote an increasing subsequence of the nodes  $1, 2, \dots, n$  such that  $s_1 = 1$ ,  $s_t = n$ , and  $\alpha_i = s_{i+1} - s_i - 1 \geq 1$  for  $1 \leq i < t$ . Let  $S_n$  denote the strong tournament in which  $i < j$  if and only if  $i < j$  except that  $s_{i+1} \rightarrow s_i$  for  $1 \leq i < t$ . It is not difficult to show, by an extension of the argument just used for the case  $t=2$ , that

$$h(S_n) = \prod_{i=1}^{t-1} (1 + 2^{\alpha_i})$$

for  $n \geq 3$ . (Notice that  $h(S_n)$  is always odd, in accordance with Rédei's theorem.)

To minimize  $h(S_n)$  for given values of  $n$ , notice that

$$(1 + 2^{\alpha+3}) > (1 + 2^2)(1 + 2^\alpha) \quad \text{for } \alpha \geq 1,$$

$$(1 + 2^3)^2 > (1 + 2^1)(1 + 2^2)^2,$$

and

$$(1 + 2^1)(1 + 2^3) = (1 + 2^1)^3 > (1 + 2^2)^2.$$

Thus if  $n = 3k+1$  the minimum occurs when  $t = k+1$  and all the  $\alpha$ 's equal two; if  $n = 3k+2$  it occurs when  $t = k+1$  and one  $\alpha$  equals three and the rest equal two or when  $t = k+2$  and two  $\alpha$ 's equal one and the rest equal two; if  $n = 3k+3$  it occurs when  $t = k+2$  and one  $\alpha$  equals one and the rest equal two. It follows, therefore, that the minimum value of  $h(S_n)$  is  $5^k$ ,  $9 \cdot 5^{k-1}$ , or  $3 \cdot 5^k$  according as  $n = 3k+1$ ,  $3k+2$ , or  $3k+3$ . This implies the upper bound in inequality (1).

**3. A lower bound for  $h(n)$ .** Before proceeding we remark that if node  $y$  is not contained in a path  $P$  and  $y$  beats some node  $x$  of  $P$  then  $y$  can be inserted in the path  $P$  before  $x$ ; in particular,  $y$  can be inserted immediately before the first node of  $P$  it beats. Similarly, if  $y$  loses to  $x$  then  $y$  can be inserted in  $P$  after  $x$ .

It is not difficult to verify that  $h(1)=1$ ,  $h(3)=3$ ,  $h(4)=5$ , and  $h(5)=9$  so the lower bound in inequality (1) certainly holds for small values of  $n$ . Suppose the inequality has been established for  $3 \leq n \leq m-1$  and consider a strong tournament  $T_m$  such that  $h(m) = h(T_m)$ .

Let  $K$  denote any minimal subset of nodes of  $T_m$  whose removal, along with all incident arcs, leaves a weak subtournament  $R$ ; such a subset  $K$  exists since, for example the removal of the nodes that lose to any given node of  $T_m$  leaves a weak subtournament. Since  $R$  is weak there exist subtournaments  $A$ ,  $B$  and  $C$  where  $A$  and  $C$  are strong such that  $R = A + B + C$ ; it may be that  $B$  is vacuous. Let  $a$ ,  $b$ ,  $c$  and  $k$  denote the number of nodes in  $A$ ,  $B$ ,  $C$  and  $K$  so that  $a + b + c + k = m$ . It follows from the minimality property of  $K$  and the fact that  $T_m$  is strong that each node of  $K$  loses to at least one node of  $C$  and beats at least one node of  $A$ . We now construct two families of spanning paths of  $T_m$  and thus obtain a lower bound for  $h(m)$ .

Let  $P_1$ ,  $P_2$ , and  $P_3$  denote any spanning paths of  $A$ ,  $B$ , and  $C$ ; there are at least  $h(a)h(c)$  spanning paths of  $R$  and they all are of the type  $P_1 + P_2 + P_3$ . Now let  $X$  and  $Y$  denote any partition of the nodes of  $K$  into two subsets. Since each node of  $X$  beats some node of  $A$  and each node of  $Y$  loses to some node of  $C$  it follows that the nodes of  $X$  and  $Y$  can be inserted in the path  $P_1 + P_2 + P_3$  so as to form a spanning path  $Q$  of  $T_m$  that first passes through the nodes of  $X$  and  $A$ ,

then passes through the nodes of  $B$ , and finally passes through the nodes of  $Y$  and  $C$ . Different paths  $P_1$  and  $P_3$  and different partitions  $X$  and  $Y$  yield different paths  $Q$  so  $T_m$  must have at least  $2^k h(a)h(c)$  spanning paths of this first type.

Let  $x$  denote any given node of  $K$  and let  $u$  and  $v$  denote given nodes of  $A$  and  $C$  that lose to  $x$  and beat  $x$ , respectively. For any spanning path  $P_1$  of  $A$  let  $P_{11}$  denote the (possibly empty) subpath determined by the nodes that come before  $u$  in  $P_1$  and let  $P_{12}$  denote the subpath determined by  $u$  and the nodes that come after  $u$  in  $P_1$ ; for any spanning path  $P_3$  of  $C$  let  $P_{31}$  denote the subpath determined by  $v$  and the nodes that come before  $v$  in  $P_3$  and let  $P_{32}$  denote the (possibly empty) subpath determined by the nodes that come after  $v$  in  $P_3$ ; for any partition of the nodes of  $B$  into two subsets  $W$  and  $Z$ , let  $P_{21}$  and  $P_{22}$  denote spanning paths of the subtournaments determined by  $W$  and  $Z$ . It is not difficult to see that for each partition  $W$  and  $Z$  there are at least  $h(a)h(c)$  paths of the type  $P^* = P_{11} + P_{21} + P_{31} + \{x\} + P_{12} + P_{22} + P_{32}$  that pass through  $x$  and the nodes of  $R$ . We now estimate the number of ways of inserting the remaining nodes of  $K$  into such a path  $P^*$ .

If  $k \geq 2$ , let  $y$  denote any node of  $K - \{x\}$ . Since  $y$  beats some node of  $A$  it follows that  $y$  can be inserted in  $P^*$  before the last node of  $P_{12}$ ; it can also be inserted after the last node of  $P_{12}$  unless it beats that node and all the nodes of  $P_{22}$  and  $P_{32}$ . Since  $y$  loses to some node of  $C$  it follows that  $y$  can be inserted in  $P^*$  after the first node of  $P_{31}$ ; it can also be inserted before the first node of  $P_{31}$  unless it loses to that node and to all nodes of  $P_{11}$  and  $P_{21}$ . Thus if  $y$  can't be inserted in  $P^*$  in at least two ways it must be that  $y \rightarrow Z$  and  $W \rightarrow y$ , among other things. Notice that this last statement also applies when  $P^*$  is replaced by any path obtained by inserting other nodes of  $K - \{x\}$  in  $P^*$ .

Let the partitions of  $B$  into two subsets be numbered from  $i=1$  to  $i=t=2^b$ . For the  $i$ -th partition,  $W_i$  and  $Z_i$ , let  $\gamma_i$  denote the number of nodes  $y$  of  $K - \{x\}$  such that  $y \rightarrow Z_i$  and  $W_i \rightarrow y$ . It follows from the observations in the last paragraph that there are at least  $2^{k-1-\gamma_i}$  ways to insert the nodes of  $K - \{x\}$  in any path  $P^*$  associated with the partition  $W_i$  and  $Z_i$  so as to form a spanning path of  $T_m$ . If we apply this construction for all spanning paths of  $A$  and  $C$  and for all partitions of  $B$  we obtain at least

$$(2) \quad 2^{k-1} (2^{-\gamma_1} + \dots + 2^{-\gamma_t}) h(a)h(c)$$

different spanning paths of  $T_m$ . (Some node of  $C$  comes before some node of  $A$  in each of these paths so they are not the same as any of the paths enumerated earlier.) Since the nodes of  $B$  that beat and lose to any node  $y$  of  $K - \{x\}$  are fixed, each such node  $y$  contributes one to just one  $\gamma_i$  and  $\gamma_1 + \dots + \gamma_t = k-1$ . Therefore, the sum in (2) is minimized when all the  $\gamma_i$ 's equal  $(k-1)2^{-b}$  and  $T_m$  must have at least  $h(a)h(c)2^{b+(k-1)(1-2^{-b})}$  spanning paths of this second type.

It follows from these two constructions that  $h(m) \geq (2^k + 2^{b+(k-1)(1-2^{-b})})h(a)h(c)$ . In order to show that this implies that  $h(m) \geq \alpha^{m-1} = \alpha^{a+b+c+k-1}$ , it suffices to show, in view of the induction hypothesis, that

$$(3) \quad 2^k + 2^{b+(k-1)(1-2^{-b})} \geq \alpha^{b+k+1} = 6^{(b+k+1)/4}$$

for all integers  $b$  and  $k$  such that  $b \geq 0$  and  $k \geq 1$ .

Since  $\log_2 \alpha < \frac{2}{3}$ , inequality (3) certainly holds if  $k \geq 2(b+k+1)/3$  or  $k \geq 2b+2$  and it can easily be verified directly for  $1 \leq k \leq 2b+1$  when  $0 \leq b \leq 2$ . (Notice that equality holds in (3) when  $k=1$  and  $b=2$ .) If  $b \geq 3$ , then inequality (3) holds if

$$(4) \quad 2^k + 2^{b+7(k-1)/8} \geq \alpha^{b+k+1}$$

for  $b \geq 3$  and  $k \geq 1$ . This inequality certainly holds if  $b+7(k-1)/8 \geq 2(b+k+1)/3$  or  $5k+8b \geq 37$ ; if  $b \geq 4$  this holds for  $k \geq 1$  and if  $b=3$  it holds if  $5k \geq 13$  or if  $k \geq 3$ . In the remaining cases, when  $b=3$  and  $k=1$  or  $2$ , inequality (4) can be verified directly. This suffices to complete the proof of the lower bound of inequality (1) by induction.

### References

- [1] L. RÉDEI, Ein kombinatorischer Satz, *Acta Litt. Szeged*, 7 (1934), 39—43.
- [2] T. SZELE, Kombinatorikai vizsgálatok az irányított teljes gráffal kapcsolatban, *Mat. Fiz. Lapok*, 50, (1943), 223—256. Reprinted as Kombinatorische Untersuchungen über gerichtete vollständige Graphen, *Publ. Math. Debrecen*, 13, (1966), 145—168.

The preparation of this paper was assisted by a grant from the National Research Council of Canada.

(Received September 1, 1970.)