

Boundedness on a set of positive measure and the fundamental equation of information

By G. T. DIDERRICH*) (Milwaukee, Wis.)

1. Introduction. An information function f is a real valued function defined on $[0, 1]$ which satisfies the following functional equation and boundary conditions

$$f(x) + (1-x)f\left(\frac{y}{1-x}\right) = f(y) + (1-y)f\left(\frac{x}{1-y}\right)$$

whenever $x, y \in [0, 1[$ and $x+y \leq 1$

with $f(0) = f(1)$ and $f\left(\frac{1}{2}\right) = 1$.

These conditions imply that $f(x) = f(1-x)$, $x \in [0, 1]$ and $f(0) = 0$. The above functional equation is called the fundamental equation of information because the problem of characterizing Shannon's measure of entropy can be reduced to a study of it. (For details and terminology see ACZÉL and DARÓCZY [1]). The Shannon measure of entropy in a 2-event space

$$S(x) = -x \log x - (1-x) \log (1-x), \quad x \in [0, 1]$$

(with the conventions $\log = \log_2$ and $0 \log 0 = 0$) is the most important information function.

In [2] we have proved the following result: If f is an information function and f is bounded on an arbitrarily small nonvanishing interval contained in $]0, 1[$, then $f = S$. The following question seems quite natural: Can one replace "bounded on an arbitrary small nonvanishing interval" by "bounded on a set of positive measure" in the above statement (measurable is meant in the Lebesgue sense with μ denoting the measure). In this connection there is a result in [2].

Corollary 1.1. *Suppose that, for every $\varepsilon > 0$, there is a positive constant K and a measurable set E_K on which the information function f is bounded by K and $\mu(E_K) \geq 1 - \varepsilon$, then $f = S$.*

In this paper we give a positive answer to the above question.
We prove the following:

*) Research Conducted Under a National Research Council Grant at the University of Waterloo, Waterloo, Ontario, Canada in the Pure Mathematics Department.

Theorem. Let f be an information function. If f is bounded on a set of positive measure, then $f=S$.

2. Proof of the Theorem. The proof is carried out by imitating the procedure in [1] (Section 3.4) and introducing a weaker definition of the multiplicative group G_f . The idea is to spread out the boundedness to larger and larger portions of $]0, 1[$ where intervals are replaced by measurable subsets together with the concept of a "right-sided density point". Some definitions follow.

Let E be a Borel subset on the real line. A point x is said to be a two-sided density of E , if

$$\lim_{\delta \rightarrow 0+} \frac{\mu(E \cap]-\delta+x, x+\delta])}{2\delta} = 1.$$

Almost all points of E are density points (see SAKS [7] Chapter 4 and RUDIN [6] Chapter 8). In an analogous way we say x is a right-sided density point, if

$$\lim_{\delta \rightarrow 0+} \frac{\mu(E \cap [x, x+\delta])}{\delta} = 1$$

with a similar definition for the concept of left-sided density point. We now proceed to the lemmas.

Lemma 2.1. (See JOU and BAKER [3].) Let $E_1, E_2, \dots, E_n$ and I be Borel subsets of the real line with $0 < \mu(I) < \infty$. Then

$$1 - \frac{\mu(E_1 \cap \dots \cap E_n \cap I)}{\mu(I)} \leq \sum_{i=1}^n \left[1 - \frac{\mu(E_i \cap I)}{\mu(I)} \right].$$

From this lemma it is clear that a two-sided density point is also a right (left)-sided density point.

Lemma 2.2. Let E_1, E_2 be Borel subsets of the real line. Suppose that x is a two-sided (right, left) density point of E_1 and E_2 . Then x is a two-sided (right, left) density point of $E_1 \cap E_2$.

PROOF. We consider the two-sided case; the other cases follow in a similar fashion. From the definition

$$\lim_{\delta \rightarrow 0+} \frac{\mu(E_i \cap I_\delta)}{2\delta} = 1 \quad (i = 1, 2)$$

where $I_\delta =]-\delta+x, x+\delta[$. The preceding lemma gives

$$0 \leq 1 - \frac{\mu(E_1 \cap E_2 \cap I_\delta)}{2\delta} \leq 1 - \frac{\mu(E_1 \cap I_\delta)}{2\delta} + 1 - \frac{\mu(E_2 \cap I_\delta)}{2\delta}$$

and hence the present lemma.

Lemma 2.3. Let T be a continuously differentiable transformation from an open interval $V =]a, b[$ to an open interval $W =]c, d[$. Assume that $|T'| \neq 0$ on V . Let E be a set of positive measure in V and suppose that $x \in E$ is a two-sided density point of E . Then Tx is a two-sided density point of TE . In particular, if $T' > 0$ ($T' < 0$)

and $x \in E$ is a right-sided density point of E , then Tx is a right (left)-sided density point of TE .

PROOF. We consider the $T' > 0$ case in the last sentence of the lemma. The other cases follow in a similar way. Let x be a right-sided density point of E , so

$$\lim_{\delta \rightarrow 0+} \frac{\mu(E \cap I_\delta)}{\delta} = 1$$

where $I_\delta = [x, x + \delta[$. Clearly T is a 1-1 open mapping of V into W , hence $T(E \cap I_\delta) = TE \cap TI_\delta$. Furthermore TI_δ is an interval in W of the form $[Tx, Tx + \delta_1[$ where $\delta_1 = \mu(TI_\delta)$. We have by [5] (Theorem 8.26) that

$$(1) \quad \mu(TE \cap TI_\delta) = \int_{E \cap I_\delta} |T'(y)| d\mu(y).$$

It suffices to prove that

$$\lim_{\delta_1 \rightarrow 0} \frac{\mu(TE \cap TI_\delta)}{\mu(TI_\delta)} = 1$$

to reach the desired conclusion. From (1) we obtain

$$\inf_{y \in E \cap I_\delta} |T'(y)| \mu(E \cap I_\delta) \leq \mu(TE \cap TI_\delta) \leq \sup_{y \in E \cap I_\delta} |T'(y)| \mu(E \cap I_\delta).$$

Therefore

$$(2) \quad \frac{\frac{l(E \cap I_\delta)}{\mu(I_\delta)} \frac{\mu(E \cap I_\delta)}{\mu(I_\delta)}}{\frac{\mu(TI_\delta)}{\mu(I_\delta)}} \leq \frac{\mu(TE \cap TI_\delta)}{\mu(TI_\delta)} \leq \frac{\frac{L(E \cap I_\delta)}{\mu(I_\delta)} \frac{\mu(E \cap I_\delta)}{\mu(I_\delta)}}{\frac{\mu(TI_\delta)}{\mu(I_\delta)}}$$

where $l(E \cap I_\delta) = \lim_{y \in E \cap I_\delta} |T'(y)|$ and $L(E \cap I_\delta) = \sup_{y \in E \cap I_\delta} |T'(y)|$. But it is clear, using the continuity of T' and the definitions, that the left and right extremities of (2) go to 1 as $\delta_1 \rightarrow 0$. This completes the proof.

Definition. Let f be a real-valued function defined on $[0, 1]$. The positive real number λ belongs to G_f , if there exist positive numbers k_λ , δ_λ and a Borel set I_λ contained in $[0, 1]$ with $0 \in I_\lambda$ as a right-sided density point such that

$$\left| f(y) - \lambda f\left(\frac{y}{\lambda}\right) \right| < k_\lambda$$

for every $y \in I_\lambda \cap [0, \delta[$ ($0 < \delta \leq \delta_\lambda$).

Roughly speaking, λ belongs to G_f if $\left| f(y) - \lambda f\left(\frac{y}{\lambda}\right) \right|$ becomes eventually bounded near $[0, \delta[$ on a set whose measure becomes closer and closer to δ as $\delta \rightarrow 0$.

Lemma 2.4. G_f is a multiplicative subgroup of $]0, \infty[$, the multiplicative group of positive reals.

PROOF.

(a) Clearly $1 \in G_f$.

(b) Suppose that $\lambda \in G_f$, we want to show that $\frac{1}{\lambda} \in G_f$. The definition yields

positive numbers k_1 , δ_1 and a Borel set I_1 with $0 \in I_1$ as a right-sided density point such that

$$(3) \quad \left| f(y) - \lambda f\left(\frac{y}{\lambda}\right) \right| < k_1$$

for each $y \in I_1 \cap [0, \delta[$ ($0 < \delta \leq \delta_1$). Now put $I_0 = \frac{I_1}{\lambda}$. Lemma 2.3 implies that $0 \in I_0$ is a right-sided density point. Choose $0 < \delta_0 \leq \frac{\delta_1}{\lambda}$. Hence, $x \in I_0 \cap [0, \delta_0[$ implies that $x = \frac{y}{\lambda}$ where $y \in I_1 \cap [0, \delta_1[$. Thus, dividing (3) by λ , we obtain

$$(4) \quad \left| f(x) - \frac{1}{\lambda} f(x\lambda) \right| = \left| f\left(\frac{y}{\lambda}\right) - \frac{1}{\lambda} f(y) \right| < \frac{k_1}{\lambda}$$

for each $x \in I_0 \cap [0, \delta[$ ($0 < \delta \leq \delta_0$). Hence we proved $\frac{1}{\lambda} \in G_f$.

(c) We now prove $\lambda_1, \lambda_2 \in G_f$ imply $\lambda_1 \lambda_2 \in G_f$. Our hypothesis gives

$$(5) \quad \left| f(y) - \lambda_i f\left(\frac{y}{\lambda_i}\right) \right| < k_i$$

for $y \in I_i \cap [0, \delta[$ where $0 < \delta \leq \delta_i$ ($i=1, 2$). Put $I_0 = I_1 \cap I_2 \lambda_1$ and $\delta_0 = \min \{\delta_1, \delta_2 \lambda_1\}$. It follows from Lemma 2.2 and Lemma 2.3 that 0 is a right-sided density point of I_0 . We claim that

$$(6) \quad \left| \lambda_1 f\left(\frac{x}{\lambda_1}\right) - \lambda_1 \lambda_2 f\left(\frac{x}{\lambda_1 \lambda_2}\right) \right| < \lambda_1 k_2$$

for every $x \in I_0 \cap [0, \delta[$ where $0 < \delta \leq \delta_0$. This follows by multiplying (5), in the $i=2$ case, by λ_1 and noting that $x \in I_0 \subset I_2 \lambda_1$ implies $\frac{x}{\lambda_1} \in I_2$. Also (5), in the $i=1$ case, yields

$$(7) \quad \left| f(x) - \lambda_1 f\left(\frac{x}{\lambda_1}\right) \right| < k_1$$

for every $x \in I_0 \cap [0, \delta[$ ($0 < \delta \leq \delta_0$). Adding (6) and (7) results in

$$\left| f(x) - \lambda_1 \lambda_2 f\left(\frac{x}{\lambda_1 \lambda_2}\right) \right| < k_1 + \lambda_1 k_2$$

for every $x \in I_0 \cap [0, \delta[$ ($0 < \delta \leq \delta_0$). This completes the proof of the lemma.

Lemma 2.5. *If an information function f is bounded on a set of positive measure E , then $G_f =]0, \infty[$.*

PROOF. Almost all points in E are two-sided density points, a fortiori, right-sided density points. Fix a density point x in $E \cap]0, 1[$. Assume that $y \rightarrow 0+$ such

that $x+y \leq 1$. Then the fundamental equation gives

$$(8) \quad f(x) + (1-x)f\left(\frac{y}{1-x}\right) = f(y) + (1-y)f\left(\frac{x}{1-y}\right).$$

Now $\frac{x}{1-y} \in E$ if and only if $y \in I = (1 - xE^{-1})$. An application of Lemma 2.3, with $T(z) = 1 - xz^{-1}$, shows that $0 \in I$ is a right-sided density point of I . Equation (8) gives

$$\left| f(x) - (1-y)f\left(\frac{x}{1-y}\right) \right| = \left| f(y) - (1-x)f\left(\frac{y}{1-x}\right) \right| < 2k$$

for $y \in I \cap [0, \delta[$ and small enough $\delta > 0$ where $k > 0$ is the bound for f on E . The preceding argument clearly proves that $1 - x \in G_f$. Hence, $G_f \supset 1 - E^*$ is the set of density points of $E \cap]0, 1[$.

The group $(G_f, \cdot)$ is isomorphic to a subgroup $(\mathbf{R}_f, +)$ of the additive group of reals $(\mathbf{R}, +)$ via the mapping $x \rightarrow \log x$. But $(\mathbf{R}_f, +)$ contains a set of positive measure, $F = \log(1 - E^*)$. The linear Steinhaus theorem (see KEMPERMAN [4], Section 2) implies that $F + F$ contains a non-degenerate interval. So $(\mathbf{R}_f, +) = (\mathbf{R}, +)$, hence $G_f =]0, \infty[$ which completes the proof of the lemma.

Lemma 2.6. *If $G_f =], \infty[$, then every point x in $]0, 1[$ is in a set E_x of positive measure with x as a right-sided density point on which f is bounded. Indeed, $E_x = x(1 - I_{1-x})^{-1}$ where I_{1-x} is the Borel set associated with $1 - x$ in the definition of G_f .*

PROOF. Now $1 - x \in G_f$ ($0 < x < 1$). Therefore the definition of G_f yields positive numbers k_1, δ_1 and a Borel set I with $0 \in I$ as a right-sided density point such that

$$(9) \quad \left| f(y) - (1-x)f\left(\frac{y}{1-x}\right) \right| < k_1$$

for $y \in I \cap [0, \delta_1[$ ($0 < \delta_1 \leq \delta_1$). The fundamental equation (8) together with (9) then gives

$$(10) \quad \left| f(x) - (1-y)f\left(\frac{x}{1-y}\right) \right| < k_1$$

for $y \in I \cap [0, \delta_1[$. But Lemma 2.3, using $T(z) = x(1 - z)^{-1}$, shows that $E_x = x(1 - I)^{-1}$ with (10) has the desired properties.

Lemma 2.7. *If f is bounded on a Borel set of positive measure in $]0, 1[$, then f is bounded on Borel sets of measure arbitrarily close to 1.*

PROOF. We use a modified Heine—Borel type of argument. Let $F = [a, b]$ ($a < b$) be a given closed interval in $]0, 1[$. Let $N \geq 2$ be a given positive integer. Let T denote the set of real numbers t such that $a \leq t \leq b$ and there is a Borel set E contained in $[a, t]$ with $\mu(E) \geq (t - a)\left(1 - \frac{1}{N}\right)$ on which f is bounded. Let $c = \sup T$. By our hypothesis and Lemma 2.6, it is clear that $c > a$.

We now argue that $c \in T$. Lemma 2.6 shows that c is a left-sided density point of $E'_c = 1 - E_{1-c}$ on which f is bounded. Hence for t sufficiently close to c ($t < c$) we can find a measurable set E_1 contained in $[t, c]$ on which f is bounded and

$$(11) \quad \mu(E_1) \cong (c-t) \left(1 - \frac{1}{N}\right).$$

But by the definition of sup and by our hypothesis, we can find a measurable set E_2 contained in $[a, t]$ on which f is bounded and

$$(12) \quad \mu(E_2) \cong (t-a) \left(1 - \frac{1}{N}\right).$$

Thus (11) and (12) yield the set $E = E_1 \cup E_2 \subset [a, c]$ on which f is bounded and

$$\mu(E) \cong (c-a) \left(1 - \frac{1}{N}\right)$$

which demonstrates that $c \in T$.

Let $E_1 \subset [a, c]$ be the set associated with c , so

$$(13) \quad \mu(E_1) \cong (c-a) \left(1 - \frac{1}{N}\right)$$

and f is bounded on E_1 . Again by Lemma 2.6, we can find a Borel set E_2 , disjoint from E_1 , with c as a right-sided density point such that $E_2 \subset [c, c_1]$ for some $c_1 > c$ with

$$(14) \quad \mu(E_2) \cong (c_1 - c) \left(1 - \frac{1}{N}\right)$$

on which f is bounded. Clearly f is bounded on $E = E_1 \cup E_2 \subset [a, c_1]$ and (13) and (14) yield

$$\mu(E) = \mu(E_1) + \mu(E_2) \cong (c-a) \left(1 - \frac{1}{N}\right) + (c_1 - c) \left(1 - \frac{1}{N}\right) \cong (c_1 - a) \left(1 - \frac{1}{N}\right).$$

But this contradicts the definition of c and, as a consequence, c cannot be an interior point, so $c = b$. Hence we proved that every closed interval $[a, b]$ contains a measurable set E such that $\mu(E) \cong (b-a) \left(1 - \frac{1}{N}\right)$ on which f is bounded. To complete the proof of the lemma, take $[a, b] = \left[\frac{1}{k}, 1 - \frac{1}{k}\right]$ with k sufficiently large and N sufficiently large.

The theorem now follows from Corollary 1.1 of [2]. The reader is referred to [5] for simplified proofs of the main result here and the result in [2].

References

- [1] J. ACZÉL and Z. DARÓCZY, On measures of information and their characterizations, *Academic Press, New York*, 1975.
- [2] G. T. DIDERRICH, Local boundedness and the Shannon entropy, *Information and Control*, **36** (1978), 292—308.
- [3] S. S. JOU and J. BAKER, On transformations of measurable sets in locally compact topological groups, *Glasnik Mat.* **9** (29) (1974), 35—41.
- [4] J. H. B. KEMPERMAN, A general functional equation, *Trans. Amer. Math. Soc.* **86** (1957), 28—56.
- [5] GY. MAKSA, Bounded symmetric information functions, *C. R. Math. Rep. Acad. Sci. Canada* — **II** (1980) *No. 5*, 247—252.
- [6] W. RUDIN, Real and complex analysis, *McGraw-Hill Book Company, New York* 1974.
- [7] S. SAKS, Theory of the integral, *Hafner Publishing Company, New York* 1937.

JOHNSON CONTROLS, INC.
P.O. BOX 423
MILWAUKEE, WIS. 53201
U.S.A.

(Received July 6, 1982)