

Completeness of Finsler manifolds

By CONSTANTIN UDRIȘTE* (Bucharest)

Dedicated to Professor Lajos Tamásy on his 70th birthday

Abstract. This paper analyses some constructions that produce complete Finsler manifolds:

- 1) Let the Finsler manifolds $(M, g(x, y))$ and $(M, \bar{g}(x, y))$ be given. Then $(M, \bar{g}(x, y))$ is complete if $(M, g(x, y))$ is complete and the tensor field $\bar{g} - g$ is positive semi-definite.
- 2) If $(M, g(x, y))$ is a Finsler manifold and $f : M \rightarrow \mathbb{R}$ is a proper function then the Finsler manifold $(M, g(x, y) + df(x) \otimes df(x))$ is complete. Using this construction we prove that a Finsler manifold which supports a proper function whose differential has bounded relative length is complete.
- 3) Let the Finsler manifolds $(M_1, g_1(x_1, y_1))$ and $(M_2, g_2(x_2, y_2))$ be given and suppose that $f > 0$ is a differentiable function on M_1 . The warped product $(M_1 \times M_2, g_1 + fg_2)$ is complete if and only if $(M_1, g_1(x_1, y_1))$ and $(M_2, g_2(x_2, y_2))$ are complete.

§1. Complete Finsler manifolds [2], [4], [5]

Let M be an n -dimensional connected C^3 -manifold and TM its tangent bundle. Denote by (x, y) an arbitrary point in TM and by x the corresponding point in M .

Definition 1.1. A Finsler tensor field $g(x, y)$ of type $(0, 2)$ which is symmetric, positive definite and whose components $g_{ij}(x, y)$ are homogeneous functions of degree zero with respect to y is called a *Finsler metric* on M . The pair $(M, g(x, y))$ is called a Finsler manifold.

The function

$$L : TM \rightarrow \mathbb{R}, \quad L(x, y) = \sqrt{g_{ij}(x, y)y^i y^j}$$

1980 *Mathematics Subject Classification* (1985 *Revision*): 53C60, 54E50.

Keywords: Finsler manifolds, criteria for completeness, warped products.

*This paper was presented at the Conference on Finsler Geometry and its Application to Physics and Control Theory, August 26–31, 1991, Debrecen, Hungary.

is called fundamental Finsler function and L^2 is called *absolute Finsler energy*.

For a vector field $V = V^i(x) \frac{\partial}{\partial x^i}$ on M we have two kinds of lengths: the absolute length

$$L(x, V(x)) = \sqrt{g_{ij}(x, V(x)) V^i(x) V^j(x)}$$

and the relative length

$$\|V(x)\|_y = \sqrt{g_{ij}(x, y) V^i(x) V^j(x)}.$$

Remark. If $g_{ij}(x, V(x)) - g_{ij}(x, y)$ is negative semidefinite, then the absolute length of $V(x)$ is the minimum of the relative length of $V(x)$.

In a Finsler manifold $(M, g(x, y))$ the length ℓ of a curve arc $\gamma : [0, 1] \rightarrow M$ is given by

$$\ell(\gamma) = \int_0^1 L(\gamma(t), \dot{\gamma}(t)) dt.$$

Definition 1.2. let $(M, g(x, y))$ be a Finsler manifold. The function

$$\exp_x : 0_x \subset T_x M \rightarrow M, \quad X \rightarrow \exp_x X,$$

where $\exp_x X$ is the terminal point $\gamma(1)$ of the geodesic $\gamma : [0, 1] \rightarrow M$, $\gamma(0) = x$, $\dot{\gamma}(0) = X$ is called the *exponential map*.

The curve $\gamma : [0, 1] \rightarrow M$, $\gamma(t) = \exp_x(tX)$, $X \in T_x M$ is a geodesic which joins the points x and $\exp_x X$. The length of this geodesic is $L(x, X)$.

Definition 1.3. The *distance* $d(x, x')$ between the points $x, x' \in M$ is the infimum of the lengths of all curves from x to x' .

This definition is correctly in sense that the properties:

- 1) $d(x, x') \geq 0$, $\forall x, x' \in M$
- 2) $d(x, x') = 0$, if and only if $x = x'$
- 3) $d(x, x') = d(x', x)$, $\forall x, x' \in M$
- 4) $d(x, x') \leq d(x, x'') + d(x'', x')$, $\forall x, x', x'' \in M$

are satisfied. Also, the topology of M induced by the distance d coincides with the manifold topology of M .

Definition 1.4. The Finsler manifold $(M, g(x, y))$ is called *geodesically complete* if the exponential map $\exp_x$ is defined on the whole of $T_x M$ for any point of M .

Definition 1.5. The Finsler manifold $(M, g(x, y))$ is called *metrically complete* if the metric space (M, d) is complete.

Theorem 1.1. [2] *For a Finsler manifold $(M, g(x, y))$ the following three conditions are equivalent:*

- 1) $(M, g(x, y))$ is geodesically complete.
- 2) $(M, g(x, y))$ is metrically complete.
- 3) Any bounded closed subset of M is compact.

Remarks. 1) Let (M, g) and $(M, \bar{g})$ be two Finsler manifolds. Then $(M, \bar{g})$ is complete if (M, g) is complete and the tensor field $\bar{g} - g$ is positive semidefinite.

2) Let (M_1, d_1) and (M_2, d_2) be complete metric spaces. The product space $(M_1 \times M_2, d_1 + d_2)$ is complete.

3) Let $g_{ij}(x, y) = \gamma_{ij}(x) + c^{-2}y_i y_j$, where γ_{ij} is a Riemann metric tensor, c is the universal speed-of light constant, $\dot{x}^i$ is the tangent vector supported by a point $x = (x^i)$ and $y_i = \gamma_{ij}(x)\dot{x}^j$. If the Riemann manifold $(M, \gamma_{ij}(x))$ is complete, then the generalized Lagrange manifold $(M, g_{ij}(x, y))$ is complete. This generalized Lagrange manifold is not reducible to a Lagrange manifold, neither to a Finsler manifold nor to a Riemannian manifold [3].

§2. Analytical criterion for completeness

Definition 2.1. A continuous function $f : M \rightarrow \mathbb{R}$ is called *proper* if $f^{-1}(K)$ is a compact set whenever K is compact.

Theorem 2.1. *Let $(M, g(x, y))$ be a Finsler C^3 -manifold (not necessarily complete) and $f : M \rightarrow \mathbb{R}$ a proper C^3 function.*

The Finsler manifold $(M, \tilde{g}(x, y) = g(x, y) + df(x) \otimes df(x))$ is complete.

PROOF. We consider the Finsler manifold $(M \times \mathbb{R}, h)$, where $h_{ij} = g_{ij}$, $h_{i n+1} = 0$, $i, j = 1, 2, \dots, n$, $h_{n+1 n+1} = 1$. The graph

$$G(f) = \{(x, f(x)) | x \in M\}$$

is a submanifold of the product manifold $M \times \mathbb{R}$, diffeomorphic to M .

The Finsler metric $h_{\alpha\beta}$ induces on $G(f)$ the Finsler metric $\tilde{g} = g + df \otimes df$.

If $\{(x_n, f(x_n))\}$ is a Cauchy sequence of elements in $G(f)$, then $\{f(x_n)\}$ is a Cauchy sequence in $\mathbb{R}$ because

$$\begin{aligned} d_{G(f)}[(x, f(x)), (x', f(x'))] &\geq d_{M \times \mathbb{R}}[(x, f(x)), (x', f(x'))] \geq \\ &\geq |f(x) - f(x')|. \end{aligned}$$

Then there exists $z \in \mathbb{R}$ with $z = \lim_{n \rightarrow \infty} f(x_n)$ and hence $\{z, f(x_1), \dots, f(x_n), \dots\}$ is a compact set in $\mathbb{R}$. But $\{x_1, \dots, x_n, \dots\} \subset f^{-1}(\{z, f(x_1), \dots\})$

$\dots, f(x_n), \dots\}$) and f is proper. Hence the sequence $\{x_n\}$ contains a convergent subsequence since $\{x_n\}$ is contained in a compact set.

So the sequence $\{(x_n, f(x_n))\}$ is covergent and hence $(M, \tilde{g}(x, y))$ is complete.

Theorem 2.2. *Let $(M, g(x, y))$ be a Finsler C^3 -manifold. If there exists a proper C^3 function $f : M \rightarrow \mathbb{R}$ such that the Finsler tensor field $g(x, y) - df(x) \otimes df(x)$ is positive definite, then $(M, g(x, y))$ is complete.*

PROOF. Put, $\tilde{g} = g - df \otimes df$. If $\tilde{g}$ is positive definite, then $(M, \tilde{g}(x, y))$ is a Finsler C^3 -manifold. As $f : M \rightarrow \mathbb{R}$ is a proper function, we apply theorem 2.1 which says that $(M, \tilde{g} + df \otimes df)$ is complete. But $\tilde{g} + df \otimes df = g$. Hence $(M, g(x, y))$ is complete.

Theorem 2.3. [1]. *Any C^3 -manifold M supports a proper C^3 function $f : M \rightarrow \mathbb{R}$.*

Theorem 2.4. *Let $(M, g(x, y))$ be a Finsler C^3 -manifold and $f : M \rightarrow \mathbb{R}$ a C^3 function. Then the Finsler tensor field*

$$\tilde{g}(x, y) = g(x, y) - df(x) \otimes df(x)$$

is positive definite iff $\|df(x)\|_y < 1$, for any vector y .

PROOF. Let $V^i(x, y) = g^{ij}(x, y) \frac{\partial f(x)}{\partial x^j}$ and $V(x, y) = V^i(x, y) \frac{\delta}{\delta x^i}$. Here $\frac{\delta}{\delta x^i} = \frac{\partial}{\partial x^i} - N_i^j \frac{\partial}{\partial y^j}$ is a local base adapted to the horizontal canonical distribution $N = (N_i^j(x, y))$ of the manifold (M, g) .

Suppose that $\tilde{g}(x, y)$ is positive definite and x is not a critical point of f . Hence

$$\begin{aligned} 0 < \tilde{g}_{ij}(x, y) V^i(x, y) V^j(x, y) &= g_{ij}(x, y) g^{ik}(x, y) \frac{\partial f(x)}{\partial x^k} g^{j\ell}(x, y) \frac{\partial f(x)}{\partial x^\ell} - \\ &\quad - \frac{\partial f(x)}{\partial x^i} \frac{\partial f(x)}{\partial x^j} g^{ik}(x, y) \frac{\partial f(x)}{\partial x^k} g^{j\ell}(x, y) \frac{\partial f(x)}{\partial x^\ell} = \\ &= \delta_j^k g^{j\ell}(x, y) \frac{\partial f(x)}{\partial x^\ell} \frac{\partial f(x)}{\partial x^k} - g^{ik}(x, y) \frac{\partial f(x)}{\partial x^i} \frac{\partial f(x)}{\partial x^k} g^{j\ell}(x, y) \frac{\partial f(x)}{\partial x^j} \frac{\partial f(x)}{\partial x^\ell} = \\ &= \|df(x)\|_y^2 (1 - \|df(x)\|_y^2). \end{aligned}$$

Thus $\|df(x)\|_y < 1, \forall y$.

Now suppose that $\|df(x)\|_y < 1, \forall y$. Then for any vector field $X(x) = X^i(x) \frac{\partial}{\partial x^i}$ we have

$$\begin{aligned}
\tilde{g}_{ij}(x, y)X^i(x)X^j(x) &= g_{ij}(x, y)X^i(x)X^j(x) - \frac{\partial f(x)}{\partial x^i}X^i(x)\frac{\partial f(x)}{\partial x^j}X^j(x) = \\
&= \|X(x)\|_y^2 - \delta_k^i \frac{\partial f(x)}{\partial x^i}X^k(x)\delta_\ell^j \frac{\partial f(x)}{\partial x^j}X^\ell(x) = \|X(x)\|_y^2 - \\
&- g_{km}(x, y)g^{im}(x, y)\frac{\partial f(x)}{\partial x^i}X^k(x)g_{\ell s}(x, y)g^{js}(x, y)\frac{\partial f(x)}{\partial x^j}X^\ell(x) = \\
&= \|X(x)\|_y^2 - (g_{km}(x, y)g^{im}(x, y)\frac{\partial f(x)}{\partial x^i}X^k(x))^2 \geq \\
&\geq \|X(x)\|_y^2 - (g_{km}(x, y)X^k(x)X^m(x))^2 \left(g^{is}(x, y)\frac{\partial f(x)}{\partial x^i}\frac{\partial f(x)}{\partial x^s} \right)^2 = \\
&= \|X(x)\|_y^2(1 - \|df(x)\|_y^2),
\end{aligned}$$

and hence $\tilde{g}(x, y)$ is positive definite.

From theorems 2.2 and 2.4 follows

Theorem 2.5. *A Finsler C^3 -manifold $(M, g(x, y))$ which supports a proper C^3 function $f : M \rightarrow \mathbb{R}$ such that $\|df(x)\|_y < 1, \forall y$, is complete.*

Theorem 2.6. *Let $(M, g(x, y))$ be a Finsler C^3 -manifold and $f : M \rightarrow \mathbb{R}$ a proper C^3 function. Then*

$$(M, \tilde{g}(x, y) = e^{\|df(x)\|_y^2} g(x, y))$$

is a complete Finsler manifold.

PROOF. Obviously $\tilde{g}(x, y)$ is symmetric, positive definite and its components $\tilde{g}_{ij}(x, y)$ are homogeneous functions of degree zero with respect to y . On the other hand

$$\|\widetilde{df(x)}\|_y^2 = e^{-\|df(x)\|_y^2} \|df(x)\|_y^2 \leq \frac{1}{e}.$$

So the proper function $\varphi : M \rightarrow \mathbb{R}$, $\varphi = \sqrt{e}f$ satisfies $\|\widetilde{d\varphi}\|_y^2 < 1, \forall y$. Applying theorem 2.5 we obtain the desired result.

§3. Warped products of complete Finsler manifolds

Let $(M_1, g_1(x_1, y_1))$ and $(M_2, g_2(x_2, y_2))$ be Finsler manifolds and $f > 0$ a differentiable function on M_1 . Consider the product manifold $M_1 \times M_2$ with its projections

$$\pi : M_1 \times M_2 \rightarrow M_1, \quad \eta : M_1 \times M_2 \rightarrow M_2.$$

The Finsler manifold

$$(M_1 \times M_2, g_1 + fg_2)$$

is called the warped product between (M_1, g_1) and (M_2, g_2) .

Theorem 3.1. $(M_1 \times M_2, g_1 + fg_2)$ is complete if and only if (M_1, g_1) and (M_2, g_2) are complete.

PROOF. If $(M_1 \times M_2, g_1 + fg_2)$ is complete, then a Cauchy sequence in (M_1, g_1) or (M_2, g_2) imbeds in a (horizontal) leaf or a (vertical) fiber as a Cauchy sequence, and hence converges.

If (M_1, g_1) and (M_2, g_2) are complete, let $\{p_i = (p_{1i}, p_{2i})\}$ be a Cauchy sequence in $(M_1 \times M_2, g_1 + fg_2)$. Denote by α_{ij} a curve from p_i to p_j in $(M_1 \times M_2, g_1 + fg_2)$ having length at most $2d(p_i, p_j)$. We can assume that all projections $\pi \circ \alpha_{ij}$ lie in a compact region in M_1 , and on this we have $f \geq c > 0$. Consequently the speed of α_{ij} at each point is at least c times the speed of $\eta \circ \alpha_{ij}$. Thus

$$d(p_{2i}, p_{2j}) \leq \frac{2}{c} d(p_i, p_j)$$

showing that $\{p_{2i}\}$ is Cauchy and hence convergent.

Since π is distance-nonincreasing, the sequence $\{p_{1i}\}$ is also Cauchy, hence convergent. Thus $\{p_i\}$ is convergent, and $(M_1 \times M_2, g_1 + fg_2)$ is complete.

References

- [1] W. B. GORDON, An analytical criterion for the completeness of Riemannian manifolds, *Proc. Amer. Math. Soc.* **37**, 1 (1973), 221–225.
- [2] B. T. M. HASSAN, The theory of geodesics in Finsler spaces, Thesis, Southampton, 1967.
- [3] T. KAWAGUCHI and R. MIRON, On the generalized Lagrange spaces with the metric $\gamma_{ij}(x) + c^{-2}y_i y_j$, *Tensor, N.S.* **48** (1989), 52–63.
- [4] M. MATSUMOTO, Foundations of Finsler Geometry and Special Finsler Spaces, *Kyoto*, 1982.
- [5] R. MIRON and M. ANASTASIEI, Fibrat vectoriale. Spații Lagrange. Aplicații în teoria relativității, *Editura Academiei, București*, 1987.
- [6] C. UDRÎȘTE, Convexity of Finsler energies, *An. St. Univ. Al. I. Cuza, Iași*, **XXX** (1984–4), 91–94.
- [7] C. UDRÎȘTE and O. SANDRU, An analytical criterion for the completeness of Finsler manifolds, preprint.

CONSTANTIN UDRÎȘTE
DEPARTMENT OF MATHEMATICS I
POLYTECHNIC INSTITUTE
BUCHAREST
ROMANIA

(Received August 28, 1991; revised April 13, 1992)