

## The properties of $T^*$ -groups

By PIROSKA CSÖRGŐ (Budapest)

**Abstract.** This paper is a continuation of [1]. The aim of this work is the generalization of Zacher's theorem [2] for the solvable  $T^*$ -groups, a characterization of these groups by the normalizers of  $p$ -subgroups and the study of subnormal  $p$ -subgroups and of the Sylow subgroups of a solvable  $T^*$ -group.

Throughout this paper  $G$  will denote a finite group. A  $T$ -group is a group  $G$  whose subnormal subgroups are normal in  $G$ . In [1] M. ASAAD introduced the concept of  $T^*$ -group. A group  $G$  is called  $T^*$ -group if every subnormal subgroup of  $G$  is quasinormal in  $G$ . A subgroup of  $G$  is quasinormal in  $G$  if it permutes with every Sylow subgroup of  $G$ . In [1] we proved theorems concerning  $T$ -groups for  $T^*$ -groups. Now we are continuing the study of solvable  $T^*$ -groups. The notation used in this paper is standard.

First we generalize a theorem of ZACHER [2] for solvable  $T^*$ -groups.

**Theorem 1.** *Let  $G$  be a solvable group, the prime divisors of its order  $p_1 > p_2 > \cdots > p_k$ , and let  $P_1, \dots, P_k$  be a Sylow system with  $P_i \in \text{Syl}_{p_i}(G)$ .  $G$  is a  $T^*$ -group if and only if it satisfies the following conditions:*

- (i) *If  $1 \leq i < j \leq k$ , then  $P_j \leq N_G(P_i)$ .*
- (ii) *For all  $1 \leq i < j \leq k$ , if  $x \in P_i$ ,  $y \in P_j$  then there exists a natural number  $n$  such that  $x^y = x^n$ .*

PROOF. 1. Suppose  $G$  is a solvable  $T^*$ -group. Then by Lemma 1 of [1]  $G$  is supersolvable whence it has a Sylow tower. So  $G$  satisfies (i). As every subgroup of a solvable  $T^*$ -group is again a  $T^*$ -group by Theorem 1

of [1], it follows that  $N_G(P_i)$  is a solvable  $T^*$ -group. We have that  $\langle x \rangle$  is subnormal in  $N_G(P_i)$ , and using Lemma 2 of [1]  $P_j < N_G(\langle x \rangle)$  is true. Thus  $G$  satisfies (ii).

2. Conversely, assume  $G$  satisfies (i) and (ii). We show that every subgroup of  $P_i$  is quasinormal in  $N_G(P_i)$ . Let  $B$  be an arbitrary subgroup of  $P_i$ . By the conditions  $P_j < N_G(B)$  for all  $j > i$ . As  $P_j^y < N_G(B^y)$  for every  $y \in N_G(P_i)$ , clearly any Sylow  $p_j$ -subgroup of  $N_G(P_i)$  normalizes any subgroup of  $P_i$ . Let  $\ell < i$  and let  $D$  be an arbitrary Sylow  $p_\ell$ -subgroup of  $N_G(P_i)$ . By Hall's theorems  $(P_i D)^z \leq P_\ell P_i$  for some  $z \in G$ . As  $(P_i D)^z = P_i^z D^z$ ,  $D^z \leq P_\ell$  and  $P_i < N_G(P_\ell)$  clearly  $P_i^z < N_G(D^z)$  follows, furthermore  $D^z \leq N_G(P_i^z)$ , whence  $D$  centralizes  $P_i$ .

Thus every subgroup of  $P_i$  is quasinormal in  $N_G(P_i)$ . Using Lemma 4 of [1] it follows that either  $P_i \leq G'$  or each Sylow  $q$ -subgroup ( $q \neq p_i$ ) of  $N_G(P_i)$  centralizes  $P_i$ . So we can repeat the first part of the proof of Theorem 2 in [1]. Consequently  $G = HK$ , where  $H$  is a nilpotent normal Hall subgroup of  $G$ ,  $K$  is a nilpotent Hall subgroup of  $G$ ,  $H \cap K = 1$ , furthermore for arbitrary  $x \in H$ ,  $y \in K$  there exists a natural number  $i$  such that  $x^y = x^i$ . Then  $G$  is a solvable  $T^*$ -group by Theorem 2 of [1].

We need the following

**Lemma.** *Let  $U$  be a  $p$ -subgroup of  $G$ ,  $a \in N_G(U)$  such that  $(|a|, |U|) = 1$  furthermore  $a$  normalizes every subgroup of  $U$ . If there is an element  $b \neq 1$  of  $U$  such that  $ab = ba$ , then  $a \in C_G(U)$  follows.*

PROOF. Clearly  $C_U(a) \neq 1$ . Assume  $C_U(a) \neq U$ .

(a)  $Z(\Omega_1(U)) \not\leq C_U(a)$ .

Denote  $W = Z(\Omega_1(U))C_U(a)$ . Clearly  $\langle a \rangle$  normalizes every subgroup of  $W$  and each element of  $\langle a \rangle$  induces the identity on  $W/Z(\Omega_1(U))$  by conjugation. Applying Lemma 3 of [1]  $\langle a \rangle \leq C_G(W)$  follows, a contradiction.

(b)  $Z(\Omega_1(U)) > C_U(a)$ .

Clearly there exists a subgroup  $T \neq 1$  such that  $Z(\Omega_1(U)) = C_U(a) \times T$ . Let  $b \in T$ ,  $b \neq 1$  and  $u \in C_U(a)$ ,  $u \neq 1$ . Clearly  $a$  normalizes  $\langle b \rangle$  and  $\langle bu \rangle$ , consequently  $(bu)^a = (bu)^m$  where  $2 \leq m \leq p-1$ . As  $(bu)^a = b^a u = b^m u^m$ ,  $u^{m-1} = (b^m)^{-1} b^a$  follows. We have  $b^a = b^n$  where  $2 \leq n \leq p-1$  so  $u^{m-1} = b^{n-m}$  is true, but  $\langle u \rangle \cap \langle b \rangle = 1$  thus  $u^{m-1} = 1$ , a contradiction.

So  $Z(\Omega_1(U)) = C_U(a)$ .

(c)  $Z(\Omega_1(U)) < \Omega_1(U)$ .

Similarly to case (b) we can show that this case is impossible too.

Thus  $C_U(a) = Z(\Omega_1(U)) = \Omega_1(U)$ . Let  $\ell \in U \setminus \Omega_1(U)$ . By the conditions  $a$  normalizes  $\langle \ell \rangle$ . As  $a$  centralizes  $\Omega_1(\langle \ell \rangle)$  consequently  $a$  centralizes  $\langle \ell \rangle$ . A contradiction.

**Theorem 2.**  *$G$  is a solvable  $T^*$ -group if and only if every  $p$ -subgroup  $A$  (for all prime divisors  $p$  of the order of  $G$ ) is quasinormal in  $N_G(P_0)$  where  $P_0$  is a  $p$ -subgroup containing the subgroup  $A$ .*

PROOF. Assume  $G$  is a solvable  $T^*$ -group. Then by Theorem 1 of [1]  $N_G(P_0)$  is a solvable  $T^*$ -group too. Clearly  $A$  is subnormal in  $N_G(P_0)$ , whence  $A$  is quasinormal in  $N_G(P_0)$ .

Conversely, let  $p_1$  be the smallest prime divisor of the order of  $G$ . We show that  $G$  has a normal  $p_1$ -complement. Let  $P_1$  be a Sylow  $p_1$ -subgroup of  $G$  and let  $H$  be an arbitrary subgroup of  $P_1$ . We prove that  $N_G(H)/C_G(H)$  is a  $p_1$ -group. Assume there is an element  $b$  of  $N_G(H) \setminus C_G(H)$  of order  $q$  with  $q \neq p_1$ . Let  $a$  be an element of  $H$  of order  $p_1$ . By the conditions  $\langle a \rangle$  is quasinormal in  $N_G(H)$ . It is easy to see  $b \in N_H(\langle a \rangle)$ . As  $q > p_1$ ,  $b \in C_G(a)$  follows. Clearly every subgroup of  $H$  is quasinormal in  $N_G(H)$  by the conditions, whence  $b$  normalizes every subgroup of  $H$ . Using our Lemma  $b \in C_G(H)$  is true, a contradiction. Thus  $N_G(H)/C_G(H)$  is a  $p_1$ -group, consequently  $G$  has a normal  $p_1$ -complement. So  $G = P_1K$ ,  $K \triangleleft G$  and  $P_1 \cap K = 1$ . Consider the smallest prime divisor  $p_2$  of the order of  $K$ . Similarly we can prove that  $K$  has a normal  $p_2$ -complement.

Thus  $G$  has a tower such that the prime divisors of the order of  $G$  are  $p_1 < p_2 < \dots < p_k$  and for arbitrary  $1 \leq i \leq k$  there is a Sylow  $p_i$ -subgroup such that  $P_i < N_G(P_j)$  for all  $1 \leq i < j \leq k$ . If  $i$  and  $j$  are such as above and  $x \in P_j$ ,  $y \in P_i$ , then  $\langle x \rangle$  is quasinormal in  $N_G(P_j)$  by the conditions, whence it is easy to see that  $y \in N_G(\langle x \rangle)$ , consequently  $x^y = x^n$  for some natural number  $n$ . Applying Theorem 1  $G$  is a solvable  $T^*$ -group.

**Theorem 3.** *Let  $G$  be a solvable  $T^*$ -group. Then an arbitrary subnormal  $p$ -subgroup of  $G$  (for all prime divisors  $p$  of the order of  $G$ ) is either normal or it is centralized by all Sylow  $q$ -subgroups of  $G$  with  $q \neq p$ .*

PROOF. Let  $A$  be a subnormal  $p$ -subgroup of  $G$ . By Theorem 7 of [1]  $G = MN$  where  $M$  is a nilpotent normal Hall subgroup of  $G$ ,  $N$  is a nilpotent Hall subgroup of  $G$ ,  $M \cap N = 1$  furthermore every subgroup of prime power order of  $M$  is normal in  $G$

(a)  $A \leq M$ . By the above  $A$  is normal in  $G$

(b)  $A \leq N^y$  for some  $y \in G$ .

Let  $Q$  be a Sylow  $q$ -subgroup of  $M$  with  $q \neq p$ . By the subnormality of  $A$  there is a chain  $A \triangleleft A_1 \triangleleft \dots \triangleleft A_\ell \triangleleft A_{\ell+1} \triangleleft \dots \triangleleft A_m = G$ .

Let  $A_\ell$  be such that  $Q \leq A_\ell$  but  $Q \not\leq A_{\ell-1}$ .  $A$  normalizes every subgroup of  $Q$ . Since  $A_{\ell-1} \triangleleft A_\ell$ ,  $Q \triangleleft A_\ell$  and  $A \leq A_{\ell-1}$  it follows that each element of  $A$  induces the identity on  $Q/Q \cap A_{\ell-1}$  by conjugation. Using Lemma 3 of [1]  $A \leq C_G(Q)$  follows. As  $Q$  is an arbitrary Sylow subgroup of  $M$ ,  $A \leq C_G(M)$  is true. We have  $G = M \cdot N^y$ ,  $N^y$  is a nilpotent Hall subgroup of  $G$  and  $N^y = P \times T$  where  $P$  is a Sylow  $p$ -subgroup of  $G$ , whence  $C_G(M) \geq M \cdot T$ . As  $MT \triangleleft G$  it is easy to see that  $A$  is centralized by an arbitrary Sylow  $q$ -subgroup of  $G$  with  $q \neq p$ .

**Theorem 4.**  *$G$  is a solvable  $T^*$ -group if and only if every Sylow subgroup  $P$  satisfies one of the following conditions:*

- (a) *every subgroup of  $P$  is normal in  $G$*
- (b) *every Sylow subgroup of  $N_G(P)$  different from  $P$  centralizes  $P$ .*

PROOF. Assume  $G$  is a solvable  $T^*$ -group. By the Theorem 7 of [1]  $G = MN$  where  $M$  is a nilpotent normal Hall subgroup of  $G$ ,  $N$  is a nilpotent Hall subgroup of  $G$ ,  $M \cap N = 1$  and every subgroup of prime power order of  $M$  is normal in  $G$ . Let  $R$  be an arbitrary Sylow subgroup of  $G$ .

Assume  $R \leq M$ . By the above every subgroup of  $R$  is normal in  $G$ .

Assume  $R \leq N^y$  for some  $y \in G$ . Clearly  $N_G(R) = N^y \cdot (N_G(R) \cap M)$ . The structure of  $G$  yields  $B = N_G(R) \cap M \leq C_G(R)$  and  $N^y = R \times L$  where  $L$  is a nilpotent Hall subgroup of  $G$ , consequently  $N_G(R) = R \times (L \cdot B)$  so  $R$  satisfies (b).

Conversely, let  $G$  be a counterexample of smallest order. Let  $M_0$  be the product of every Sylow subgroup of  $G$  each subgroup of which is normal in  $G$ . Clearly  $M_0$  is a nilpotent normal Hall subgroup of  $G$ . By the Theorem of Zassenhaus there is a subgroup  $N_0$  such that  $M_0 \cdot N_0 = G$  and  $M_0 \cap N_0 = 1$ . Clearly  $N_0$  is a Hall subgroup in  $G$  and it satisfies the conditions of our theorem. By the minimality of  $G$   $N_0$  is a solvable  $T^*$ -group. Using Theorem 2 of [1]  $N_0 = A \cdot B$  where  $A$  is a nilpotent normal Hall subgroup of  $G$ ,  $B$  is a nilpotent Hall subgroup of  $G$  and  $A \cap B = 1$  furthermore for arbitrary  $a \in A$ ,  $b \in B$  there is a natural number  $i$  such that  $a^b = a^i$ . Assume  $A \neq 1$ . Let  $P$  be a Sylow subgroup of  $A$ . Using  $N_{N_0}(P) = N_0$ ,  $B \leq C_G(P)$  follows by the conditions whence  $B \leq C_G(A)$ . Thus  $N_0 = A \times B$  and  $N_0$  is a nilpotent normal Hall subgroup of  $G$ . Using Theorem 2 of [1] it follows that  $G$  is a solvable  $T^*$ -group.

**References**

- [1] M. ASAAD and P. CSÖRGŐ, On  $T^*$ -groups, (*submitted to Acta Mathematica Hungarica*).
- [2] G. ZACHER, Caratterizzazione dei  $t$ -gruppi finiti risolubili, *Ricerche Mat.* **1** (1952), 287–294.

PIROSKA CSÖRGŐ  
EÖTVÖS UNIVERSITY  
DEPARTMENT OF ALGEBRA  
MÚZEUM KRT. 6–8., 1088 BUDAPEST  
HUNGARY

*(Received July 24, 1995)*