Publ. Math. Debrecen
69/1-2 (2006), 227–243

Hybrid mean value on the difference between a quadratic residue and its inverse modulo p

By LIU HUANING (Xi'an) and ZHANG WENPENG (Xi'an)

Abstract. The main purpose of this paper is using the generalized Bernoulli numbers, Gauss sums and the mean value theorems of Dirichlet L-functions to study the asymptotic property of the difference between a quadratic residue and its inverse modulo p (a prime), and give an interesting hybrid mean value formula.

1. Introduction

Let $q > 2$ and c are two integers with $(c, q) = 1$. For each integer $1 \le a \le q$ with $(a, q) = 1$, we know that there exists one and only one integer $1 \le b \le q$ with $(b, q) = 1$ such that $ab \equiv c \bmod q$. Let

$$M(q, k, c) = \sum_{\substack{a=1 \\ ab \equiv c}}^{q}{}' \sum_{\substack{b=1 \\ \bmod q}}^{q}{}' (a - b)^{2k},$$

where $\sum_{a=1}^{q}{}'$ denotes the summation over all a such that $(a, q) = 1$. In [3], the second author used the estimates for Kloosterman sums and trigonometric sums to obtain a sharp asymptotic formula for $M(q, k, c)$, and prove the following:

Mathematics Subject Classification: 11L05, 11A07.
Key words and phrases: modular inverses, Bernoulli numbers, Gauss sums.
This work is supported by the N.S.F. and P.N.S.F. of P.R. China.

Proposition 1. *Let $q > 2$ and c are two integers with $(c, q) = 1$. Then for any positive integer k, we have the asymptotic formula*

$$M(q,k,c) = \frac{1}{(2k+1)(k+1)}\phi(q)q^{2k} + O\left(4^k q^{\frac{4k+1}{2}} d^2(q)\ln^2 q\right),$$

where $\phi(q)$ is the Euler function, and $d(q)$ is the divisor function.

The error term in Proposition 1 is the best possible. In fact for $k = 1$, let

$$M(q,1,c) = \frac{1}{6}\phi(q)q^2 + \frac{1}{3}q\prod_{p|q}(1-p) + F(q,1,c),$$

where $\prod_{p|q}$ denotes the product over all distinct prime divisors of q. The second author [5] used the properties of Dedekind sums and Cochrane sums to give a sharp mean value formula for $F(q,1,c)$. That is the following:

Proposition 2. *For any integer $q > 2$, we have the asymptotic formula*

$$\sum_{c=1}^{q}{}' F^2(q,1,c) = \frac{5}{36}q^3\phi^3(q)\prod_{p^\alpha \| q}\frac{\frac{(p+1)^3}{p(p^2+1)} - \frac{1}{p^{3\alpha-1}}}{1+\frac{1}{p}+\frac{1}{p^2}} + O\left(q^5\exp\left(\frac{4\ln q}{\ln\ln q}\right)\right),$$

where $\prod_{p^\alpha \| q}$ denotes the product over all prime divisors of q with $p^\alpha | q$ and $p^{\alpha+1} \nmid q$.

Now we consider the difference between a quadratic residue and its inverse modulo q. Let

$$N(q,k,c) = \sum_{\substack{a=1\\ a\in A\\ ab\equiv c \bmod q}}^{q}{}' \sum_{\substack{b=1\\ b\in A}}^{q}{}' (a-b)^{2k},$$

where A denotes the set of all quadratic residues modulo q. Denote

$$\begin{aligned} N(q,k,c) = {} & \frac{1}{2\phi(q)}\sum_{a=1}^{q}{}'\sum_{b=1}^{q}{}'(a-b)^{2k} \\ & + \frac{1}{\phi(q)}\sum_{a=1}^{q}{}'\sum_{b=1}^{q}{}'\chi_2(b)(a-b)^{2k} + E(q,k,c), \end{aligned}$$

where χ_2 is the quadratic character modulo q.

It is interesting that there exists some relation between $E(q,k,c)$ and the classical Kloosterman sums

$$K(m,n;q)=\sum_{a=1}^{q}{}'\,\mathrm{e}\left(\frac{ma+n\overline{a}}{q}\right),$$

where $\mathrm{e}(y)=\mathrm{e}^{2\pi iy}$, $\overline{a}$ is defined by the equation $a\overline{a}\equiv 1 \bmod q$. In this paper, we use the generalized Bernoulli numbers, Gauss sums and the mean value theorems of Dirichlet L-functions to study the hybrid mean value of $\sum\limits_{\substack{c=1\\ c\in A}}^{p-1} E(p,k,c)K(c,1;p)$, and give an interesting asymptotic formula. That is, we shall prove the following:

Theorem 1. *For any prime $p\geq 5$ and integer $k\geq 1$, we have the following:*

$$\begin{aligned}&\sum_{\substack{c=1\\ c\in A}}^{p-1} E(p,k,c)K(c,1;p)\\ &\quad=2p^{2k+2}\sum_{n=2}^{2k-2}C_{2k}^{n}\sum_{m=1}^{[\frac{n}{2}]}\sum_{l=1}^{[\frac{2k-n}{2}]}\frac{C_n^{2m-1}C_{2k-n}^{2l-1}(2m-1)!(2l-1)!}{(-1)^{n+m+l}(2\pi)^{2m+2l}}\\ &\qquad-2p^{2k+2}\sum_{n=1}^{2k-1}C_{2k}^{n}\sum_{m=0}^{[\frac{n-1}{2}]}\sum_{l=0}^{[\frac{2k-n-1}{2}]}\frac{C_n^{2m}C_{2k-n}^{2l}(2m)!(2l)!}{(-1)^{n+m+l}(2\pi)^{2m+2l+2}}+O\big(p^{2k+\frac{3}{2}+\epsilon}\big),\end{aligned}$$

where $C_n^m=\frac{n!}{m!(n-m)!}$ is the combinatorial number, ϵ is any fixed positive number, and $[y]$ denotes the greatest integer not more than y.

For general integers $q\geq 5$ and $k\geq 1$, whether there exists an asymptotic formula for

$$\sum_{\substack{c=1\\ c\in A}}^{q}{}'\,E(q,k,c)K(c,1;q)$$

is an open problem.

2. Some lemmas

To complete the proof of Theorem 1, we need the following lemmas.

Lemma 1. *Let q be a positive integer, then for any character χ modulo q we have*

$$\sum_{\substack{c=1\\c\in A}}^{q}\overline{\chi}(c)K(c,1;q)=\frac{1}{2}\tau^2(\overline{\chi})+\frac{1}{2}\tau^2(\overline{\chi}\chi_2),$$

where $\tau(\chi)=\sum_{a=1}^{q}\chi(a)e(\frac{a}{q})$ is the Gauss sum.

Proof. Noting that (see Lemma 2 of [6])

$$\sum_{c=1}^{q}\overline{\chi}(c)K(c,1;q)=\tau^2(\overline{\chi}),$$

so we have

$$\begin{aligned}\sum_{\substack{c=1\\c\in A}}^{q}\overline{\chi}(c)K(c,1;q)&=\frac{1}{2}\sum_{c=1}^{q}[1+\chi_2(c)]\overline{\chi}(c)K(c,1;q)\\&=\frac{1}{2}\tau^2(\overline{\chi})+\frac{1}{2}\tau^2(\overline{\chi}\chi_2).\end{aligned}$$

This proves Lemma 1. □

Lemma 2. *Let integers $q>2$ and $n>0$, then for any primitive character χ modulo q we have*

$$\sum_{a=1}^{q}a^n\chi(a)=\begin{cases}2q^n\tau(\chi)\displaystyle\sum_{m=1}^{[\frac{n}{2}]}\frac{C_n^{2m-1}(2m-1)!L(2m,\overline{\chi})}{(-1)^{m+1}(2\pi)^{2m}}, & \text{if}\quad \chi(-1)=1;\\[2ex] 2q^n\tau(\chi)\displaystyle\sum_{m=0}^{[\frac{n-1}{2}]}\frac{C_n^{2m}(2m)!L(2m+1,\overline{\chi})}{(-1)^{m+1}(2\pi)^{2m+1}i}, & \text{if}\quad \chi(-1)=-1,\end{cases}$$

where $L(s,\chi)$ is the Dirichlet L-function corresponding to χ.

PROOF. Usually, the generalized Bernoulli numbers $B_{k,\chi}$ are defined as following:

$$\sum_{a=1}^{q}\chi(a)\frac{te^{at}}{e^{qt}-1}=\sum_{k=0}^{\infty}\frac{B_{k,\chi}}{k!}t^k.$$

Let

$$B_{n,\chi}(x)=\sum_{k=0}^{n}C_n^kB_{k,\chi}x^{n-k},$$

then from [1] we know that

$$\begin{aligned}\sum_{a=1}^{q}a^n\chi(a)&=\frac{1}{n+1}\left(B_{n+1,\chi}(q)-B_{n+1,\chi}(0)\right)\\&=\frac{1}{n+1}\sum_{k=0}^{n}C_{n+1}^kB_{k,\chi}q^{n+1-k}.\end{aligned}\tag{1}$$

For $\chi(-1)=1$, from (1) we have

$$\sum_{a=1}^{q}a^n\chi(a)=\frac{1}{n+1}\sum_{m=1}^{[\frac{n}{2}]}C_{n+1}^{2m}B_{2m,\chi}q^{n+1-2m}.\tag{2}$$

Noting that (see [1])

$$L(2m,\overline{\chi})=\frac{(-1)^{m+1}\tau(\overline{\chi})}{2(2m)!}\left(\frac{2\pi}{q}\right)^{2m}B_{2m,\chi}\tag{3}$$

for any positive integer m, so from (2) and (3) we have

$$\sum_{a=1}^{q}a^n\chi(a)=2q^n\tau(\chi)\sum_{m=1}^{[\frac{n}{2}]}\frac{C_n^{2m-1}(2m-1)!L(2m,\overline{\chi})}{(-1)^{m+1}(2\pi)^{2m}},\quad\text{if }\chi(-1)=1.$$

For $\chi(-1)=-1$, it follows from (1) that

$$\sum_{a=1}^{q}a^n\chi(a)=B_{1,\chi}q^n+\frac{1}{n+1}\sum_{m=1}^{[\frac{n-1}{2}]}C_{n+1}^{2m+1}B_{2m+1,\chi}q^{n-2m}.\tag{4}$$

Noting that (see [1])

$$B_{1,\chi} = \frac{i\tau(\chi)L(1,\overline{\chi})}{\pi}, \tag{5}$$

and for any positive integer m,

$$L(2m+1,\overline{\chi}) = \frac{i(-1)^m\tau(\overline{\chi})}{2(2m+1)!}\left(\frac{2\pi}{q}\right)^{2m+1} B_{2m+1,\chi}, \tag{6}$$

now combining (4), (5) and (6) we have

$$\sum_{a=1}^{q} a^n\chi(a) = 2q^n\tau(\chi)\sum_{m=0}^{[\frac{n-1}{2}]}\frac{C_n^{2m}(2m)!L(2m+1,\overline{\chi})}{(-1)^{m+1}(2\pi)^{2m+1}i}, \quad \text{if } \chi(-1)=-1.$$

This completes the proof of Lemma 2. □

Lemma 3. *Let q and r be integers with $q \geq 2$ and $(r,q)=1$, χ be a Dirichlet character modulo q. Then we have the identities*

$$\sum_{\chi \bmod q}^{*}\chi(r) = \sum_{d|(q,r-1)}\mu\left(\frac{q}{d}\right)\phi(d)$$

and

$$J(q) = \sum_{d|q}\mu(d)\phi\left(\frac{q}{d}\right),$$

where $\sum_{\chi \bmod q}^{*}$ denotes the summation over all primitive characters modulo q, $\mu(n)$ is the Möbius function and $J(q)$ denotes the number of primitive characters mod q.

Proof. This is Lemma 3 of [4]. □

Lemma 4. *Let q be any integer with $q > 2$, and $u, v \geq 1$ be real numbers. Then we have*

(I) $$\sum_{\chi(-1)=1}^{*} L(u,\overline{\chi})L(v,\overline{\chi}) = \frac{J(q)}{2} + O\left(q^{\epsilon}\right);$$

(II) $$\sum_{\chi(-1)=-1}^{*} L(u,\overline{\chi})L(v,\overline{\chi}) = \frac{J(q)}{2} + O\left(q^{\epsilon}\right).$$

PROOF. We only prove (II), since similarly we can deduce (I). For any non-principal character χ modulo q, and parameter $N \geq q$, applying Abel's identity we have

$$L(u,\overline{\chi}) = \sum_{n=1}^{+\infty} \frac{\overline{\chi}(n)}{n^u} = \sum_{1\leq n\leq N} \frac{\overline{\chi}(n)}{n^u} + u\int_N^{+\infty} \frac{\sum_{N<n\leq y}\overline{\chi}(n)}{y^{u+1}}\mathrm{d}y$$
$$= \sum_{1\leq n\leq N} \frac{\overline{\chi}(n)}{n^u} + O\left(\frac{\sqrt{q}\log q}{N^u}\right). \tag{7}$$

Let $\sigma_\alpha(n) = \sum_{d|n} d^\alpha$. For $(a,q)=1$, by Lemma 3 we have

$$\sum_{\chi(-1)=-1}^{*} \chi(a) = \frac{1}{2}\sum_{\chi \bmod q}^{*} (1-\chi(-1))\chi(a) = \frac{1}{2}\sum_{\chi \bmod q}^{*} \chi(a) - \frac{1}{2}\sum_{\chi \bmod q}^{*} \chi(-a)$$
$$= \frac{1}{2}\sum_{d|(q,a-1)} \mu\left(\frac{q}{d}\right)\phi(d) - \frac{1}{2}\sum_{d|(q,a+1)} \mu\left(\frac{q}{d}\right)\phi(d).$$

Therefore

$$\sum_{\chi(-1)=-1}^{*} L(u,\overline{\chi})L(v,\overline{\chi})$$
$$= \sum_{\chi(-1)=-1}^{*} \left(\sum_{1\leq n\leq N} \frac{\overline{\chi}(n)}{n^u} + O\left(\frac{\sqrt{q}\log q}{N^u}\right)\right)\left(\sum_{1\leq m\leq N} \frac{\overline{\chi}(m)}{m^v} + O\left(\frac{\sqrt{q}\log q}{N^v}\right)\right)$$
$$= \sum_{\chi(-1)=-1}^{*} \sum_{1\leq n\leq N}\sum_{1\leq m\leq N} \frac{\overline{\chi}(n)\overline{\chi}(m)}{n^u m^v} + O\left(\frac{q^{\frac{3}{2}}\log^2 q}{N}\right)$$
$$= \sum_{1\leq k\leq N^2} \frac{\sigma_{u-v}(k)}{k^u} \sum_{\chi(-1)=-1}^{*} \overline{\chi}(k) + O\left(\frac{q^{\frac{3}{2}}\log^2 q}{N}\right)$$
$$= \frac{1}{2}\sum_{\substack{1\leq k\leq N^2\\(k,q)=1}} \frac{\sigma_{u-v}(k)}{k^u} \sum_{d|(q,k-1)} \mu\left(\frac{q}{d}\right)\phi(d)$$
$$- \frac{1}{2}\sum_{\substack{1\leq k\leq N^2\\(k,q)=1}} \frac{\sigma_{u-v}(k)}{k^u} \sum_{d|(q,k+1)} \mu\left(\frac{q}{d}\right)\phi(d) + O\left(\frac{q^{\frac{3}{2}}\log^2 q}{N}\right)$$

$$= \frac{1}{2}\sum_{d|q}\mu\left(\frac{q}{d}\right)\phi(d)\sum_{\substack{1\le k\le N^2\\(k,q)=1\\k\equiv 1 \bmod d}}\frac{\sigma_{u-v}(k)}{k^u}$$

$$-\frac{1}{2}\sum_{d|q}\mu\left(\frac{q}{d}\right)\phi(d)\sum_{\substack{1\le k\le N^2\\(k,q)=1\\k\equiv -1 \bmod d}}\frac{\sigma_{u-v}(k)}{k^u}+O\left(\frac{q^{\frac{3}{2}}\log^2 q}{N}\right)$$

$$= \frac{J(q)}{2}+O\left(\sum_{d|q}\phi(d)\sum_{1\le l\le\frac{N^2-1}{d}}\frac{\sigma_{u-v}(ld+1)}{(ld+1)^u}\right)$$

$$+O\left(\sum_{d|q}\phi(d)\sum_{1\le l\le\frac{N^2+1}{d}}\frac{\sigma_{u-v}(ld-1)}{(ld-1)^u}\right)+O\left(\frac{q^{\frac{3}{2}}\log^2 q}{N}\right)$$

$$= \frac{J(q)}{2}+O\left(\frac{q^{\frac{3}{2}}\log^2 q}{N}\right)+O\left(N^{\epsilon}\right).$$

Now taking $N = q^{\frac{3}{2}}$ in above, we immediately get

$$\sum_{\chi(-1)=-1}^{*} L(u,\overline{\chi})L(v,\overline{\chi}) = \frac{J(q)}{2}+O(q^{\epsilon}).$$

This proves Lemma 4. □

Lemma 5. *Let q be an odd prime power and let F be the field $GF(q)$. Let $f(y)$ be a non-constant polynomial over F which decomposes into the product of m distinct linear factors. Then, for the quadratic character χ,*

$$\left|\sum_{y\in F}\chi(f(y))\right| \le (m-1)q^{\frac{1}{2}}.$$

Proof. This is the well-known estimate for character sums, due to Weil [2]. □

Lemma 6. *Let prime $p \ge 5$, $u, v \ge 1$ be real numbers, χ_2 be the quadratic character modulo p. Then we have*

$$\text{(I)}\qquad \sum_{\substack{\chi\ne\chi_0\\\chi(-1)=1}}\tau^2(\overline{\chi}\chi_2)\tau^2(\chi)L(u,\overline{\chi})L(v,\overline{\chi}) \ll p^{\frac{5}{2}+\epsilon};$$

$$\text{(II)} \qquad \sum_{\chi(-1)=-1} \tau^2(\overline{\chi}\chi_2)\tau^2(\chi)L(u,\overline{\chi})L(v,\overline{\chi}) \ll p^{\frac{5}{2}+\epsilon}.$$

Proof. We only prove (II), since similarly we can deduce (I). From the properties of Gauss sums we have

$$\begin{aligned}\tau(\overline{\chi}\chi_2)\tau(\chi) &= \sum_{a=1}^{p-1}\overline{\chi}(a)\chi_2(a)\mathrm{e}\left(\frac{a}{p}\right)\sum_{b=1}^{p-1}\chi(b)\mathrm{e}\left(\frac{b}{p}\right)\\ &= \sum_{a=1}^{p-1}\chi_2(a)\mathrm{e}\left(\frac{a}{p}\right)\sum_{b=1}^{p-1}\chi(b)\mathrm{e}\left(\frac{ba}{p}\right)\\ &= \sum_{b=1}^{p-1}\chi(b)\sum_{a=1}^{p-1}\chi_2(a)\mathrm{e}\left(\frac{a(1+b)}{p}\right)\\ &= \tau(\chi_2)\sum_{b=1}^{p-1}\chi(b)\chi_2(1+b),\end{aligned}$$

therefore

$$\begin{aligned}\tau^2(\overline{\chi}\chi_2)\tau^2(\chi) &= \tau^2(\chi_2)\sum_{a=1}^{p-1}\chi(a)\chi_2(1+a)\sum_{b=1}^{p-1}\chi(b)\chi_2(1+b)\\ &= \tau^2(\chi_2)\sum_{b=1}^{p-1}\chi(b)\sum_{a=1}^{p-1}\chi_2(1+a)\chi_2(1+b\overline{a}).\end{aligned}$$

Noting that for $(ab,p)=1$, by the orthogonality relations for character sums modulo p we have

$$\sum_{\chi(-1)=-1}\chi(a)\overline{\chi}(b) = \begin{cases}\dfrac{1}{2}(p-1), & \text{if } a\equiv b \mod p;\\ -\dfrac{1}{2}(p-1), & \text{if } a\equiv -b \mod p;\\ 0, & \text{otherwise.}\end{cases} \tag{8}$$

Then for parameter $N \geq p$, by (7) and (8) we get

$$\sum_{\chi(-1)=-1}\tau^2(\overline{\chi}\chi_2)\tau^2(\chi)L(u,\overline{\chi})L(v,\overline{\chi})$$

$$= \sum_{\chi(-1)=-1} \tau^2(\overline{\chi}\chi_2)\tau^2(\chi) \sum_{1\le n\le N} \sum_{1\le m\le N} \frac{\overline{\chi}(n)\overline{\chi}(m)}{n^u m^v} + O\left(\frac{p^{\frac{7}{2}}\log^2 p}{N}\right)$$

$$= \tau^2(\chi_2) \sum_{1\le k\le N^2} \frac{\sigma_{u-v}(k)}{k^u} \sum_{b=1}^{p-1}\sum_{a=1}^{p-1} \chi_2(1+a)\chi_2(1+b\overline{a}) \sum_{\chi(-1)=-1} \chi(b)\overline{\chi}(k)$$
$$+ O\left(\frac{p^{\frac{7}{2}}\log^2 p}{N}\right)$$

$$= \frac{\tau^2(\chi_2)(p-1)}{2} \sum_{\substack{1\le k\le N^2\\(k,p)=1}} \frac{\sigma_{u-v}(k)}{k^u} \sum_{\substack{b=1\\ b\equiv k \bmod p}}^{p-1} \sum_{a=1}^{p-1} \chi_2(1+a)\chi_2(1+b\overline{a})$$
$$- \frac{\tau^2(\chi_2)(p-1)}{2} \sum_{\substack{1\le k\le N^2\\(k,p)=1}} \frac{\sigma_{u-v}(k)}{k^u} \sum_{\substack{b=1\\ b\equiv -k \bmod p}}^{p-1} \sum_{a=1}^{p-1} \chi_2(1+a)\chi_2(1+b\overline{a})$$
$$+ O\left(\frac{p^{\frac{7}{2}}\log^2 p}{N}\right).$$

On the other hand, by Lemma 5 we can get

$$\sum_{a=1}^{p-1} \chi_2(1+a)\chi_2(1+b\overline{a}) = \sum_{a=1}^{p-1} \chi_2(a^2)\chi_2(1+a)\chi_2(1+b\overline{a})$$
$$= \sum_{a=1}^{p-1} \chi_2(a)\chi_2(a^2+ab+a+b) \ll p^{\frac{1}{2}}.$$

So we have

$$\sum_{\chi(-1)=-1} \tau^2(\overline{\chi}\chi_2)\tau^2(\chi)L(u,\overline{\chi})L(v,\overline{\chi})$$
$$\ll p^{\frac{5}{2}} \sum_{1\le k\le N^2} \frac{\sigma_{u-v}(k)}{k^u} + \frac{p^{\frac{7}{2}}\log^2 p}{N} \ll p^{\frac{5}{2}}N^{\epsilon} + \frac{p^{\frac{7}{2}}\log^2 p}{N}.$$

Now taking $N = p$ in above, we immediately get

$$\sum_{\chi(-1)=-1} \tau^2(\overline{\chi}\chi_2)\tau^2(\chi)L(u,\overline{\chi})L(v,\overline{\chi}) \ll p^{\frac{5}{2}+\epsilon}.$$

This completes the proof of Lemma 6. □

Lemma 7. *Let prime $p > 2$, $u, v \geq 1$ be real numbers, χ_2 be the quadratic character modulo p. Then we have*

$$\text{(I)} \qquad \sum_{\substack{\chi \neq \chi_0 \\ \chi(-1)=1}} \tau(\overline{\chi})\tau(\chi\chi_2)L(u,\overline{\chi})L(v,\overline{\chi}\chi_2) \ll p^{\frac{3}{2}+\epsilon};$$

$$\text{(II)} \qquad \sum_{\chi(-1)=-1} \tau(\overline{\chi})\tau(\chi\chi_2)L(u,\overline{\chi})L(v,\overline{\chi}\chi_2) \ll p^{\frac{3}{2}+\epsilon}.$$

Proof. We only prove (II), since similarly we can deduce (I). From the properties of Gauss sums we have

$$\begin{aligned}\tau(\overline{\chi})\tau(\chi\chi_2) &= \sum_{a=1}^{p-1} \overline{\chi}(a)\mathrm{e}\left(\frac{a}{p}\right)\sum_{b=1}^{p-1}\chi(b)\chi_2(b)\mathrm{e}\left(\frac{b}{p}\right)\\ &= \sum_{a=1}^{p-1}\sum_{b=1}^{p-1}\chi(b)\chi_2(b)\chi_2(a)\mathrm{e}\left(\frac{a(1+b)}{p}\right)\\ &= \tau(\chi_2)\sum_{b=1}^{p-1}\chi(b)\chi_2(b)\chi_2(1+b).\end{aligned}$$

Let $\tau_\alpha(n) = \sum_{d|n}\chi_2(d)d^\alpha$. Then for $N \geq p$, by (7) and (8) we get

$$\begin{aligned}&\sum_{\chi(-1)=-1}\tau(\overline{\chi})\tau(\chi\chi_2)L(u,\overline{\chi})L(v,\overline{\chi}\chi_2)\\ &= \sum_{\chi(-1)=-1}\tau(\overline{\chi})\tau(\chi\chi_2)\sum_{1\leq n\leq N}\sum_{1\leq m\leq N}\frac{\overline{\chi}(n)\overline{\chi}(m)\chi_2(m)}{n^u m^v} + O\left(\frac{p^{\frac{5}{2}}\log^2 p}{N}\right)\\ &= \tau(\chi_2)\sum_{1\leq k\leq N^2}\frac{\tau_{u-v}(k)}{k^u}\sum_{b=1}^{p-1}\chi_2(b)\chi_2(1+b)\sum_{\chi(-1)=-1}\chi(b)\overline{\chi}(k) + O\left(\frac{p^{\frac{5}{2}}\log^2 p}{N}\right)\\ &= \frac{\tau(\chi_2)(p-1)}{2}\sum_{\substack{1\leq k\leq N^2\\(k,p)=1}}\frac{\tau_{u-v}(k)}{k^u}\sum_{\substack{b=1\\ b\equiv k \bmod p}}^{p-1}\chi_2(b)\chi_2(1+b)\\ &\quad - \frac{\tau(\chi_2)(p-1)}{2}\sum_{\substack{1\leq k\leq N^2\\(k,p)=1}}\frac{\tau_{u-v}(k)}{k^u}\sum_{\substack{b=1\\ b\equiv -k \bmod p}}^{p-1}\chi_2(b)\chi_2(1+b) + O\left(\frac{p^{\frac{5}{2}}\log^2 p}{N}\right)\end{aligned}$$

$$\ll p^{\frac{3}{2}} \sum_{1\le k\le N^2} \frac{\tau_{u-v}(k)}{k^u} + \frac{p^{\frac{5}{2}}\log^2 p}{N} \ll p^{\frac{3}{2}} N^{\epsilon} + \frac{p^{\frac{5}{2}}\log^2 p}{N}.$$

Now taking $N = p$ in above, we immediately get

$$\sum_{\chi(-1)=-1} \tau(\overline{\chi})\tau(\chi\chi_2)L(u,\overline{\chi})L(v,\overline{\chi}\chi_2) \ll p^{\frac{3}{2}+\epsilon}.$$

This proves Lemma 7. □

3. Proof of Theorem 1

In this section, we complete the proof of Theorem 1. For any prime $p \ge 5$ and integer $k \ge 1$, from the definition of $N(p,k,c)$ we have

$$\begin{aligned}
N(p,k,c) &= \sum_{\substack{a=1\\ a\in A\\ ab\equiv c \bmod p}}^{p-1} \sum_{\substack{b=1\\ b\in A}}^{p-1} (a-b)^{2k} \\
&= \frac{1}{4} \sum_{\substack{a=1\\ ab\equiv c \bmod p}}^{p-1} \sum_{b=1}^{p-1} (1+\chi_2(a))(1+\chi_2(b))(a-b)^{2k} \\
&= \frac{1}{4} \sum_{\substack{a=1\\ ab\equiv c \bmod p}}^{p-1} \sum_{b=1}^{p-1} (a-b)^{2k} + \frac{1}{2} \sum_{\substack{a=1\\ ab\equiv c \bmod p}}^{p-1} \sum_{b=1}^{p-1} \chi_2(b)(a-b)^{2k} \\
&\quad + \frac{1}{4} \sum_{\substack{a=1\\ ab\equiv c \bmod p}}^{p-1} \sum_{b=1}^{p-1} \chi_2(a)\chi_2(b)(a-b)^{2k} \\
&= \frac{1}{2(p-1)} \sum_{a=1}^{p-1}\sum_{b=1}^{p-1} (a-b)^{2k} + \frac{1}{(p-1)} \sum_{a=1}^{p-1} \sum_{b=1}^{p-1} \chi_2(b)\,(a-b)^{2k} \\
&\quad + \frac{1}{2(p-1)} \sum_{\chi\ne\chi_0} \overline{\chi}(c) \sum_{a=1}^{p-1} \sum_{b=1}^{p-1} \chi(a)\chi(b)(a-b)^{2k} \\
&\quad + \frac{1}{2(p-1)} \sum_{\chi^2\ne\chi_0} \overline{\chi}(c) \sum_{a=1}^{p-1} \sum_{b=1}^{p-1} \chi(a)\chi(b)\chi_2(b)\,(a-b)^{2k}.
\end{aligned}$$

That is,

$$E(p,k,c) = \frac{1}{2(p-1)} \sum_{\chi \neq \chi_0} \overline{\chi}(c) \sum_{a=1}^{p-1} \sum_{b=1}^{p-1} \chi(a)\chi(b) \, (a-b)^{2k}$$

$$+ \frac{1}{2(p-1)} \sum_{\chi^2 \neq \chi_0} \overline{\chi}(c) \sum_{a=1}^{p-1} \sum_{b=1}^{p-1} \chi(a)\chi(b)\chi_2(b) \, (a-b)^{2k} .$$

Then from Lemma 1 we have

$$\sum_{\substack{c=1 \\ c \in A}}^{p-1} E(p,k,c) K(c,1;p)$$

$$= \frac{1}{2(p-1)} \sum_{\chi \neq \chi_0} \left[\frac{\tau^2(\overline{\chi})}{2} + \frac{\tau^2(\overline{\chi}\chi_2)}{2} \right] \sum_{a=1}^{p-1} \sum_{b=1}^{p-1} \chi(a)\chi(b) \, (a-b)^{2k}$$

$$+ \frac{1}{2(p-1)} \sum_{\chi^2 \neq \chi_0} \left[\frac{\tau^2(\overline{\chi})}{2} + \frac{\tau^2(\overline{\chi}\chi_2)}{2} \right] \sum_{a=1}^{p-1} \sum_{b=1}^{p-1} \chi(a)\chi(b)\chi_2(b) \, (a-b)^{2k}$$

$$= \frac{1}{4(p-1)} \sum_{\chi \neq \chi_0} \tau^2(\overline{\chi}) \sum_{a=1}^{p-1} \sum_{b=1}^{p-1} \chi(a)\chi(b) \, (a-b)^{2k}$$

$$+ \frac{1}{4(p-1)} \sum_{\chi \neq \chi_0} \tau^2(\overline{\chi}\chi_2) \sum_{a=1}^{p-1} \sum_{b=1}^{p-1} \chi(a)\chi(b) \, (a-b)^{2k}$$

$$+ \frac{1}{2(p-1)} \sum_{\chi^2 \neq \chi_0} \tau^2(\overline{\chi}) \sum_{a=1}^{p-1} \sum_{b=1}^{p-1} \chi(a)\chi\chi_2(b) \, (a-b)^{2k}$$

$$= \Omega_1 + \Omega_2 + \Omega_3. \tag{9}$$

Noting that $J(p) = p-2$, every non-principal character χ modulo p is a primitive character mod p, and

$$\sum_{a=1}^{p-1} \chi(a) = 0, \quad \text{if } \chi \neq \chi_0, \qquad \sum_{a=1}^{p-1} a\chi(a) = 0, \quad \text{if } \chi(-1) = 1,$$

then by Lemma 2 and Lemma 4 we have

$$\begin{aligned}\Omega_1 &= \frac{1}{4(p-1)} \sum_{\substack{\chi\neq\chi_0\\ \chi(-1)=1}} \tau^2(\overline{\chi}) \sum_{n=2}^{2k-2} C_{2k}^n (-1)^n \left[\sum_{a=1}^{p-1} a^n\chi(a)\right]\left[\sum_{b=1}^{p-1} b^{2k-n}\chi(b)\right] \\ &\quad + \frac{1}{4(p-1)} \sum_{\chi(-1)=-1} \tau^2(\overline{\chi}) \sum_{n=1}^{2k-1} C_{2k}^n (-1)^n \left[\sum_{a=1}^{p-1} a^n\chi(a)\right]\left[\sum_{b=1}^{p-1} b^{2k-n}\chi(b)\right] \\ &= \frac{p^{2k+2}}{(p-1)} \sum_{\substack{\chi\neq\chi_0\\ \chi(-1)=1}} \sum_{n=2}^{2k-2} C_{2k}^n \sum_{m=1}^{[\frac{n}{2}]} \sum_{l=1}^{[\frac{2k-n}{2}]} \\ &\qquad \times \frac{C_n^{2m-1} C_{2k-n}^{2l-1} (2m-1)!(2l-1)! L(2m,\overline{\chi}) L(2l,\overline{\chi})}{(-1)^{n+m+l}(2\pi)^{2m+2l}} \\ &\quad - \frac{p^{2k+2}}{(p-1)} \sum_{\chi(-1)=-1} \sum_{n=1}^{2k-1} C_{2k}^n \sum_{m=0}^{[\frac{n-1}{2}]} \sum_{l=0}^{[\frac{2k-n-1}{2}]} \\ &\qquad \times \frac{C_n^{2m} C_{2k-n}^{2l} (2m)!(2l)! L(2m+1,\overline{\chi}) L(2l+1,\overline{\chi})}{(-1)^{n+m+l}(2\pi)^{2m+2l+2}} \\ &= 2p^{2k+2} \sum_{n=2}^{2k-2} C_{2k}^n \sum_{m=1}^{[\frac{n}{2}]} \sum_{l=1}^{[\frac{2k-n}{2}]} \frac{C_n^{2m-1} C_{2k-n}^{2l-1} (2m-1)!(2l-1)!}{(-1)^{n+m+l}(2\pi)^{2m+2l}} \\ &\quad - 2p^{2k+2} \sum_{n=1}^{2k-1} C_{2k}^n \sum_{m=0}^{[\frac{n-1}{2}]} \sum_{l=0}^{[\frac{2k-n-1}{2}]} \frac{C_n^{2m} C_{2k-n}^{2l} (2m)!(2l)!}{(-1)^{n+m+l}(2\pi)^{2m+2l+2}} \\ &\quad + O(p^{2k+1+\epsilon}). \end{aligned} \tag{10}$$

Similarly, from Lemma 2 and Lemma 6 we also have

$$\Omega_2 = \frac{1}{4(p-1)} \sum_{\substack{\chi\neq\chi_0\\ \chi(-1)=1}} \tau^2(\overline{\chi}\chi_2) \sum_{n=2}^{2k-2} C_{2k}^n (-1)^n \left[\sum_{a=1}^{p-1} a^n\chi(a)\right]\left[\sum_{b=1}^{p-1} b^{2k-n}\chi(b)\right]$$

$$+\frac{1}{4(p-1)}\sum_{\chi(-1)=-1}\tau^2(\overline{\chi}\chi_2)\sum_{n=1}^{2k-1}C_{2k}^n(-1)^n\left[\sum_{a=1}^{p-1}a^n\chi(a)\right]\left[\sum_{b=1}^{p-1}b^{2k-n}\chi(b)\right]$$

$$=\frac{p^{2k}}{(p-1)}\sum_{n=2}^{2k-2}C_{2k}^n\sum_{m=1}^{[\frac{n}{2}]}\sum_{l=1}^{[\frac{2k-n}{2}]}\frac{C_n^{2m-1}C_{2k-n}^{2l-1}(2m-1)!(2l-1)!}{(-1)^{n+m+l}(2\pi)^{2m+2l}}$$

$$\times\sum_{\substack{\chi\neq\chi_0\\ \chi(-1)=1}}\tau^2(\overline{\chi}\chi_2)\tau^2(\chi)L(2m,\overline{\chi})L(2l,\overline{\chi})$$

$$-\frac{p^{2k}}{(p-1)}\sum_{n=1}^{2k-1}C_{2k}^n\sum_{m=0}^{[\frac{n-1}{2}]}\sum_{l=0}^{[\frac{2k-n-1}{2}]}\frac{C_n^{2m}C_{2k-n}^{2l}(2m)!(2l)!}{(-1)^{n+m+l}(2\pi)^{2m+2l+2}}$$

$$\times\sum_{\chi(-1)=-1}\tau^2(\overline{\chi}\chi_2)\tau^2(\chi)L(2m+1,\overline{\chi})L(2l+1,\overline{\chi})$$

$$\ll p^{2k+\frac{3}{2}+\epsilon}. \tag{11}$$

Now we calculate Ω_3. If $p\equiv 1 \bmod 4$, then $\chi_2(-1)=1$. So from Lemma 2 and Lemma 7 we have

$$\Omega_3=\frac{1}{2(p-1)}\sum_{\substack{\chi^2\neq\chi_0\\ \chi(-1)=1}}\tau^2(\overline{\chi})\sum_{n=2}^{2k-2}C_{2k}^n(-1)^n\left[\sum_{a=1}^{p-1}a^n\chi(a)\right]\left[\sum_{b=1}^{p-1}b^{2k-n}\chi\chi_2(b)\right]$$

$$+\frac{1}{2(p-1)}\sum_{\chi(-1)=-1}\tau^2(\overline{\chi})\sum_{n=1}^{2k-1}C_{2k}^n(-1)^n\left[\sum_{a=1}^{p-1}a^n\chi(a)\right]\left[\sum_{b=1}^{p-1}b^{2k-n}\chi\chi_2(b)\right]$$

$$=\frac{2p^{2k+1}}{(p-1)}\sum_{n=2}^{2k-2}C_{2k}^n\sum_{m=1}^{[\frac{n}{2}]}\sum_{l=1}^{[\frac{2k-n}{2}]}\frac{C_n^{2m-1}C_{2k-n}^{2l-1}(2m-1)!(2l-1)!}{(-1)^{n+m+l}(2\pi)^{2m+2l}}$$

$$\times\sum_{\substack{\chi^2\neq\chi_0\\ \chi(-1)=1}}\tau(\overline{\chi})\tau(\chi\chi_2)L(2m,\overline{\chi})L(2l,\overline{\chi}\chi_2)$$

$$+\frac{2p^{2k+1}}{(p-1)}\sum_{n=1}^{2k-1}C_{2k}^n\sum_{m=0}^{[\frac{n-1}{2}]}\sum_{l=0}^{[\frac{2k-n-1}{2}]}\frac{C_n^{2m}C_{2k-n}^{2l}(2m)!(2l)!}{(-1)^{n+m+l}(2\pi)^{2m+2l+2}}$$

$$\times \sum_{\chi(-1)=-1} \tau(\overline{\chi})\tau(\chi\chi_2)L(2m+1,\overline{\chi})L(2l+1,\overline{\chi}\chi_2)$$

$$\ll p^{2k+\frac{3}{2}+\epsilon}. \tag{12}$$

On the other hand, if $p \equiv 3 \bmod 4$, then $\chi_2(-1) = -1$. By Lemma 2 and Lemma 7 we also have

$$\begin{aligned}\Omega_3 &= \frac{1}{2(p-1)} \sum_{\substack{\chi\neq\chi_0\\ \chi(-1)=1}} \tau^2(\overline{\chi}) \sum_{n=2}^{2k-2} C_{2k}^n (-1)^n \left[\sum_{a=1}^{p-1} a^n \chi(a)\right]\left[\sum_{b=1}^{p-1} b^{2k-n}\chi\chi_2(b)\right] \\ &\quad + \frac{1}{2(p-1)} \sum_{\substack{\chi\neq\chi_2\\ \chi(-1)=-1}} \tau^2(\overline{\chi}) \sum_{n=1}^{2k-1} C_{2k}^n (-1)^n \left[\sum_{a=1}^{p-1} a^n \chi(a)\right]\left[\sum_{b=1}^{p-1} b^{2k-n}\chi\chi_2(b)\right] \\ &= \frac{2p^{2k+1}}{(p-1)} \sum_{n=2}^{2k-2} C_{2k}^n \sum_{m=1}^{[\frac{n}{2}]} \sum_{l=0}^{[\frac{2k-n-1}{2}]} \frac{C_n^{2m-1} C_{2k-n}^{2l}(2m-1)!(2l)!}{(-1)^{n+m+l}(2\pi)^{2m+2l+1} i} \\ &\quad \times \sum_{\substack{\chi\neq\chi_0\\ \chi(-1)=1}} \tau(\overline{\chi})\tau(\chi\chi_2)L(2m,\overline{\chi})L(2l+1,\overline{\chi}\chi_2) \\ &\quad - \frac{2p^{2k+1}}{(p-1)} \sum_{n=1}^{2k-1} C_{2k}^n \sum_{m=0}^{[\frac{n-1}{2}]} \sum_{l=1}^{[\frac{2k-n}{2}]} \frac{C_n^{2m} C_{2k-n}^{2l-1}(2m)!(2l-1)!}{(-1)^{n+m+l}(2\pi)^{2m+2l+1} i} \\ &\quad \times \sum_{\substack{\chi\neq\chi_2\\ \chi(-1)=-1}} \tau(\overline{\chi})\tau(\chi\chi_2)L(2m+1,\overline{\chi})L(2l,\overline{\chi}\chi_2) \ll p^{2k+\frac{3}{2}+\epsilon}. \end{aligned} \tag{13}$$

Now combining formulae (9)–(13), we immediately get Theorem 1.

Acknowledgments

The authors express their gratitude to the referee for his very helpful comments in improving this paper.

References

[1] T. Masao, On certain character sums, *Acta Arithmetica* **LV** (1990), 229–232.

[2] A. Weil, Sur les courbes algébriques et les variéstés qui sèn déduisent, *Actualités Sci. Ind.* **1041** (1948).

[3] Wenpeng Zhang, On the difference between an integer and its inverse modulo n, *Journal of Number Theory* **52** (1995), 1–6.

[4] Wenpeng Zhang, On a Cochrane sum and its hybrid mean value formula, *Journal of Mathematical Analysis and Applications* **267** (2002), 89–96.

[5] Wenpeng Zhang, On the difference between an integer and its inverse modulo n (II), *Science in China (Series A)* **46** (2003), 229–238.

[6] Wenpeng Zhang and Huaning Liu, A Note on the Cochrane Sum and its Hybrid Mean Value Formula, *Journal of Mathematical Analysis and Applications* **288** (2003), 646–659.

LIU HUANING
DEPARTMENT OF MATHEMATICS
NORTHWEST UNIVERSITY
XI'AN, SHAANXI
P.R. CHINA

E-mail: hnliu@nwu.edu.cn

ZHANG WENPENG
DEPARTMENT OF MATHEMATICS
NORTHWEST UNIVERSITY
XI'AN, SHAANXI
P.R. CHINA

E-mail: wpzhang@nwu.edu.cn

(Received March 3, 2005; revised May 9, 2005)