

A note on the diagonal mapping in spaces of analytic functions in the unit polydisc

By MIROLJUB JEVTIĆ (Belgrade), MIROSLAV PAVLOVIĆ (Belgrade)
and ROMI F. SHAMOYAN (Bryansk)

Abstract. We define two spaces $K^{p,q,\alpha,\beta}$ and $M^{p,\alpha}$ of analytic functions in the unit polydisc U^n of C^n , closely related to the mixed norm and the Bergman spaces on U^n , and for any holomorphic function F in $K^{p,q,\alpha,\beta}$ or in $M^{p,\alpha}$ we consider its restriction to the diagonal, i.e., the function in the unit disc U of C defined by $DF(z) = F(z, \dots, z)$, and prove that the diagonal mapping D maps $K^{p,q,\alpha,\beta}$ onto the mixed-norm space $H^{p,q,\beta+\frac{q}{p}(|\alpha|+2n-1)}(U)$ and the space $M^{p,\alpha}$ onto the Bergman space $A^{p,|\alpha|+2n-1}(U)$.

Introduction

Let U^n be the unit polydisc in C^n and T^n be its Shilov boundary (see [R]) ($U^1 = U$ and $T^1 = T$). Denote by dm_n the normalized volume measure in U^n , and by $d\sigma_n$ the normalized surface measure on T^n .

For any Lebesgue measurable function f in U^n , we define

$$M_p(r, f) = \left(\int_{T^n} |f(r\xi)|^p d\sigma_n(\xi) \right)^{1/p},$$

where $0 < p < \infty$ and $r\xi = (r_1\xi_1, \dots, r_n\xi_n)$.

If $0 < p < \infty$, $0 < q < \infty$, and $\alpha = (\alpha_1, \dots, \alpha_n)$, $\alpha_j > -1$, $j = 1, \dots, n$, let

$$\|f\|_{p,q,\alpha}^q = \int_{I^n} \left(\prod_{j=1}^n (1 - r_j^2)^{\alpha_j} M_p(r, f)^q \right) dr,$$

Mathematics Subject Classification: 32A35, 32A37.

Key words and phrases: Bergman spaces, mixed norm spaces on the polydisc, diagonal mapping. The research of the first two authors was supported by a grant from MNS ON144010, Serbia.

where $I^n = [0, 1]^n$ and $dr = dr_1 \dots dr_n$. The mixed norm space $H^{p,q,\alpha} = H^{p,q,\alpha}(U^n)$ is then defined to be the space of functions f holomorphic in U^n , ($f \in H(U^n)$) such that $\|f\|_{p,q,\alpha} < \infty$.

First we give a new characterization of the mixed norm spaces $H^{p,q,\alpha}(U)$.

Theorem 1. *Let $0 < p, q < \infty$ and $-1 < \beta, \gamma < \infty$. A function $f \in H(U)$ belongs to $H^{p,q,\beta+q/p(\gamma+1)}(U)$ if and only if*

$$\int_0^1 \left(\int_{|z|<r} |f(z)|^p (1-|z|)^\gamma dm_1(z) \right)^{q/p} (1-r)^\beta dr < \infty.$$

Motivated by this characterization we define the spaces $K^{p,q,\alpha,\beta}(U^n)$ as follows.

The space $K^{p,q,\alpha,\beta} = K^{p,q,\alpha,\beta}(U^n)$ consists of all $f \in H(U^n)$ such that

$$\|f\|_{p,q,\alpha,\beta}^q = \int_0^1 \left(\int_{|z_1|<r} \dots \int_{|z_n|<r} |f(z)|^p \prod_{j=1}^n (1-|z_j|^2)^{\alpha_j} dm_n(z) \right)^{q/p} (1-r)^\beta dr < \infty,$$

where, $0 < p, q < \infty$, $-1 < \beta < \infty$ and $\alpha = (\alpha_1, \dots, \alpha_n)$, $\alpha_j > -1$, $j = 1, \dots, n$.

We note that the spaces $K^{p,q,\alpha,\beta}(U)$ were studied by [AJ] and [J].

To each $F \in H(U^n)$, we associate a function DF , defined on the unit disc U of C , by

$$DF(z) = F(z, \dots, z), \quad z \in U.$$

The problem of description of diagonal of subspaces of $H(U^n)$ was studied by many authors (see [DS], [MR], [HO], [RS], [Sh1], [Sh2], [S].)

Now we are ready to state the main result of this paper.

Theorem 2. *Let $0 < p, q < \infty$, $-1 < \beta < \infty$, $\alpha = (\alpha_1, \dots, \alpha_n)$, $\alpha_j > -1$, $j = 1, \dots, n$. Then*

$$DK^{p,q,\alpha,\beta}(U^n) = H^{p,q,\beta+q/p(|\alpha|+2n-1)}(U),$$

where $|\alpha| = \alpha_1 + \dots + \alpha_n$.

Note that in [RS] it was shown that

$$DH^{p,q,\alpha}(U^n) = H^{p,q,|\alpha|+(q/p+1)(n-1)}(U).$$

A special case of mixed norm spaces $H^{p,q,\alpha}(U^n)$ are the Bergman spaces $A^{p,\alpha}(U^n)$, $0 < p < \infty$ and $\alpha = (\alpha_1, \dots, \alpha_n)$, $\alpha_j > -1$, $j = 1, \dots, n$,

$$A^{p,\alpha} = A^{p,\alpha}(U^n) = H^{p,p,\alpha}(U^n).$$

J. SHAPIRO [S] and F. SHAMOYAN [Sh2] proved that

$$DA^{p,\alpha}(U^n) = A^{p,|\alpha|+2n-2}(U).$$

For example, if $\alpha' = (\alpha_1, \dots, \alpha_{j-1}, \alpha_j + 1, \alpha_{j+1}, \dots, \alpha_n)$, then

$$DA^{p,\alpha'}(U^n) = A^{p,|\alpha|+2n-1}(U).$$

In this paper we define the space $M^{p,\alpha}(U^n)$, $0 < p < \infty$, $\alpha = (\alpha_1, \dots, \alpha_n)$, $\alpha_j > -1$, $j = 1, \dots, n$, that contains $A^{p,\alpha'}(U^n)$, for $\alpha' = (\alpha_1, \dots, \alpha_{j-1}, \alpha_j + 1, \alpha_{j+1}, \dots, \alpha_n)$ and show that $DM^{p,\alpha}(U^n) = A^{p,|\alpha|+2n-1}(U)$.

Let $0 < p < \infty$ and $\alpha = (\alpha_1, \dots, \alpha_n)$, $\alpha_j > -1$, $j = 1, \dots, n$. Define $M^{p,\alpha} = M^{p,\alpha}(U^n)$ as the space of all $f \in H(U^n)$ such that

$$\|f\|_{p,\alpha}^p = \int_T \int_{\Gamma_t(\xi)} \dots \int_{\Gamma_t(\xi)} |f(z)|^p \prod_{k=1}^n (1 - |z_k|^2)^{\alpha_k} dm_n(z) d\sigma_1(\xi) < \infty,$$

where $\Gamma_t(\xi) = \{z \in U : |z - \xi| < t(1 - |z|)\}$, $t > 1$, $\xi \in T$ is the Stolz angle with vertex ξ .

Theorem 3. *Let $0 < p < \infty$, $\alpha = (\alpha_1, \dots, \alpha_n)$, $\alpha_j > -1$, $j = 1, \dots, n$. Then*

$$DM^{p,\alpha}(U^n) = A^{p,|\alpha|+2n-1}(U),$$

where $|\alpha| = \alpha_1 + \dots + \alpha_n$.

1. Preliminaries

In this Section we gather several partially well-known lemmas that will be used in the proofs of our results.

Lemma 1. *Let $b > a > 0$, $c > 0$ and let $g : [0, 1] \rightarrow [0, \infty)$ be measurable. Assume either $0 < k < 1$ and g is increasing or $1 \leq k < \infty$. Then*

$$\int_0^1 (1-r)^{ka-1} \left(\int_0^1 \frac{g(\rho)(1-\rho)^{c-1} d\rho}{(1-r\rho)^b} \right)^k dr \leq C \int_0^1 (1-r)^{k(a+c-b)-1} g^k(r) dr.$$

See [RS, Proposition 3.1]. See also [AJ] and [M].

Lemma 2. *Let $0 < p < \infty$ and $\beta > 0$. Let (a_n) be a sequence of nonnegative real numbers such that $\sum_{n=1}^{\infty} 2^{-n\beta} a_n^p < \infty$. Then there is a constant $C > 0$, depending only on p and β , such that*

$$\sum_{n=1}^{\infty} 2^{-n\beta} a_n^p \leq C \left(a_0^p + \sum_{n=1}^{\infty} 2^{-n\beta} |a_n - a_{n-1}|^p \right).$$

See [MP, Lemma HL].

Lemma 3. *Let $0 < p < 1$, $\beta > -1$ and $\beta p + 2p > 1$. If $f \in H^{p,p,\beta p+2p-2}(U)$, then $f \in H^{1,1,\beta}(U)$. Moreover, there is a constant $C > 0$, depending only on p and β , such that*

$$\|f\|_{1,1,\beta} \leq C \|f\|_{p,p,\beta p+2p-2}.$$

PROOF. By using the standard estimate

$$M_1(r^2, f) \leq C(1-r)^{1-(1/p)} M_p(r, f),$$

(see [D]), we find that

$$\int_U |f(z)|(1-|z|)^\beta dm_1(z) \leq C \int_0^1 M_p(r, f)(1-r)^{\beta-(1/p)+1} dr.$$

Let $r_n = 1 - 2^{-n}$, $n = 0, 1, 2, \dots$. Then we have

$$\begin{aligned} \|f\|_{1,1,\beta}^p &\leq C \left(\sum_{n=1}^{\infty} \int_{r_{n-1}}^{r_n} M_p(r, f)(1-r)^{\beta-(1/p)+1} dr \right)^p \\ &\leq C \sum_{n=1}^{\infty} 2^{-np(\beta-(1/p)+2)} M_p^p(r_n, f) \\ &\leq C \sum_{n=0}^{\infty} \int_{r_n}^{r_{n+1}} M_p^p(r, f)(1-r)^{p(\beta-(1/p)+2)-1} dr \\ &\leq C \int_U |f(z)|^p (1-|z|)^{\beta p+2p-2} dm_1(z). \end{aligned} \quad \square$$

An integration in polar coordinates shows that the following is true.

Lemma 4. *Let $\beta > -1$ and $\gamma - \beta > 2$. Then there is a constant C , depending only on β and γ , such that*

$$\int_{|z|<r} \frac{(1-|z|)^\beta}{|1-\bar{w}z|^\gamma} dm_1(z) \leq \frac{C}{(1-r|w|)^{\gamma-\beta-2}}, \quad |w| < 1, \quad 0 < r \leq 1.$$

Lemma 5. *Let $\beta > -1$, $\gamma > \beta + 2$ and $t > 1$. Then*

$$\int_{\Gamma_t(\xi)} \frac{(1 - |z|)^\beta}{|1 - \bar{z}w|^\gamma} dm_1(z) \leq \frac{C}{|1 - \xi\bar{w}|^{\gamma-\beta-2}}, \quad \xi \in T, |w| < 1,$$

where C is a constant depending only on β , γ and t .

Lemma 6. *Let $\beta > 1$. Then*

$$\int_T \frac{d\sigma_1(\xi)}{|1 - \xi z|^\beta} \leq \frac{C}{(1 - |z|)^{\beta-1}}, \quad |z| < 1.$$

See [D].

2. On characterization of the mixed norm spaces in the unit polydisc U

In this section we prove Theorem 1.

PROOF OF THEOREM 1. First, assume that $\|f\|_{p,q,\beta+q/p(\gamma+1)} < \infty$. For $0 < t < 1$ define $f_t(z) = f(tz)$, $z \in U$. Let $r_n = 1 - 2^{-n}$, $n = 0, 1, \dots$. Then

$$\begin{aligned} & \int_0^1 (1-r)^\beta \left(\int_{|z|<r} |f_t(z)|^p (1-|z|)^\gamma dm_1(z) \right)^{q/p} dr \\ &= \sum_{n=1}^{\infty} \int_{r_{n-1}}^{r_n} (1-r)^\beta \left(\int_{|z|<r} |f_t(z)|^p (1-|z|)^\gamma dm_1(z) \right)^{q/p} dr \leq C \sum_{n=1}^{\infty} 2^{-n(\beta+1)} A_n^{q/p}, \end{aligned}$$

where

$$A_n = \int_{|z|<r_n} |f_t(z)|^p (1-|z|)^\gamma dm_1(z).$$

Now by using Lemma 2 we find that

$$\begin{aligned} & \int_0^1 (1-r)^\beta \left(\int_{|z|<r} |f_t(z)|^p (1-|z|)^\gamma dm_1(z) \right)^{q/p} dr \\ & \leq C \sum_{n=1}^{\infty} 2^{-n(\beta+1)} \left(\int_{r_{n-1} \leq |z| < r_n} |f_t(z)|^p (1-|z|)^\gamma dm_1(z) \right)^{q/p} \\ & \leq C \sum_{n=1}^{\infty} 2^{-n(\beta+1)} M_p(r_n, f_t)^q 2^{-n(\gamma+1)q/p} \end{aligned}$$

$$\begin{aligned}
&\leq C \sum_{n=1}^{\infty} \int_{r_n}^{r_{n+1}} M_p(r, f_t)^q (1-r)^{\beta+q/p(\gamma+1)} dr \\
&\leq C \int_0^1 (1-r)^{\beta+q/p(\gamma+1)} M_p(r, f_t)^q dr.
\end{aligned}$$

Letting $t \rightarrow 1$, we get

$$\int_0^1 (1-r)^{\beta} \left(\int_{|z|<r} |f(z)|^p (1-|z|)^{\gamma} dm_1(z) \right)^{q/p} dr \leq C \|f\|_{p,q,\beta+q/p(\gamma+1)}^q.$$

Conversely,

$$\begin{aligned}
\|f\|_{p,q,\beta+q/p(\gamma+1)}^q &= \int_0^1 (1-r)^{\beta+q/p(\gamma+1)} M_p(r, f)^q dr \\
&= \sum_{n=1}^{\infty} \int_{r_{n-1}}^{r_n} (1-r)^{\beta+q/p(\gamma+1)} M_p(r, f)^q dr \\
&\leq C \sum_{n=1}^{\infty} 2^{-n(\beta+q/p(\gamma+1)+1)} M_p(r_n, f)^q \\
&\leq C \sum_{n=1}^{\infty} \left(\int_{r_n < |z| < r_{n+1}} |f(z)|^p (1-|z|)^{\gamma} dm_1(z) \right)^{q/p} 2^{-n(\beta+1)} \\
&\leq C \sum_{n=1}^{\infty} \left(\int_{r_n < |z| < r_{n+1}} |f(z)|^p (1-|z|)^{\gamma} dm_1(z) \right)^{q/p} \int_{r_{n+1}}^{r_{n+2}} (1-r)^{\beta} dr \\
&\leq C \sum_{n=1}^{\infty} \int_{r_{n+1}}^{r_{n+2}} (1-r)^{\beta} \left(\int_{|z|<r} |f(z)|^p (1-|z|)^{\gamma} dm_1(z) \right)^{q/p} dr \\
&\leq C \int_0^1 (1-r)^{\beta} \left(\int_{|z|<r} |f(z)|^p (1-|z|)^{\gamma} dm_1(z) \right)^{q/p} dr.
\end{aligned}$$

3. On the diagonal of $K^{p,q,\alpha,\beta}(U^n)$

In this section we prove Theorem 2.

PROOF OF THEOREM 2. For $k = 0, 1, \dots$, let

$$U_k = \{U_{k,j} : j = 0, 1, \dots, 2^{k+1} - 1\},$$

where

$$U_{k,j} = \left\{ z \in U : 1 - \frac{1}{2^k} \leq |z| < 1 - \frac{1}{2^{k+1}}, \frac{2\pi j}{2^{k+1}} \leq \arg z < \frac{2\pi(j+1)}{2^{k+1}} \right\},$$

and let

$$V_k = \{V_{k,j} : j = 0, 1, \dots, 2^{k+1} - 1\},$$

where

$$V_{k,j} = \left\{ z \in U : 1 - \frac{1}{2^{k-1}} \leq |z| < 1 - \frac{1}{2^{k+2}}, \frac{2\pi(j-1/2)}{2^{k+1}} \leq \arg z < \frac{2\pi(j+3/2)}{2^{k+1}} \right\}.$$

To prove Theorem 2, we will use the dyadic decomposition of the polydisc (see [DjS]):

$$U_{k,l_1,\dots,l_n} = U_{k,l_1} \times \dots \times U_{k,l_n},$$

where $k = 0, 1, \dots$ and $U_{k,l_j} \in U_k$, $j = 1, \dots, n$.

Now, let $r_m = 1 - \frac{1}{2^m}$, $m = 0, 1, \dots$. Then by using Theorem 1 we find that

$$\begin{aligned} & \|Df\|_{p,q,\beta+q/p(|\alpha|+2n-1)}^q \\ & \leq C \int_0^1 (1-r)^\beta \left(\int_{|z|<r} |Df(z)|^p (1-|z|^2)^{|\alpha|+2n-2} dm_1(z) \right)^{q/p} dr \\ & = C \sum_{m=0}^{\infty} \int_{r_m}^{r_{m+1}} (1-r)^\beta \left(\int_{|z|<r} |Df(z)|^p (1-|z|^2)^{|\alpha|+2n-2} dm_1(z) \right)^{q/p} dr \\ & \leq C \sum_{m=0}^{\infty} 2^{-m(\beta+1)} \left(\int_{|z|<r_{m+1}} |Df(z)|^p (1-|z|^2)^{|\alpha|+2n-2} dm_1(z) \right)^{q/p} \\ & \leq C \sum_{m=0}^{\infty} 2^{-m(\beta+1)} \left(\sum_{k \leq m} \sum_{l_1, \dots, l_n} \left(\max_{z \in U_{k,l_1, \dots, l_n}} |f(z)|^p \right) 2^{-k(|\alpha|+2n-2)} 2^{-2k} \right)^{q/p}. \end{aligned} \quad (1)$$

From the n -subharmonicity of $|f(z)|^p$, $0 < p < \infty$, it follows that

$$\max_{z \in U_{k,l_1, \dots, l_n}} |f(z)|^p \leq C 2^{2kn} \int_{V_{k,l_1, \dots, l_n}} |f(w_1, \dots, w_n)|^p dm_n(w). \quad (2)$$

See [DjS].

The family of enlarged sets $V_{k,j}$, $k = 0, 1, \dots$, $j = 0, 1, \dots, 2^{k+1} - 1$, is a finite covering of U^n . Thus,

$$\begin{aligned} & \sum_{k \leq m} \sum_{l_1, \dots, l_n} \int_{V_{k,l_1, \dots, l_n}} |f(w_1, \dots, w_n)|^p \prod_{j=1}^n (1-|w_j|^2)^{\alpha_j} dm_n(w) \\ & \leq C \int_{|w_1| < r_{m+2}} \dots \int_{|w_n| < r_{m+2}} |f(w_1, \dots, w_n)|^p \prod_{j=1}^n (1-|w_j|^2)^{\alpha_j} dm_n(w). \end{aligned} \quad (3)$$

Combining (1), (2) and (3) we find that

$$\begin{aligned}
& \|Df\|_{\beta+q/p(\alpha+2n-1)}^q \\
& \leq C \sum_{m=0}^{\infty} 2^{-m(\beta+1)} \int_{|w_1| < r_{m+2}} \cdots \int_{|w_n| < r_{m+2}} |f(w_1, \dots, w_n)|^p \\
& \quad \times \prod_{j=1}^n (1 - |w_j|^2)^{\alpha_j} dm_n(w) \\
& \leq C \sum_{m=0}^{\infty} \int_{r_{m+2}}^{r_{m+3}} (1-r)^{\beta} \left(\int_{|w_1| < r} \cdots \int_{|w_n| < r} |f(w_1, \dots, w_n)|^p \right. \\
& \quad \times \left. \prod_{j=1}^n (1 - |w_j|^2)^{\alpha_j} dm_n(w) \right)^{q/p} dr \\
& \leq C \int_0^1 (1-r)^{\beta} \left(\int_{|w_1| < r} \cdots \int_{|w_n| < r} |f(w)|^p \prod_{j=1}^n (1 - |w_j|^2)^{\alpha_j} dm_n(w) \right)^{q/p} dr.
\end{aligned}$$

It remains to show that

$$D : K^{p,q,\alpha,\beta}(U^n) \rightarrow H^{p,q,\beta+q/p(\alpha+2n-1)}(U)$$

is onto.

Let $f \in H^{p,q,\beta+q/p(\alpha+2n-1)}(U)$. Define a function F on U^n by

$$F(z_1, \dots, z_n) = s \int_U \frac{f(w)(1 - |w|^2)^{s-1}}{\prod_{j=1}^n (1 - z_j \bar{w})^{(s+1)/n}} dm_1(w),$$

where s is a positive real number. Then F is holomorphic in U^n and it is easy to see that

$$DF(z) = F(z, \dots, z) = f(z), \quad z \in U.$$

Let $0 < p \leq 1$. We may assume that $p(s+1) - n \max_{1 \leq j \leq n} \alpha_j - 2n > (\beta+1)p/q$. Then using Lemma 3 and Lemma 4 we find that

$$\begin{aligned}
\|F\|_{p,q,\alpha,\beta}^q &= \int_0^1 \left(\int_{|z_1| < r} \cdots \int_{|z_n| < r} |F(z)|^p \prod_{j=1}^n (1 - |z_j|^2)^{\alpha_j} dm_n(z) \right)^{q/p} (1-r)^{\beta} dr \\
&\leq C \int_0^1 \left(\int_{|z_1| < r} \cdots \int_{|z_n| < r} \left(\int_U \frac{|f(w)|^p (1 - |w|)^{p(s-1)+2p-2} dm_1(w)}{\prod_{j=1}^n |1 - z_j \bar{w}|^{(s+1)p/n}} \right) \right. \\
&\quad \times \left. \prod_{j=1}^n (1 - |z_j|)^{\alpha_j} dm_n(z) \right)^{q/p} (1-r)^{\beta} dr
\end{aligned}$$

$$\leq C \int_0^1 \left(\int_0^1 M_p^p(\rho, f) \frac{(1-\rho)^{p(s-1)+2p-2}}{(1-r\rho)^{p(s+1)-|\alpha|-2n}} d\rho \right)^{q/p} (1-r)^\beta dr.$$

An application of Lemma 1 gives

$$\begin{aligned} \|F\|_{p,q,\alpha,\beta}^q &\leq C \int_0^1 (1-r)^{\beta+q/p(|\alpha|+2n-1)} M_p(r, f)^q dr \\ &= C \|f\|_{p,q,\beta+q/p(|\alpha|+2n-1)}^q. \end{aligned}$$

Thus, $F \in K^{p,q,\alpha,\beta}(U^n)$ and $DF = f$.

Assume now that $1 < p < \infty$ and $(s+1) > n \max_{1 \leq k \leq n} (\alpha_k + 2)$. Let γ_1 and γ_2 be positive real numbers such that $\gamma_1 + \gamma_2 = (s+1)/n$, and

$$\frac{s+1}{pn} - \frac{\min_{1 \leq k \leq n} \alpha_k + 1}{p} < \gamma_1 < \frac{s+1}{pn}.$$

Let q denote the conjugate of p , i.e. $1/p + 1/q = 1$. Then we have

$$\begin{aligned} \|F\|_{p,q,\alpha,\beta}^q &\leq C \int_0^1 \left(\int_{|z_1| < r} \cdots \int_{|z_n| < r} \left(\int_U \frac{|f(w)|(1-|w|^2)^{s-1} dm_1(w)}{\prod_{j=1}^n |1 - z_j \bar{w}|^{(s+1)/n}} \right)^p \right. \\ &\quad \times \left. \prod_{j=1}^n (1 - |z_j|^2)^{\alpha_j} dm_n(z) \right)^{q/p} (1-r)^\beta dr. \end{aligned} \quad (4)$$

By using Hölder's inequality we find that

$$\begin{aligned} \left(\int_U \frac{|f(w)|(1-|w|^2)^{s-1} dm_1(w)}{\prod_{j=1}^n |1 - z_j \bar{w}|^{(s+1)/n}} \right)^p &\leq \int_U \frac{|f(w)|^p (1-|w|^2)^{s-1} dm_1(w)}{\prod_{j=1}^n |1 - z_j \bar{w}|^{\gamma_1 p}} \\ &\quad \times \left(\int_U \frac{(1-|w|^2)^{s-1} dm_1(w)}{\prod_{j=1}^n |1 - z_j \bar{w}|^{\gamma_2 q}} \right)^{p/q}. \end{aligned} \quad (5)$$

Using the Hölder's inequality and Lemma 4 we see that

$$\begin{aligned} \left(\int_U \frac{(1-|w|^2)^{s-1} dm_1(w)}{\prod_{j=1}^n |1 - z_j \bar{w}|^{\gamma_2 q}} \right)^{p/q} &\leq \prod_{j=1}^n \left(\int_U \frac{(1-|w|^2)^{s-1} dm_1(w)}{|1 - z_j \bar{w}|^{n\gamma_2 q}} \right)^{p/(qn)} \\ &\leq \prod_{j=1}^n \frac{1}{(1 - |z_j|)^{(s+1)/n - p\gamma_1}}. \end{aligned} \quad (6)$$

Thus, from (4), (5) and (6) it follows that

$$\|F\|_{p,q,\alpha,\beta}^q \leq C \int_0^1 \left(\int_{|z_1| < r} \cdots \int_{|z_n| < r} \int_U \frac{|f(w)|^p (1-|w|^2)^{s-1} dm_1(w)}{\prod_{j=1}^n |1 - z_j \bar{w}|^{\gamma_1 p}} \right)$$

$$\begin{aligned}
& \times \prod_{j=1}^n (1 - |z_j|^2)^{\alpha_j - (s+1)/n + \gamma_1 p} dm_n(z) \Big)^{q/p} (1-r)^\beta dr \\
& \leq C \int_0^1 \left(\int_U |f(w)|^p (1 - |w|^2)^{s-1} \right. \\
& \quad \times \left. \left(\prod_{j=1}^n \int_{|z_j| < r} \frac{(1 - |z_j|)^{\alpha_j - (s+1)/n + \gamma_1 p} dm_1(z_j)}{|1 - z_j \bar{w}|^{\gamma_1 p}} \right) dm_1(w) \right)^{q/p} (1-r)^\beta dr.
\end{aligned}$$

Now by using Lemma 4 and Lemma 1 we get

$$\begin{aligned}
\|F\|_{p,q,\alpha,\beta}^q & \leq C \int_0^1 \left(\int_U \frac{|f(w)|^p (1 - |w|^2)^{s-1} dm_1(w)}{(1 - r|w|)^{s+1-|\alpha|-2n}} \right)^{q/p} (1-r)^\beta dr \\
& \leq C \int_0^1 M_p(r, f)^q (1-r)^{\beta+q/p(|\alpha|+2n-1)} dr \\
& = C \|f\|_{p,q,\beta+q/p(|\alpha|+2n-1)}^q.
\end{aligned}$$

This finishes the proof of Theorem 2.

4. On the diagonal of $M^{p,\alpha}(U^n)$

PROOF OF THEOREM 3. Let $f \in H^{p,\alpha}(U^n)$. To show that

$$\|f\|_{p,\alpha}^p \geq C \int_U |Df(z)|^p (1 - |z|^2)^{|\alpha|+2n-1} dm_1(z) \quad (7)$$

we will use the following inequality

$$\begin{aligned}
& \int_U |Df(z)|^p (1 - |z|^2)^{|\alpha|+2n-1} dm_1(z) \\
& = \sum_{k,j} \int_{U_{k,j}} |Df(z)|^p (1 - |z|^2)^{|\alpha|+2n-1} dm_1(z) \\
& \leq C \sum_{k,j} 2^{-k(|\alpha|+2n)} \int_{2\pi j/2^{k+1}}^{2\pi(j+1)/2^{k+1}} \sup_{1-2^{-k} \leq \rho < 1-2^{-(k+1)}} |f(\rho e^{i\theta}, \dots, \rho e^{i\theta})|^p d\theta \\
& \leq C \sum_k 2^{-k(|\alpha|+2n)} \int_0^{2\pi} \sup_{1-2^{-k} \leq \rho < 1-2^{-(k+1)}} |f(\rho e^{i\theta}, \dots, \rho e^{i\theta})|^p d\theta \\
& \leq C \sum_{k_1, \dots, k_n} \int_{1-2^{-(k_1+1)}}^{1-2^{-(k_1+2)}} (1 - \rho_1^2)^{\alpha_1+1} d\rho_1 \cdots \int_{1-2^{-(k_n+1)}}^{1-2^{-(k_n+2)}} (1 - \rho_n^2)^{\alpha_n+1} d\rho_n
\end{aligned}$$

$$\begin{aligned}
& \times \int_0^{2\pi} |f(\rho_1 e^{i\theta}, \dots, \rho_n e^{i\theta})|^p d\theta \\
& \leq C \int_T \left(\int_0^1 \cdots \int_0^1 |f(\rho_1 \xi, \dots, \rho_n \xi)|^p \prod_{j=1}^n (1 - \rho_j^2)^{\alpha_j+1} d\rho_1 \cdots d\rho_n \right) d\sigma_1(\xi). \quad (8)
\end{aligned}$$

If u is a function defined on U , and $0 < \alpha, p < \infty$, than as a special case of the following known asymptotic relation (see [C])

$$\int_U |g(z)|^p d\mu(z) \cong \int_T \left(\int_{\Gamma_t(\xi)} \frac{|g(z)|^p}{1 - |z|} d\mu(z) \right) d\sigma_1(\xi),$$

where $d\mu$ is a positive Borel measure on U , we have

$$\int_0^1 |u(re^{i\theta})|^p (1-r)^\alpha dr \leq C \int_{\Gamma_t(e^{i\theta})} |u(z)|^p (1-|z|^2)^{\alpha-1} dm_1(z).$$

Using this inequality, by each variable, we get the following

$$\begin{aligned}
& \int_0^1 \cdots \int_0^1 |f(\rho_1 e^{i\theta}, \dots, \rho_n e^{i\theta})|^p \prod_{j=1}^n (1 - \rho_j^2)^{\alpha_j+1} d\rho_1 \cdots d\rho_n \\
& \leq C \int_{\Gamma_t(e^{i\theta})} \cdots \int_{\Gamma_t(e^{i\theta})} |f(z)|^p \prod_{j=1}^n (1 - |z_j|^2)^{\alpha_j} dm_n(z).
\end{aligned}$$

Integrating both sides by T and using (8) we get (7). The argument given here is taken from [LS].

The inequality (7) shows that the diagonal mapping D maps $M^{p,\alpha}(U^n)$ into $A^{p,|\alpha|+2n-1}(U)$. That D is onto follows from the following two facts:

$$A^{p,\alpha'}(U^n) \subset M^{p,\alpha}(U^n) \quad \text{and} \quad DA^{p,\alpha'}(U^n) = A^{p,|\alpha|+2n-1}(U),$$

where $\alpha = (\alpha_1, \dots, \alpha_n)$ and $\alpha' = (\alpha_1, \dots, \alpha_{j-1}, \alpha_j + 1, \alpha_{j+1}, \dots, \alpha_n)$.

For a reader convenience, we show that an extension operator, the right inverse of the diagonal operator, is bounded from $A^{p,|\alpha|+2n-1}(U)$ into $M^{p,\alpha}$ and consequently $D : M^{p,\alpha}(U^n) \rightarrow A^{p,|\alpha|+2n-1}(U)$ is onto.

Let $f \in A^{p,|\alpha|+2n-1}(U)$. Define a function F on U^n by

$$F(z_1, \dots, z_n) = s \int_U \frac{f(w)(1 - |w|^2)^{s-1}}{\prod_{j=1}^n (1 - z_j \bar{w})^{(s+1)/n}} dm_1(w),$$

where s is a positive real number. Obviously, $DF = f$, by the known Bergman representation formulas.

First, assume that $0 < p \leq 1$. We may assume that $p(s+1) > n \max_{1 \leq j \leq n} \alpha_j + 2n + 1$. By using Lemma 3, Lemma 5 and Lemma 6 we obtain

$$\begin{aligned}
\|F\|_{p,\alpha}^p &\leq C \int_T \left(\int_{\Gamma_t(\xi)} \cdots \int_{\Gamma_t(\xi)} \int_U \frac{|f(w)|^p (1 - |w|^2)^{p(s-1)+2p-2} dm_1(w)}{\prod_{j=1}^n |1 - z_j \bar{w}|^{(s+1)p/n}} \right. \\
&\quad \times \left. \prod_{j=1}^n (1 - |z_j|^2)^{\alpha_j} dm_1(z_j) \right) d\sigma_1(\xi) \\
&\leq C \int_T \left(\int_U |f(w)|^p (1 - |w|^2)^{p(s-1)+2p-2} \right. \\
&\quad \times \left. \prod_{j=1}^n \int_{\Gamma_t(\xi)} \frac{(1 - |z_j|^2)^{\alpha_j} dm_1(z_j)}{|1 - z_j \bar{w}|^{(s+1)p/n}} dm_1(w) \right) d\sigma_1(\xi) \\
&\leq C \int_T \left(\int_U \frac{|f(w)|^p (1 - |w|^2)^{p(s-1)+2p-2}}{|1 - \xi \bar{w}|^{p(s+1)-|\alpha|-2n}} dm_1(w) \right) d\sigma_1(\xi) \\
&\leq C \int_U |f(w)|^p (1 - |w|^2)^{|\alpha|+2n-1} dm_1(w).
\end{aligned}$$

Case 1 $1 < p < \infty$. We may assume that $s > n \max_{1 \leq j \leq n} \alpha_j + 2n$.

Let q be the conjugate of p and let γ_1 and γ_2 be positive numbers such that $\gamma_1 + \gamma_2 = (s+1)/n$. and

$$\frac{s+1}{pn} - \frac{\alpha_j + 1}{p} < \gamma_1 < \frac{s+1}{pn}, \quad \text{for } j = 1, \dots, n.$$

By using Hölder's inequality we find that

$$\begin{aligned}
\|F\|_{p,\alpha}^p &\leq C \int_T \left(\int_{\Gamma_t(\xi)} \cdots \int_{\Gamma_t(\xi)} \left(\int_U \frac{|f(w)|^p (1 - |w|^2)^{s-1}}{\prod_{j=1}^n |1 - z_j \bar{w}|^{\gamma_1 p}} dm_1(w) \right. \right. \\
&\quad \times \left. \left. \left(\int_U \frac{(1 - |w|^2)^{s-1} dm_1(w)}{\prod_{j=1}^n |1 - z_j \bar{w}|^{\gamma_2 q}} \right)^{p/q} \right) \prod_{j=1}^n (1 - |z_j|^2)^{\alpha_j} dm_1(z_j) \right) d\sigma_1(\xi). \quad (9)
\end{aligned}$$

An application of Hölders inequality and Lemma 4 gives

$$\begin{aligned}
&\left(\int_U \frac{(1 - |w|^2)^{s-1} dm_1(w)}{\prod_{j=1}^n |1 - z_j \bar{w}|^{\gamma_2 q}} \right)^{p/q} \\
&\leq \prod_{j=1}^n \left(\int_U \frac{(1 - |w|^2)^{s-1} dm_1(w)}{|1 - z_j \bar{w}|^{n \gamma_2 q}} \right)^{p/(qn)} \leq C \prod_{j=1}^n (1 - |z_j|)^{\gamma_1 p - (s+1)/n}. \quad (10)
\end{aligned}$$

By using (9), (10) and Lemma 6 we see that

$$\begin{aligned}
\|F\|_{p,\alpha}^p &\leq C \int_T \left(\int_{\Gamma_t(\xi)} \cdots \int_{\Gamma_t(\xi)} \int_U \frac{|f(w)|^p (1 - |w|^2)^{s-1} dm_1(w)}{\prod_{j=1}^n |1 - z_j \bar{w}|^{\gamma_1 p}} \right. \\
&\quad \left. \times \prod_{j=1}^n (1 - |z_j|^2)^{\alpha_j + \gamma_1 p - (s+1)/n} dm_1(z_j) \right) d\sigma_1(\xi) \\
&= C \int_T \left(\int_U |f(w)|^p (1 - |w|^2)^{s-1} dm_1(w) \right. \\
&\quad \left. \times \prod_{j=1}^n \int_{\Gamma_t(\xi)} \frac{(1 - |z_j|^2)^{\alpha_j + \gamma_1 p - (s+1)/n}}{|1 - z_j \bar{w}|^{\gamma_1 p}} dm_1(z_j) \right) d\sigma_1(\xi) \\
&\leq C \int_T \left(\int_U \frac{|f(w)|^p (1 - |w|^2)^{s-1}}{|1 - \xi \bar{w}|^{s+1-|\alpha|-2n}} dm_1(w) \right) d\sigma_1(\xi) \\
&= C \int_U |f(w)|^p (1 - |w|^2)^{s-1} \left(\int_T \frac{d\sigma_1(\xi)}{|1 - \xi \bar{w}|^{s+1-|\alpha|-2n}} \right) dm_1(w) \\
&\leq C \int_U |f(w)|^p (1 - |w|^2)^{|\alpha|+2n-1} dm_1(w).
\end{aligned}$$

References

- [AJ] P. AHERN, M. JEVIĆ, Duality and multipliers for mixed norm spaces, *Michigan Math. J.* **30** (1983), 53–64.
- [C] W. COHN, Generalized area operators on Hardy spaces, *J. Math. Anal. Appl.* **216** (1997), 112–121.
- [DjS] A. E. DJRBASHIAN and F. A. SHAMOYAN, Topics in the Theory of A_α^p Spaces, *Leipzig, Teubner*, 1988.
- [D] P. L. DUREN, Theory of H^p Spaces, *Academic Press, New York*, 1970, reprinted by Dover, Mineola, NY 2000.
- [DS] P. L. DUREN and A. L. SHIELDS, Restrictions of H^p functions to the diagonal of the polydisc, *Duke Math. J.* **42** (1975), 751–753.
- [HO] C. HOROWITZ and D. OBERLIN, Restriction of H^p functions to the diagonal of U^n , *Indiana Univ. Math. J.* **24** (1975), 767–772.
- [J] M. JEVIĆ, Sur la derivee de la fonction atomique, *C. R. Acad. Sci. Paris Sér. I Math.* **292** (1981), 201–203 (in *French. English summary*).
- [LS] S. LI and R. SHAMOYAN, On a differential operator connected with finite sum of functions analytic in the polydisk, (*preprint*).
- [MP] M. MATELJEVIĆ and M. PAVLOVIĆ, Multipliers of H^p and $BMOA$, *Pacific Journal of Mathematics* **146** (1990), 71–84.
- [M] V. MAZYA, Sobolev Spaces, *Moscow*, 1985.
- [MR] B. MOULIN and J. P. ROSAY, Sur la restriction des fonctions pluri-sous-harmoniques a la diagonale du polydisque, *Indiana Univ. Math. J.* **26** (1977), 869–873 (in *French*).

- [RS] G. REN and J. SHI, The diagonal mapping in mixed norm spaces, *Studia Mathematica* **163**(2) (2004), 103–117.
- [R] W. RUDIN, Function Theory in Polydiscs, *W. A. Benjamin, Inc., New York – Amsterdam*, 1969.
- [Sh1] F. A. SHAMOYAN, Diagonal mapping and problems of representation in anisotropic spaces of functions that are holomorphic in a polydisc, *Siberian Math. J.* **31** (1990), 350–365.
- [Sh2] F. A. SHAMOYAN, Applications of Dzhrbashyan integral representations to some problems of analysis, *Dokl. Akad. Nauk SSSR* **261**, no. 3 (1981), 557–561 (in *Russian*).
- [RSh] R. F. SHAMOYAN, The generalized Hardy transformation and Toeplitz operators in BMOA-type spaces, *Ukrain. Mat. Zh.* **53**, no. 9 (2001), 1260–1271 (in *Russian*).
- [S] J. SHAPIRO, Mackey topologies, reproducing kernels and diagonal maps on the Hardy and Bergman spaces, *Duke Mathematical J.* **43** (1976), 187–202.

MIROLJUB JEVTIĆ
MATEMATIČKI FAKULTET
PP. 550, 11000 BELGRADE
SERBIA

E-mail: jevtic@matf.bg.ac.yu

MIROSLAV PAVLOVIĆ
MATEMATIČKI FAKULTET
PP. 550, 11000 BELGRADE
SERBIA

E-mail: pavlovic@matf.bg.ac.yu

ROMI F. SHAMOYAN
BRYANSK STATE UNIVERSITY
BRYANSK, 241050
RUSSIA

E-mail: rsham@mail.ru

(Received February 21, 2008; revised September 21, 2008)