

On absolutely conformal mappings

By DAVID KALAJ (Podgorica) and MIODRAG MATELJEVIĆ (Belgrade)

Abstract. Let Ω be a domain in $\mathbb{R}^n$. It is proved that, if $u \in C^1(\Omega; \mathbb{R}^n)$ and there holds the formula $\|\nabla u(x)\|^n = n^{n/2} |\det \nabla u(x)|$ in Ω , then for $n \geq 3$ u is a restriction of a Möbius transformation, and for $n = 2$, u is an analytic function. This extends, partially, the well-known Liouville theorem ([6]), which states that if $u \in ACL^n(\Omega; \mathbb{R}^n)$, $n \geq 3$, and the condition $\|\nabla u(x)\|^n = n^{n/2} |\det \nabla u(x)|$ is satisfied a.e. in Ω , then u is a restriction of a Möbius transformation.

1. Introduction

Let Ω be an open set of the Euclidean space $\mathbb{R}^n$. For $x = (x_1, \dots, x_n) \in \mathbb{R}^n$, we denote by $|x| = (x_1^2 + \dots + x_n^2)^{1/2}$ the norm of x . Let $m = m_n$ denote the usual Lebesgue measure on $\mathbb{R}^n$. Sometimes we use notation $dx = dx_1 \dots dx_n$ and $|D|$ instead of dm and $m(D)$, where $x = (x_1, \dots, x_n) \in \mathbb{R}^n$ and D is a Lebesgue measurable set in $\mathbb{R}^n$, respectively. For a given domain $\Omega \subset \mathbb{R}^n$, we say that a continuous mapping $u : \Omega \rightarrow \mathbb{R}^n$ is quasiregular (abbreviated qr) if

- (1) u is ACL^n , and
- (2) there exists a real number K , $K \geq 1$, such that

$$|u'(x)|^n \leq K J_u(x) \quad \text{a.e. on } \Omega, \quad (1.1)$$

where $|u'(x)| = \max_{|h|=1} |u'(x)h|$.

In this setting we shortly write that u is a K -qr mapping. For properties of qr-mappings see [1], [2], [3], [7] and [8]. If u is a K -qr and homeomorphic mapping then it is called K -quasiconformal or shortly K -q.c.

Mathematics Subject Classification: Primary 31B05 ; Secondary 31B10, 30C65.

Key words and phrases: quasiregular mappings, Möbius transformations.

Research of the second author was partially supported by MNTRS, Serbia, Grant No. 144 020.

Let

$$\|u'(x)\| = \sqrt{\sum_{i,j=1}^n (\partial_j u_i(x))^2}$$

denote the Hilbert–Schmidt norm of $u'(x)$, where $\partial_j = \partial_{x_j}$ denotes j –th partial derivative.

It is well known that if u is a K -qr mapping on Ω , then

$$\|u'(x)\|^n \leq n^{n/2} K J_u(x) \quad \text{a.e. on } \Omega. \quad (1.2)$$

In this paper we will consider generalized 1-quasiregular mappings, i.e. continuous mappings u satisfying the conditions $u \in ACL^n$ and

$$\|u'(x)\|^n \leq n^{n/2} |J_u(x)| \quad \text{a.e. on } \Omega. \quad (1.3)$$

2. The main result

Proposition 2.1 ([4, Section V.3]). *Let Ω be an open subset of $\mathbb{R}^n$ and let $u : \Omega \rightarrow \mathbb{R}^n$ be a ACL^n mapping satisfying the Lusin’s condition (N) (The condition (N) means that a mapping maps sets of measure zero to sets of measure zero). Then the function $y \mapsto N(y, u)$ is measurable on $\mathbb{R}^n$ and*

$$\int_{\mathbb{R}^n} N(y, u) dy = \int_{\Omega} |J_u(x)| dx, \quad (2.1)$$

where $J_u(x)$ is the Jacobian of u at x and $N(y, u)$ denotes the cardinal number of the set $u^{-1}(y)$ if the last set is finite and it is $+\infty$ in the other case.

Corollary 2.2. *Under the condition of the previous proposition there holds the inequality*

$$\int_{\Omega} |J_u(x)| dx \geq |u(\Omega)|. \quad (2.2)$$

The equality holds in (2.2) if and only if u is univalent on Ω .

For 1-qr mapping we also say generalized conformal mapping. The generalized Liouville theorem ([6]) states: for $n \geq 3$ every 1-qr mapping on a domain $\Omega \subset \mathbb{R}^n$, is a restriction of a Möbius transformation or a constant.

We extend (partially) this theorem as follows:

Theorem 2.3. *Let Ω be a domain in $\mathbb{R}^n$ and let $u : \Omega \rightarrow \mathbb{R}^n$ be a C^1 mapping such that*

$$\|u'(x)\|^n = n^{n/2} |J_u(x)|, \quad x \in \Omega. \quad (\text{We say that } u \text{ is absolutely conformal}). \quad (2.3)$$

Then, for $n = 2$, u is analytic or anti-analytic function. For $n \geq 3$, u is a restriction of a Möbius transformation or a constant.

PROOF. We first consider the case $n = 2$. Let $\Omega_0 = \{z \in \Omega : J_u(z) = 0\}$, $\Omega_1 = \{z \in \Omega : J_u(z) > 0\}$, $\Omega_2 = \{z \in \Omega : J_u(z) < 0\}$. Let u^1 and u^2 be the restrictions of u on Ω_1 and Ω_2 . Put $p = u_z$. Suppose that u is smooth and non constant function and the equality holds in (2.3). Then $u_z = u_{\bar{z}} = 0$ on Ω_0 , u^1 is conformal on Ω_1 , and u^2 is anti conformal on Ω_2 (and therefore $p = 0$ on Ω_2). Hence p is continuous on Ω and analytic on $\Omega_1 \cup \Omega_2$.

We will prove that p is analytic on Ω .

There are two cases:

- (a) If $z_0 \in \Omega_0$ is an interior point of Ω_0 , then p is analytic at z_0 .
- (b) If $z_0 \in \Omega_0$ is not an interior point of Ω_0 then $z_0 \in \partial\Omega_0 \setminus \partial\Omega = \partial(\Omega_1 \cup \Omega_2) \setminus \partial\Omega$.

Hence then there exists a sequence $z_n \in \Omega_1 \cup \Omega_2$ such that $\lim_{n \rightarrow \infty} z_n = z_0$.

It follows that $p_{\bar{z}}(z_n) = 0$ and therefore $\lim_{n \rightarrow \infty} p_{\bar{z}}(z_n) = 0$ and hence, since p is continuous, we find $p_{\bar{z}}(z_0) = 0$. Hence p is analytic in Ω . If Ω_2 is not empty set, according to the uniqueness theorem, this gives that $u_z(z) \equiv 0$ on Ω and hence u is anti analytic on Ω . Since u is analytic on Ω_1 , we first conclude that u is constant on Ω_1 and therefore that Ω_1 is empty set. In a similar way, if Ω_1 is not empty set, we conclude that u is analytic on Ω .

Hence u is analytic in Ω or it is anti-analytic in Ω . Note that the set $\Omega \setminus (\Omega_1 \cup \Omega_2)$ is discrete or it is equal to the set Ω .

We now consider the case $n > 2$.

Let $\Omega_0 = \{x \in \Omega : J_u(x) = 0\}$, $\Omega_1 = \{x \in \Omega : J_u(x) > 0\}$, $\Omega_2 = \{x \in \Omega : J_u(x) < 0\}$ and $\Omega^* = \Omega \setminus \Omega_0$. If u is a C^1 mapping, then the Hadamard inequality gives

$$|J_u| \leq \prod_{k=1}^n |\partial_k u| \quad (2.4)$$

and hence:

$$|J_u| \leq \left(\frac{\sum_{k=1}^n |\partial_k u|^2}{n} \right)^{\frac{n}{2}}, \quad \text{that is } n^{n/2} |J_u(x)| \leq \|u'(x)\|^n, \quad x \in \Omega. \quad (2.5)$$

Using the geometric interpretation of $J_u(x)$ one can show that the equality holds in (2.4) at a point $x \in \Omega^*$ if and only if the vectors $\partial_i u(x)$, $i = 1, 2, \dots, n$,

are orthogonal. Observe that if the equality holds in (2.5), then the equality holds in (2.4). Hence, for $x \in \Omega^*$, the equality holds in (2.5) if and only if

$$\langle \partial_i u, \partial_j u \rangle = \begin{cases} 0 & \text{if } i \neq j, \\ s^2 & \text{if } i = j, \end{cases}$$

where $s = s(x)$ is a real function on Ω^* . Hence $\|u'(x)\|^n = n^{n/2} |J_u(x)|$ for $x \in \Omega^*$ and since $u \in C^1$, $\|u'(x)\|^n = n^{n/2} |J_u(x)|$ for $x \in \Omega$.

If u is not a constant function, then Ω^* is not empty set. Suppose, for example, that Ω_1 is not an empty set and let $\hat{\Omega}$ be a component of Ω_1 .

Thus u is a generalized conformal mapping in $\hat{\Omega}$. Hence, by the generalized Liouville theorem i.e. GEHRING-RESHETNYAK's theorem see ([6]), every generalized conformal mapping in the space is a Möbius transformation. Hence u is the restriction to $\hat{\Omega}$ of a Möbius transformation A . Let $\omega \in \hat{\Omega}$. Then there exists a sequence $z_n \in \hat{\Omega}$, $n \in \mathbb{N}$, such that z_n tends ω . Since u is a C^1 function, $J_u(\omega) = \lim J_u(z_n) = \lim J_A(z_n)$. It is clear that $J_A(\omega) = \lim J_A(z_n)$ and therefore $J_u(\omega) > 0$ $\omega \in \hat{\Omega}$. Hence $\hat{\Omega}$ is closed-open in Ω and therefore $\Omega = \hat{\Omega}$. $\square$

Example 1. Let $n \geq 3$ and $x = (x_1, \dots, x_n)$,

$$u(x) = \begin{cases} (x_1, \dots, x_{n-1}, x_n) & \text{if } x_n \geq 0 \\ (x_1, \dots, x_{n-1}, -x_n) & \text{if } x_n \leq 0. \end{cases}$$

Then there hold (2.3) almost everywhere but u is not a Möbius transformation. This means that the condition u is C^1 in Theorem 2.3 is important.

Corollary 2.4. *Let Ω be a domain in the Euclidean space $\mathbb{R}^n$, $n \geq 2$, and $u \in ACL^n(\Omega)$ satisfying the condition (N).*

Then

$$\int_{\Omega} \|u'(x)\|^n dx \geq n^{n/2} |u(\Omega)|. \quad (2.6)$$

If u is a C^1 mapping, then the equation in (2.6) holds if and only if u is an injective conformal mapping or a constant mapping.

PROOF. Using (2.2) and (2.5) and Corollary 2.2 we obtain:

$$D_n(u) := \int_{\Omega} \left(\sum_{i=1}^n |\partial_i u|^2 \right)^{n/2} dm(x) \geq n^{n/2} \int_{\Omega} |J_u(x)| dm(x) \geq n^{n/2} |u(\Omega)|,$$

and consequently

$$D_n(u) \geq n^{n/2} |u(\Omega)|.$$

Suppose now that the equality holds in (2.6) and $u \in C^1$. Since u is C^1 , u satisfies the condition (N) and the equality holds in (2.5) for every $x \in \Omega$ (that is u satisfies condition (2.3) and therefore u is absolutely conformal). Thus, by Theorem 2.3, it is an analytic (or anti-analytic) function in the plane or a Möbius transformation in the space. Thus if $n \geq 3$, then u is a Möbius transformation.

It remains to finish the proof in the case $n = 2$; that is to prove that if u is non-constant analytic (or anti-analytic) and

$$D_2(u) = 2|u(\Omega)|$$

then it is a univalent conformal (or anti-conformal) mapping.

Using the fact that u is an open mapping the assumption $u_0 = u(z_1) = u(z_2)$ for $z_1 \neq z_2$ has the consequence that there exist two disjoint open sets $U_1, U_2 \subset \Omega$ and a disk $D(u_0, r') \subset u(\Omega)$ such that $D(u_0, r') = u(U_1) = u(U_2)$. Hence

$$D_2(u) > \int_{\Omega \setminus U_1} \|u'(z)\|^2 dx dy \geq 2|(u(\Omega \setminus U_1))| = 2|\Omega'|$$

which is a contradiction. $\square$

AN OPEN QUESTION. From the proofs of the main theorems of the paper a question emerges: Does there exist a number $K_o = K_o(n) > 1$ such that if $u \in C^1$ satisfies $|u'(x)|^n \leq K|J(x, u)|$ and $K < K_o$, then u is a K -quasiregular mapping?

ACKNOWLEDGMENT. We thank the referee for useful comments and suggestions related to this paper. We thank professors M. VUORINEN, O. MARTIO and V. ZORICH for showing interest in this paper and giving comments concerning Theorem 2.3.

References

- [1] V. M. GOL'DSHTEIN and YU. G. RESHETNYAK, Introduction to the Theory of Functions With Generalized Derivatives, and Quasiconformal Mappings, *Nauka, Moscow*, 1983 (in Russian).
- [2] T. IWANIEC and G. MARTIN, Geometric Function Theory and Nonlinear Analysis, Oxford Mathematical Monographs, *The Clarendon Press, Oxford University Press, New York*, 2001.
- [3] J. MALÝ and O. MARTIO, Lusin's condition (N) and mappings of the class $W^{1,n}$, *J. Reine Angew. Math.* **458** (1995), 19–36.
- [4] T. RADO and P. V. REICHELDERFER, Continuous Transformations in Analysis with an Introduction to Algebraic Topology, *Springer-Verlag, Berlin – Göttingen – Heidelberg*, 1955.

- [5] YU. G. RESHETNYAK, Some geometrical properties and mappings with generalized derivatives, *Sibirsk. Mat. Z.* **7** (1966), 886–919 (in *Russian*).
- [6] YU. G. RESHETNYAK, Liouville’s conformal theorem under minimal regularity hypotheses, *Sibirsk. Mat. Z.* **8** (1967), 835–840 (in *Russian*).
- [7] S. RICKMAN, Quasiregular Maps, *Springer-Verlag, Berlin – Heidelberg*, 1993.
- [8] M. VUORINEN, Conformal Geometry and Quasiregular Mappings, *Lecture Notes in Math.* **1319**, *Springer-Verlag, Berlin – New York*, 1988.

DAVID KALAJ
UNIVERSITY OF MONTENEGRO
FACULTY OF NATURAL SCIENCES
AND MATHEMATICS
CETINJSKI PUT B.B.
81000 PODGORICA
MONTENEGRO

E-mail: davidk@t-com.me

MIODRAG MATELJEVIĆ
UNIVERSITY OF BELGRADE
FACULTY OF MATHEMATICS
STUDENTSKI TRG 16
11000 BELGRADE
SERBIA

E-mail: miodrag@matf.bg.ac.rs

(Received February 16, 2009; revised January 2, 2010)