

Derivations in commutators with power central values in rings

By YI-QIU DU (Changchun) and YU WANG (Shanghai)

Abstract. Let R be a prime ring of characteristic different from 2 and I a nonzero ideal of R , d a nonzero derivation of R such that $[d(x^k), x^k]^n$ is central, for all $x \in I$ where k, n are fixed positive integers. Then R satisfies s_4 , the standard identity in 4 variables.

1. Introduction

Throughout this article, R is always a prime ring with center Z . For any $x, y \in R$, we set $[x, y]_1 = [x, y] = xy - yx$ and $[x, y]_n = [[x, y]_{n-1}, y]$ where $n > 1$ is a positive integer. By s_4 we denote the standard identity in 4 variables. By d we denote a nonzero derivation of R .

A well-known result proved by POSNER [14] states that R must be commutative if $[d(x), x] \in Z$ for all $x \in R$. In [12] LEE and LEE generalized Posner's result by showing that if $\text{char}(R) \neq 2$ and $[d(x), x] \in Z$ for all x in a noncentral Lie ideal of R , then R is commutative. As to the case when $\text{char}(R) = 2$, LANSKI obtained the same conclusion except when R satisfies s_4 (see [9]). In [2] CARINI and DE FILIPPIS studied the situation when $[d(x), x]^n \in Z$ for all x in a noncommutative Lie ideal of R with $\text{char}(R) \neq 2$. In [16] the second author and You removed the assumption of $\text{char}(R) \neq 2$.

Mathematics Subject Classification: 16W25, 16R50, 16U80, 16N60.

Key words and phrases: derivation, generalized polynomial identity, differential identity, prime ring.

The second author has been partially supported by the Leading Academic Discipline Project of Shanghai Normal University (No.DZL803).

The second author is the corresponding author.

In [6] FELZENSZWALB proved that R is commutative if $d(x^k) = 0$ for all $x \in R$, where k is a fixed positive integer. A significant extension of [6] shows that R is commutative if $[d(x^k), x^k]_n = 0$ for all x in a nonzero left ideal of R (see [11, Theorem 1]). In [15] SHIUE discussed the situation when $a[d(x^k), x^k]_n = 0$ for all x in an one-sided ideal of R , where $0 \neq a \in R$.

The purpose of this paper is to investigate the situation when $[d(x^k), x^k]^n \in Z$ for all x in a nonzero ideal of R . The main result is the following

Theorem 1. *Let R be a prime ring of characteristic different from 2 with its center Z , I a nonzero ideal of R , and d a nonzero derivation of R such that $[d(x^k), x^k]^n \in Z$ for all $x \in I$ where k, n are fixed positive integers. Then R satisfies s_4 , the standard identity in 4 variables.*

The following counterexample shows that Theorem 1 is not valid on some one-sided ideals.

Example 1. Let F be a field and $R = M_m(F)$, the ring of all $m \times m$ matrix algebra over F with $m > 2$. Let e_{ij} be the matrix unit with 1 in (i, j) -entry and zero elsewhere. It is easy to check that $([e_{11}, x^k]_2)^n = 0$ for all $x \in Re_{22}$ (or, $e_{22}R$), where $n > 1$.

2. The proof of Theorem 1

By Q we denote the Martindale quotient ring of R and C the extended centroid R . The definitions and properties of these objects can be found in [1, Chapter 2].

We begin with the following easy result.

Lemma 1. *Let $R = M_2(F)$, the ring of all 2×2 matrices over a field F with $\text{char}(F) \neq 2$. If a is a nonzero element of R such that $([a, x^k]_2)^n = 0$ for all $x \in R$, then $a \in F \cdot I_2$.*

PROOF. Let $a = \sum_{i,j} a_{ij}e_{ij}$ with $a_{ij} \in F$. We first claim that a is a diagonal matrix. By assumption we get

$$0 = ([a, e_{11}]_2)^{2n} = (a_{12}a_{21})^n e_{11} + (a_{12}a_{21})^n e_{22},$$

thus $a_{12}a_{21} = 0$. Without loss of generality we may assume that $a_{21} = 0$. Let $\varphi \in \text{Aut}_F(M_2(F))$ such that $\varphi(x) = (1 + e_{21})x(1 - e_{21})$. In particular, we have

$$\varphi(a) = (a_{11} - a_{12})e_{11} + a_{12}e_{12} + (a_{11} - a_{12} - a_{22})e_{21} + (a_{12} + a_{22})e_{22}.$$

Since $([\varphi(a), x^k]_2)^n = 0$ for all $x \in R$, as above we can get that $a_{12}(a_{11} - a_{12} - a_{22}) = 0$. That is, either $a_{12} = 0$ or $a_{11} - a_{12} - a_{22} = 0$. If $a_{11} - a_{12} - a_{22} = 0$, then

$$[a, e_{11} + e_{21}]_2 = -a_{12}e_{11} + a_{12}e_{12} - 3a_{12}e_{21} + a_{12}e_{22}.$$

By assumption we get

$$0 = ([a, e_{11} + e_{21}]_2)^{2n} = (-2a_{12}^2)^n e_{11} + (-2a_{12}^2)^n e_{22}$$

and so $a_{12} = 0$, this implies that a is a diagonal matrix.

Write $a = \sum_{i=1}^2 a_{ii}e_{ii}$, we see as above that $\varphi(a) = \sum_{i=1}^2 a_{ii}e_{ii} + (a_{11} - a_{22})e_{21}$ is also a diagonal matrix. Therefore $a_{11} = a_{22}$ and so $a \in F \cdot I_2$ as desired. $\square$

If $([a, x^k]_2)^n \in F \cdot I_2$ for all $x \in M_2(F)$, one can not expect to obtain that $a \in F \cdot I_2$. For example, it is easy to check that $([e_{11}, x]_2)^2 \in F \cdot I_2$ for all $x \in M_2(F)$.

Lemma 2. *Let $R = M_m(F)$, the ring of all $m \times m$ matrices over a field F with $\text{char}(F) \neq 2$. If a is a noncentral element of R such that $([a, x^k]_2)^n \in F \cdot I_m$ for all $x \in R$, then $m \leq 2$.*

PROOF. Suppose on the contrary that $m > 2$. Let $a = \sum a_{ij}e_{ij}$ with $a_{ij} \in F$. Write $a = \begin{pmatrix} a_{11} & A \\ B & C \end{pmatrix}$, where $A = (a_{12}, \dots, a_{1m})$, $B = (a_{21}, \dots, a_{m1})^T$, and $C = (a_{ij})$ with $2 \leq i, j \leq m$. Since $[a, e_{11}]_2 = \begin{pmatrix} 0 & A \\ B & 0 \end{pmatrix}$, by assumption we have

$$([a, e_{11}]_2)^{2n} = \begin{pmatrix} (AB)^n & 0 \\ 0 & (BA)^n \end{pmatrix} \in F \cdot I_m.$$

Set $\alpha = AB \in F$. Then $\begin{pmatrix} \alpha^n & 0 \\ 0 & \alpha^{n-1}BA \end{pmatrix} \in F \cdot I_m$. If $\alpha \neq 0$, then

$$\begin{pmatrix} \alpha & 0 \\ 0 & BA \end{pmatrix} \in F \cdot I_m.$$

Thus, $\alpha = a_{21}a_{12} = a_{31}a_{13}$ and $a_{21}a_{13} = 0$. Thus $\alpha = 0$, a contradiction. Hence $AB = 0$.

Let φ_{ij} be an inner automorphism of R given by $\varphi_{ij}(x) = (1 + e_{ij})x(1 - e_{ij})$ for all $x \in R$. Write $1 + e_{21} = \begin{pmatrix} 1 & 0 \\ E_2 & I_{m-1} \end{pmatrix}$, where $E_2 = (1, 0, \dots, 0)^T$ and I_{m-1} is the $(m-1)$ -identity matrix. So

$$\varphi_{21}(a) = \begin{pmatrix} a_{11} - a_{12} & A \\ a_{11}E_2 - a_{12}E_2 + B - CE_2 & E_2A + C \end{pmatrix}.$$

Since $([\varphi_{21}(a), x^k]_2)^n \in F \cdot I_m$ for all $x \in R$, as above we have

$$A(a_{11}E_2 - a_{12}E_2 + B + CE_2) = 0.$$

Recalling $AB = 0$ we get from the last relation that $a_{11}a_{12} - a_{12}^2 - ACE_2 = 0$.

Since $[a, e_{11} + e_{21}]_2 = \begin{pmatrix} -a_{12} & A \\ D & E_2A \end{pmatrix}$, where $D = B + CE_2 - (a_{11} + 2a_{12})E_2$, we get

$$([a, e_{11} + e_{21}]_2)^2 = \begin{pmatrix} a_{12}^2 + AD & 0 \\ -a_{12}D + E_2AD & DA + a_{12}E_2A \end{pmatrix}.$$

Making use of both $AB = 0$ and $a_{11}a_{12} - a_{12}^2 - ACE_2 = 0$ we get $AD = -3a_{12}^2$.

Thus

$$([a, e_{11} + e_{21}]_2)^2 = \begin{pmatrix} -2a_{12}^2 & 0 \\ -a_{12}D - 3a_{12}^2E_2 & DA + a_{12}E_2A \end{pmatrix},$$

Therefore

$$([a, e_{11} + e_{21}]_2)^{2n} = \begin{pmatrix} (-2a_{12}^2)^n & 0 \\ U & (DA + a_{12}E_2A)^n \end{pmatrix} \in F \cdot I_m$$

where U is a $(m-1) \times 1$ matrix. Since $\text{rank}((DA + a_{12}E_2A)^n) \leq \text{rank}(A) \leq 1$ and $m > 2$, we infer that $(-2a_{12}^2)^n = 0$ and so $a_{12} = 0$.

Now we claim that a is a diagonal matrix. Since $([\varphi_{j2}(a), x^k]_2)^n \in F \cdot I_m$ for all $x \in R$, where $j > 2$, as above we have that $-a_{1j} = \varphi_{j1}(a)_{12} = 0$. So $a_{1j} = 0$ for $j > 1$. For $1 < j < t \leq m$, as above we get from $([\varphi_{1j}(a), x^k]_2)^n \in F \cdot I_m$ for all $x \in R$, that $a_{jt} = \varphi_{1j}(a)_{1t} = 0$. This shows that a must be lower triangular. Since the transpose of a satisfies the same condition, a is indeed diagonal.

We have showed that $a = \sum_{i=1}^m a_{ii}e_{ii}$ with $a_{ii} \in F$. For $1 \leq i \neq j \leq m$, as above we get that $\varphi_{ij}(a)$ is a diagonal matrix. On the other hand $\varphi(a) = a + (a_{jj} - a_{ii})e_{ij}$, we infer that $a_{jj} = a_{ii}$ and so a is central in R , which is a contradiction. The proof is thereby complete. $\square$

The following result is a special case of Theorem 1, which is of independent interest.

Lemma 3. *Let R be a prime ring with $\text{char}(R) \neq 2$ and I a nonzero ideal of R , d a nonzero derivation of R such that $[d(x^k), x^k]^n = 0$ for all $x \in I$ where k, n are fixed positive integers. Then R is commutative.*

PROOF. By assumption we see that I satisfies the differential identity

$$\left[\sum_{i=0}^{k-1} x^i d(x) x^{k-i-1}, x^k \right]^n = 0.$$

If d is not Q -inner, by KHARCHENKO's theorem [7], I satisfies the polynomial identity $\left[\sum_{i=0}^{k-1} x^i y x^{k-i-1}, x^k\right]^n = 0$ and so for R too. It is well known that there exists a field F such that R and F_m satisfy the same polynomial identities [8, p. 57 and p. 89]. Suppose that $m \geq 2$. If we choose $x = e_{11}, y = e_{12} + e_{21}$, then we get a contradiction as follows

$$0 = \left[\sum_{i=0}^{k-1} e_{11}^i (e_{12} + e_{21}) e_{11}^{k-i-1}, e_{11} \right]^{2n} = [e_{12} + e_{21}, e_{11}]^{2n} = (-1)^n (e_{11} + e_{22}) \neq 0.$$

Thus $m = 1$ and so R is commutative.

Assume next that d is Q -inner, that is, $d(x) = [a, x]$ for all $x \in R$, where a is a noncentral element in Q . By assumption we get $([a, x^k]_2)^n = 0$ for all $x \in I$. By a theorem of CHUANG [4, Theorem 2], $([a, x^k]_2)^n = 0$ for all $x \in Q$. In case C is infinite, we have $([a, x^k]_2)^n = 0$ for all $x \in Q \otimes_C \bar{C}$, where $\bar{C}$ is the algebraic closure of C . Since both Q and $Q \otimes_C \bar{C}$ are centrally closed [5, Theorems 2.5 and 3.5], we may replace R by Q or $Q \otimes_C \bar{C}$ according as C is finite or infinite. Thus we may assume that R is centrally closed over C which is either finite or algebraically closed and $([a, x^k]_2)^n = 0$ for all $x \in R$. By MARTINDALE's theorem [13], R is a primitive ring and so isomorphic to a dense subring of linear transformations on a vector space V over C .

If V is infinite dimensional over C , for any given $v \in V$ we claim that v and va are C -dependent. Suppose on the contrary that v and va are C -independent. We choose $v_1, \dots, v_{2k-1}$ such that $v, va, v_1, \dots, v_{2k-1}$ are C -independent. By the density of R on ${}_C V$, there exists $x \in R$ such that $vx = 0$, $va x = v_1$, $v_i x = v_{i+1}$, $v_{2k-1} x = v$, where $i = 1, \dots, 2k-2$. Thus

$$v[a, x^k]_2 = va x^{2k} = v_1 x^{2k-1} = \dots = v_{2k-1} x = v$$

and so $0 = v([a, x^k]_2)^n = v$, a contradiction. Therefore v and va are C -dependent for any $v \in V$. A standard argument shows that $a \in C$, a contradiction. So V must be of finite dimension. That is, $R \cong M_s(C)$ for some s . In view of both Lemma 1 and Lemma 2 we get that $a \in C$, a contradiction. The proof is now complete. $\square$

THE PROOF OF THEOREM 1. Suppose on the contrary that $\dim_C RC > 4$. By assumption we have $[d(x^k), x^k]^n \in Z$ for all $x \in I$, that is, I satisfies the following differential identity

$$\left[\left[\sum_{i=0}^{k-1} x^i d(x) x^{k-i-1}, x^k \right]^n, y \right] = 0. \quad (1)$$

If $[d(x^k), x^k]^n = 0$ for all $x \in I$, the result follows from Lemma 3. Otherwise, there exists $r \in I$ such that $[d(r^k), r^k]^n \neq 0$. Thus I satisfies the central differential identity $[d(x^k), x^k]^n$. By [3, Theorem 1] we get that R is a PI-prime ring and so is Q .

If d is not Q -inner, applying Kharchenko's theorem to (1) we get that I satisfies the polynomial identity $[[\sum_{i=0}^{k-1} x^i y x^{k-i-1}, x^k]^n, r] = 0$ and so for R too. It is well known that there exists a field F such that R and F_m satisfy the same polynomial identities. Thus $[\sum_{i=0}^{k-1} x^i y x^{k-i-1}, x^k]^n \in F \cdot I_m$. Note that $m > 2$. If we choose $x = e_{11}, y = e_{12} + e_{21}$, then

$$\left[\sum_{i=0}^{k-1} e_{11}^i (e_{12} + e_{21}) e_{11}^{k-i-1}, e_{11}^k \right]^{2n} = (-1)^n (e_{11} + e_{22}) \in F \cdot I_m.$$

This is a contradiction.

We next assume that d is an Q -inner derivation induced by a noncentral element $b \in Q$. It follows from (1) that

$$\left[\left[\sum_{i=0}^{k-1} x^i [b, x] x^{k-i-1}, x^k \right]^n, y \right] = 0 \quad \text{for all } x, y \in I. \quad (2)$$

In view of [4, Theorem 2] we have

$$\left[\left[\sum_{i=0}^{k-1} x^i [b, x] x^{k-i-1}, x^k \right]^n, y \right] = 0 \quad \text{for all } x, y \in Q. \quad (3)$$

Since there exists $r \in R$ such that $[d(r^k), r^k]^n \neq 0$, we see that (3) is a non trivial generalized polynomial identity on Q . By MARTINDALE's theorem [13] Q is primitive ring. Since Q is a PI-ring, by the famous Kaplansky's theorem [1, Theorem 6.1.10] we see that Q is a finite dimensional central simple algebra over C . It follows from [10, Lemma 2] that there exists a suitable field F of $\text{char}(F) \neq 2$ such that $Q \subseteq M_m(F)$ and moreover $M_m(F)$ satisfies the same generalized polynomial identity (3). Then Lemma 2 tells us that $m \leq 2$, which is a contradiction. The proof is thereby complete. $\square$

ACKNOWLEDGEMENT. The authors would like to express their sincere thanks to the referee for mending a gap in the proof of Lemma 2 and pointing out several misprints.

References

- [1] K. I. BEIDAR, W. S. MARTINDALE 3rd and A. V. MIKHALEV, Rings with Generalized Identities, *Dekker, New York*, 1996.

- [2] L. CARINI and V. DE FILIPPIS., Commutators with power central values on a Lie ideal, *Pacific J. Math.* **193** (2000), 269–278.
- [3] M. CHANG and T. K. LEE, Annihilators of power values of derivations in prime rings, *Comm. Algebra* **26** (1998), 2091–2113.
- [4] C. L. CHUANG, GPIs having coefficients in Utumi quotient rings, *Proc. Amer. Math. Soc.* **103** (1988), 723–728.
- [5] T. S. ERICKSON, W. S. MARTINDALE 3rd and J. M. OSBORN, Prime nonassociative algebras, *Pacific J. Math.* **60** (1975), 49–63.
- [6] B. FELZENSZWALB, Derivations in prime rings, *Proc. Amer. Math. Soc.* **84** (1982), 16–20.
- [7] V. K. KHARCHENKO, Differential identities of prime rings, *Algebra and Logic* **17** (1978), 155–168.
- [8] N. JACOBSON, PI-Algebras, An introduction, Lecture Notes in Math., *Springer, Verlag, New York*, 1975.
- [9] C. LANSKI, Differential identities, Lie ideals, and Posner's theorems, *Pacific J. Math.* **134** (1988), 275–297.
- [10] C. LANSKI, An Engel condition with derivation, *Proc. Amer. Math. Soc.* **118** (1993), 731–734.
- [11] C. LANSKI, An Engel condition with derivation for left ideals, *Proc. Amer. Math. Soc.* **125** (1997), 339–345.
- [12] P. H. LEE and T. K. LEE, Lie ideals of prime rings with derivations, *Bull. Ins. Math. Acad. Sinica* **11** (1983), 75–80.
- [13] W. S. MARTINDALE 3rd, Prime rings satisfying a generalized polynomial identity, *J. Algebra* **12** (1969), 576–584.
- [14] E. C. POSNER, Derivation in prime rings, *Proc. Amer. Math. Soc.* **8** (1957), 1093–1100.
- [15] E. K. SHIUE, Annihilators of derivations with Engel conditions on one-sided ideals, *Publ. Math. Debrecen* **62** (2003), 237–243.
- [16] Y. WANG and H. YOU, Commutators with power center values on Lie ideals in rings, *Acta Math. Sinica, English series* **22** (2006), 1715–1720.

YI-QIU DU
 INSTITUTE OF MATHEMATICS
 JILIN UNIVERSITY
 CHANGCHUN 130012
 CHINA
 AND
 COLLEGE OF MATHEMATICS
 JILIN NORMAL UNIVERSITY
 SIPING 136000
 CHINA

E-mail: duyiqu-2006@163.com

YU WANG
 MATHEMATICS AND SCIENCE COLLEGE
 SHANGHAI NORMAL UNIVERSITY
 SHANGHAI 200234
 CHINA

E-mail: ywang2004@126.com

(Received July 31, 2009; revised November 10, 2009)