

A finiteness condition for verbal conjugacy classes in a group

By JOSE M. MUÑOZ-ESCOLANO (Zaragoza) and PAVEL SHUMYATSKY (Brasilia)

Abstract. Given a group G and a word w , we denote by G_w the set of all w -values in G and by $w(G)$ the corresponding verbal subgroup. The main result of the paper is the following theorem. Let k be a positive integer and let w be either the word γ_k or the word δ_k . Suppose that G is a group in which $\langle x^{G^w} \rangle$ is Chernikov for all $x \in G$. Then $\langle x^{w(G)} \rangle$ is Chernikov for all $x \in G$ as well.

1. Introduction

Let w be a word in n variables, and let G be a group. The verbal subgroup $w(G)$ of G determined by w is the subgroup generated by the set G_w consisting of all values $w(g_1, \dots, g_n)$, where $g_1, \dots, g_n$ are elements of G . A word w is said to be concise if whenever G_w is finite for a group G , it always follows that $w(G)$ is finite. P. Hall asked whether every word is concise, but it was later proved that this problem has a negative solution in its general form (see [4], p. 439). On the other hand, many relevant words are known to be concise. For instance, TURNER-SMITH [7] showed that the lower central words γ_k and the derived words δ_k are concise; here the words γ_k and δ_k are defined by the positions $\gamma_1 = \delta_0 = x$, $\gamma_{k+1} = [\gamma_k, \gamma_1]$ and $\delta_{k+1} = [\delta_k, \delta_k]$. The corresponding verbal subgroups for these words are the familiar k th term of the lower central series of G denoted by $\gamma_k(G)$ and the k th derived group of G denoted by $G^{(k)}$.

Mathematics Subject Classification: 20E45, 20F24.

Key words and phrases: words, verbal subgroups, conjugacy classes.

The first author was supported by Proyecto MTM2010-19938-C03-03 of Dirección General de Investigación of MICINN (Spain). The second author was supported by CNPq-Brazil.

There are several natural ways to look at Hall's question from a different angle. The circle of problems arising in this context can be characterized as follows.

Given a word w and a group G , assume that certain restrictions are imposed on the set G_w . How does this influence the properties of the verbal subgroup $w(G)$?

If X and Y are non-empty subsets of a group G , we will write X^Y to denote the set $\{y^{-1}xy \mid x \in X, y \in Y\}$. In [2] groups G with the property that x^{G_w} is finite for all $x \in G$ were called $FC(w)$ -groups. Recall that FC -groups are precisely groups with finite conjugacy classes. The main result of [2] tells us that if w is a concise word, then a group G is an $FC(w)$ -group if and only if $x^{w(G)}$ is finite for all $x \in G$. In particular, it follows that if w is a concise word and G is an $FC(w)$ -group, then the verbal subgroup $w(G)$ is FC . Later it was shown in [1] that there exists a function $f = f(m, w)$ such that if, under the hypothesis of the above theorem, x^{G_w} has at most m elements for all $x \in G$, then $x^{w(G)}$ has at most f elements for all $x \in G$. In view of these results we would like to consider the following question.

Given a concise word w and a group G , assume that for all $x \in G$ the subgroup $\langle x^{G_w} \rangle$ satisfies a certain finiteness condition. Is it true that a similar condition is also satisfied by $\langle x^{w(G)} \rangle$ for all $x \in G$?

Here and throughout the paper $\langle M \rangle$ denotes the subgroup generated by the set M . The main result of the present paper is as follows.

Theorem 1.1. *Let k be a positive integer and let w be either the word γ_k or the word δ_k . Suppose that G is a group in which $\langle x^{G_w} \rangle$ is Chernikov for all $x \in G$. Then $\langle x^{w(G)} \rangle$ is Chernikov for all $x \in G$ as well.*

Recall that a group G is Chernikov if it has a subgroup of finite index that is a direct product of finitely many groups of type C_{p^∞} for various primes p (quasicyclic p -groups). By a deep result obtained independently by SHUNKOV [6] and KEGEL and WEHRFRITZ [3] Chernikov groups are precisely the locally finite groups satisfying the minimal condition on subgroups, that is, any non-empty set of subgroups possesses a minimal subgroup. The minimal subgroup of finite index of a Chernikov group G is called the radicable part of G . In general a group G is called radicable if the equation $x^n = a$ has a solution in G for every positive integer n and every $a \in G$. It is well-known that a periodic abelian radicable group is a direct product of quasicyclic p -subgroups.

A proof of Theorem 1.1 in the case where $w = \gamma_k$ can be obtained from

the case $w = \delta_k$ by simply replacing everywhere in the proof the term “ δ_k -commutators” by “ γ_k -commutators”. That is why we do not provide an explicit proof for the case $w = \gamma_k$ concentrating instead on proving Theorem 1.1 in the case $w = \delta_k$.

Most of this work was done while the first author was visiting the Department of Mathematics of the University of Brasilia. He thanks the Department of Mathematics for support and hospitality.

2. Preliminary lemmas

We start the section with the following well-known lemma (see for example [5, Lemma 3.13]).

Lemma 2.1. *Suppose that R is a radicable abelian normal subgroup of the group G and suppose that H is a subgroup of G such that $[R, \underbrace{H, \dots, H}_r] = 1$ for some natural number r . If H/H' is periodic, then $[R, H] = 1$.*

From this we can easily deduce the following useful corollaries.

Corollary 2.2. *In a periodic nilpotent group G every radicable abelian subgroup Q is central.*

PROOF. Arguing by induction on the nilpotency class of G we assume that the image of Q in $G/Z(G)$ is central. Therefore Q is contained in a normal abelian subgroup of G . In particular $\langle Q^G \rangle$ is a normal abelian radicable subgroup and the result is now immediate from Lemma 2.1. $\square$

Let G be a group acted on by a group A . As usual, $[G, A]$ denotes the subgroup generated by all elements of the form $x^{-1}x^a$, where $x \in G$, $a \in A$. It is well-known that $[G, A]$ is a normal subgroup of G . If B is a normal subset of A such that $A = \langle B \rangle$, then $[G, A] = \langle [G, b]; b \in B \rangle$.

Corollary 2.3. *Let A be a periodic group acting on a periodic radicable abelian group G . Then $[G, A, A] = [G, A]$.*

PROOF. To show this, we can assume that $[G, A, A] = 1$. In this case Lemma 2.1 yields at once that $[G, A] = 1$. $\square$

Lemma 2.4. *Let A be a finite group acting on a periodic radicable abelian group G . Then $[G, A]$ is radicable.*

PROOF. Since $[G, A] = \prod_{a \in A} [G, a]$, it is sufficient to show that $[G, a]$ is radicable for every $a \in A$. Let $x \in [G, a]$ and let n be a positive integer. Then there exist $g \in G$ such that $x = [g, a]$ and $g_1 \in G$ such that $g_1^n = g$. Since G is abelian, we have $[g_1, a]^n = [g_1^n, a]$. Hence for every $x \in [G, a]$ and every positive integer n , there exists an element $[g_1, a] \in [G, a]$ such that $x = [g_1, a]^n$; that is, $[G, a]$ is radicable, as required. $\square$

Lemma 2.5. *Let A be a radicable Chernikov group acting on a Chernikov group B . Then $[B, A, A] = 1$.*

PROOF. Denote by B_0 the radicable part of B . By [5, Theorem 3.29.2], $A/C_A(B_0)$ is finite. Since A is radicable, it follows that A has no subgroups of finite index and so $[B_0, A] = 1$. On the other hand, B/B_0 is finite and therefore $A/C_A(B/B_0)$ is also finite. Again, since A has no subgroups of finite index, it follows that $[B, A] \leq B_0$. Hence $[B, A, A] \leq [B_0, A] = 1$. $\square$

Lemma 2.6. *Let G be a group and y an element of G . Suppose that $x_1, \dots, x_k \in G$ are δ_k -commutators for $k \geq 0$. Then $[y, x_1, \dots, x_k]$ is a δ_k -commutator as well.*

PROOF. Note that $x_1, \dots, x_k$ can be viewed as δ_i -commutators for each $i \leq k$. It is clear that $[y, x_1]$ is a δ_1 -commutator. Arguing by induction on k assume that $k \geq 1$ and $[y, x_1, \dots, x_{k-1}]$ is a δ_{k-1} -commutator. Then $[y, x_1, \dots, x_k] = [[y, x_1, \dots, x_{k-1}], x_k]$ is a δ_k -commutator. $\square$

Throughout the paper, whenever G is a Chernikov group we denote by G_0 the radicable part of G and by G^* the subgroup $[G_0, G]$.

Lemma 2.7. *Let G be a Chernikov group for which there exists a positive integer m such that G can be generated by elements of order dividing m . If $G^* = 1$, then G is finite.*

PROOF. Since $G^* = 1$, it follows that G_0 is central. The Schur Theorem [5, Theorem 4.12] yields that G' is finite. Since G can be generated by elements of order dividing m , we conclude that G has finite exponent. In particular G has no subgroups of type C_{p^∞} . Thus, G must be finite. $\square$

Lemma 2.8. *Let G be a group such that $\langle x^G \rangle$ is Chernikov for every $x \in G$. Then all abelian radicable subgroups of G generate an abelian radicable subgroup.*

PROOF. Let T be the subgroup of G generated by all abelian radicable subgroups. Let A be an arbitrary abelian radicable subgroup in G and choose $x \in G$. Then Lemma 2.1 together with Lemma 2.5 shows that the product $\langle x^G \rangle_0 A$ is

abelian. Thus, all subgroups of the form $\langle x^G \rangle_0$ lie in the center of T . Therefore $G/Z(T)$ is a periodic FC -group. Since T has no subgroups of finite index, it centralizes every finite normal subgroup and we conclude that the image of T in $G/Z(T)$ is central. Therefore T is nilpotent of class at most two. Corollary 2.2 now enables us to deduce that T is abelian, as required. $\square$

We will also require the following lemma.

Lemma 2.9. *Let X be a normal set in a locally finite group G . Let $a \in G$ and assume that the set a^X is finite. Then the set $a^{\langle X \rangle}$ is likewise finite.*

PROOF. Let $x_1, \dots, x_n$ be elements of X with the property that $a^X = \{a^{x_1}, \dots, a^{x_n}\}$ and let $Y = \langle x_1, \dots, x_n \rangle$. Since Y is finite, the class a^Y is finite as well. Let $N = \langle X \rangle$. We will show that $a^N = a^Y$. Choose $y \in N$. Then y can be written as a product $y = y_1 \dots y_m$, where $y_i \in X$. It is sufficient to show that $a^y \in a^Y$. If $m = 1$, then $y \in X$ and so $a^y \in \{a^{x_1}, \dots, a^{x_n}\} \subseteq a^Y$. Thus, assume that $m \geq 2$ and use induction on m . Suppose that $a^{y_1} = a^{x_1}$. Set $z_i = x_1 y_i x_1^{-1}$ for $i = 2, \dots, m$. Since X is a normal set of G , $z_i \in X$. Write

$$a^y = a^{x_1 y_2 \dots y_m} = a^{x_1 y_2 \dots y_m x_1^{-1} x_1} = a^{z_2 \dots z_m x_1}.$$

By induction $a^{z_2 \dots z_m} \in a^Y$. Since $x_1 \in Y$, it follows that $a^y \in a^Y$. This completes the proof. $\square$

3. Proof of Theorem 1.1

Assume the hypothesis of Theorem 1.1 with $w = \delta_k$ and let X denote the set of all δ_k -commutators in G . By the hypothesis $\langle a^X \rangle$ is Chernikov for all $a \in G$. Set $H = G^{(k)}$. We wish to show that $\langle a^H \rangle$ is Chernikov for all $a \in G$. First we will deal with the particular case where $a \in X$. Thus, choose $a \in X$ and let $D = \langle a^X \rangle$. As usual, the normal closure of a subset $S \subseteq G$ is the minimal normal subgroup of G containing S .

Lemma 3.1. *With the above notation, the normal closure of D^* in G is an abelian radicable subgroup.*

PROOF. By Corollary 2.3 $D^* = [D_0, \underbrace{D, \dots, D}_k]$. Since $a \in X$, every element of a^X is also a δ_k -commutator. It follows that D is generated by the normal set $X \cap D$. Therefore the subgroup D^* is generated by subgroups of the form $[D_0, b_1, \dots, b_k]$, where $b_1, \dots, b_k \in X \cap D$.

Let us show that for every choice of $b_1, \dots, b_k \in X \cap D$ the subgroup $[D_0, b_1, \dots, b_k]$ is contained in a normal abelian radicable subgroup of G . Thus, fix $b_1, \dots, b_k \in X \cap D$ and put $K = [D_0, b_1, \dots, b_k]$. Since D_0 is abelian, it is clear that for every $d_1, d_2 \in D_0$ we have

$$[d_1, b_1, \dots, b_k][d_2, b_1, \dots, b_k] = [d_1 d_2, b_1, \dots, b_k].$$

Now, Lemma 2.6 shows that every element of K is a δ_k -commutator and Lemma 2.4 yields that K is radicable. Since $\langle g^K \rangle$ is Chernikov for every $g \in G$, it follows from Lemma 2.5 that $[g^K, K, K] = 1$. In particular $[g, K, K] = 1$ and we conclude that K commutes with K^g for every $g \in G$. Therefore $\langle K^G \rangle$ is abelian. Since $\langle K^G \rangle$ is generated by radicable subgroups, it follows that $\langle K^G \rangle$ is radicable.

Now choose other elements $b'_1, \dots, b'_k \in X \cap D$ and set $K_1 = [D_0, b'_1, \dots, b'_k]$. Repeating the above argument we conclude that $\langle K_1^G \rangle$ is abelian and radicable. Thus, the product $\langle K^G \rangle \langle K_1^G \rangle$ is nilpotent of class at most two and Corollary 2.2 tells us that $\langle K^G \rangle \langle K_1^G \rangle$ is abelian. Thus, all subgroups of the form $\langle [D_0, x_1, \dots, x_k]^G \rangle$, where $x_1, \dots, x_k \in X \cap D$, commute and the lemma follows. $\square$

Set $R = \langle \langle y^X \rangle^*; y \in X \rangle$. This notation will be kept throughout the rest of the paper.

Corollary 3.2. *The subgroup R is abelian and radicable.*

PROOF. Choose $y_1, y_2 \in X$. Let R_1 be the normal closure of $\langle y_1^X \rangle^*$ and R_2 that of $\langle y_2^X \rangle^*$. By Lemma 3.1 both R_1 and R_2 are abelian radicable subgroups. We conclude that the product $R_1 R_2$ is nilpotent of class at most two and Corollary 2.2 shows that $R_1 R_2$ is abelian. The result follows. $\square$

In the next lemma we use terminology and some results from the paper [2]. For the reader's convenience we will briefly explain it. Let w be a word, G a group and H a subgroup of $w(G)$. We say that H has finite w -index if the elements of G_w lie in finitely many right cosets of H in $w(G)$. A group G is an $FC(w)$ -group if and only if the subgroup $C_{w(G)}(x)$ has finite w -index for every element x of G .

Lemma 3.3. *The group G is locally finite.*

PROOF. First of all we notice that G is torsion since $\langle y^X \rangle$ is torsion for every $y \in G$. Since R is abelian (Corollary 3.2), it is sufficient to prove the local finiteness of G under the assumption that $R = 1$. Choose $y \in X$. By the above assumption $\langle y^X \rangle^* = 1$. Since $\langle y^X \rangle$ is generated by conjugates of y , it follows from Lemma 2.7 that $\langle y^X \rangle$ is finite. This implies that $C_G(y) \cap H$ has finite δ_k -index.

This happens for every choice of $y \in X$. Since every element of H is a product of finitely many δ_k -commutators, [2, Lemma 2.1] shows that H is an $FC(\delta_k)$ -group. In particular, the main result of [2] tells us that the k th derived group of H is an FC -group. Now the local finiteness of G is obvious. $\square$

Lemma 3.4. *The subgroup $\langle a^H \rangle$ is Chernikov.*

PROOF. Recall that $a \in X$ and $D = \langle a^X \rangle$. Set $E = \langle a^H \rangle$. We wish to show that E is Chernikov. Let us show first that ER/R is finite. It suffices to show this under the additional assumption that $R = 1$. In this case $D^* = 1$ and so D is finite by Lemma 2.7. In particular a^X is finite and since G is locally finite, we use Lemma 2.9 to conclude that E is finite. Thus, indeed ER/R is finite. Set $R_1 = E \cap R$. Choose elements $e_1, \dots, e_s \in X$ such that every conjugate of a in H belong to a coset $e_i R_1$ for some $i = 1, \dots, s$. We have $E = \langle R_1, e_1, \dots, e_s \rangle$. Since the set $e_1, \dots, e_s$ generates E and is normal in H modulo R_1 , it follows that

$$[R_1, E] = [R_1, e_1] \dots [R_1, e_s].$$

By Lemma 2.4 $[R_1, e_i] = [R_1, e_i, e_i]$ and Lemma 2.6 shows that $[R_1, e_i] \subseteq X$. We conclude that $[R_1, e_i] = [R_1, e_i, e_i] \leq [X, e_i] \leq \langle e_i^X \rangle$ and so the subgroups $[R_1, e_i]$ are Chernikov for every $i = 1, \dots, s$. Therefore $[R_1, E]$ is Chernikov and we can pass to the quotient $H\langle a \rangle/[R_1, E]$. Without loss of generality we assume that $[R_1, E] = 1$. In this case, $R_1 \leq Z(E)$ and $E/Z(E)$ is finitely generated. Lemma 3.3 shows that $E/Z(E)$ is finite. The Schur Theorem now tells us that the derived group E' is finite. We see that D is a Chernikov group generated by elements of the same order and its derived group D' is finite. It follows that D is finite. Now Lemma 2.9 enables us to deduce that E is finite. This completes the proof. $\square$

Corollary 3.5. *If $g \in H$, then $\langle g^H \rangle$ is Chernikov.*

PROOF. This follows directly from Lemma 3.4 and the fact that every element of H is a product of finitely many elements from X . $\square$

We are now ready to complete the proof of Theorem 1.1.

PROOF OF THEOREM 1.1. Combining Corollary 3.5 with Lemma 2.8 we deduce that all abelian radicable subgroups of H generate an abelian radicable subgroup. This will be denoted by T .

To complete the proof of Theorem 1.1 we need to show that $\langle b^H \rangle$ is Chernikov for every $b \in G$. Thus, let $b \in G$. Set $B = \langle b^X \rangle$ and $C = \langle b^H \rangle$. By the hypothesis, B is Chernikov. Since T contains all the abelian radicable subgroups of H , the

image of B in $H\langle b \rangle/T$ is finite. Therefore Lemma 2.9 shows that also the image of C is finite. Let us define

$$S = \langle [T, b_1, \dots, b_k] \mid b_i \in X \rangle.$$

For every choice $b_1, \dots, b_k \in X$ the subgroup $[T, b_1, \dots, b_k]$ is a radicable subgroup (Lemma 2.4) contained in X (Lemma 2.6). Thus, S is a normal radicable subgroup of G . Let $\{S_\lambda\}_{\lambda \in \Lambda}$ be the list of the radicable subgroups contained in $S \cap X$. Then $S = \langle S_\lambda \mid \lambda \in \Lambda \rangle$. Since $S_\lambda \subseteq X$, we have $[S_\lambda, b] \leq [X, b] \leq B$ for every λ and so we deduce that $[S, b] \leq B$. In particular, $[S, x]$ is Chernikov for every $x \in G$. Set $T_1 = C \cap T$. Now choose in C finitely many conjugates of b , say $c_1, \dots, c_n$, such that $C = \langle T_1, c_1, \dots, c_n \rangle$ and the set $c_1 T_1, \dots, c_n T_1$ is normal in C/T_1 . Then $[S, C] = [S, c_1] \dots [S, c_n]$. Since every subgroup $[S, c_i]$ is Chernikov, so is $[S, C]$. Moreover the subgroup $[S, C]$ is normal in $H\langle b \rangle$ and so we can consider the quotient $H\langle b \rangle/[S, C]$. Thus, we assume that $[S, C] = 1$.

Suppose temporarily that $S = 1$. Then T is contained in the k th term of the upper central series of H and Lemma 2.1 shows that actually $T \leq Z(H)$. In this case B_0 , the radicable part of B , is normal in $H\langle b \rangle$ and so we can consider the quotient $H\langle b \rangle/B_0$. The image of B in the quotient is finite. By Lemma 2.9 the image of C must be finite as well. This proves the theorem in the particular case where $S = 1$.

Let us now drop the assumption that $S = 1$. The above argument shows that the image of C in $H\langle b \rangle/S$ is Chernikov. Taking into account that $[S, C] = 1$ we deduce that $C/Z(C)$ is Chernikov. Polovickii's Theorem [5, p. 129] now tells us that C' , the derived group of C , is Chernikov and we can pass to the quotient $H\langle b \rangle/C'$. Thus, we assume that $C' = 1$. Now C is an abelian group generated by elements of the same order (namely, of order equal to that of b). We deduce that C has finite exponent. Taking into account that $B \leq C$ we observe that B is a Chernikov group of finite exponent. Hence, B is finite. But then by Lemma 2.9, C must be finite as well. The proof is now complete. $\square$

References

- [1] S. BRAZIL, A. KRASILNIKOV and P. SHUMYATSKY, Groups with bounded verbal conjugacy classes, *J. Group Theory* **9** (2006), 127–137.
- [2] S. FRANCIOSI, F. DE GIOVANNI and P. SHUMYATSKY, On group with finite verbal conjugacy classes, *Houston J. Math.* **28**(4) (2002), 683–689.
- [3] O. H. KEGEL and B. A. F. WEHRFRITZ, Strong finiteness conditions in locally finite groups, *Math. Z.* **117** (1970), 309–324.

- [4] A. YU, Olshanskii, Geometry of Defining Relations in Groups, *Kluwer Academic Publishers*, 1991.
- [5] D. J. S. ROBINSON, Finiteness Conditions and Generalized Soluble Groups, Part 1, *Springer, Berlin*, 1973.
- [6] V. P. SHUNKOV, On the minimality problem for locally finite groups, *Algebra Logic* **9** (1970), 137–151.
- [7] R. F. TURNER-SMITH, Finiteness conditions for verbal subgroups, *J. London Math. Soc.* **41** (1966), 166–176.

JOSE M. MUÑOZ-ESCOLANO
DEPARTMENT OF MATHEMATICS – IUMA
UNIVERSITY OF ZARAGOZA
PEDRO CERBUNA 12
50009 ZARAGOZA
SPAIN

E-mail: jmescola@unizar.es

PAVEL SHUMYATSKY
DEPARTMENT OF MATHEMATICS
UNIVERSITY OF BRASILIA
BRASILIA-DF, 70910-900
BRAZIL

E-mail: pavel@mat.unb.br

(Received July 14, 2011; revised October 24, 2011)