

Grünwald shift spaces

By JOSEF MIKEŠ (Olomouc) and KARL STRAMBACH (Erlangen)

Abstract. An n -dimensional differentiable shift space $\mathcal{S}$ for which in case $n = 2$ there exists an affine connection if $\mathcal{S}$ is a Grünwald plane (cf. [13, § 4]) admits for $n \geq 3$ no affine connection. In contrast to this the set of all images of the system of curves arising by shifting the argument from a Grünwald curve $\mathcal{C}$ under the translation group of $\mathbb{R}^n$ is a system of geodesics with respect to a metrizable affine connection if and only if $\mathcal{C}$ is a curve corresponding to parabolas in a suitable coordinate system.

1. Introduction

The investigation of systems $\mathfrak{S}$ of curves in the plane $\mathbb{R}^2$ such that any two different points are incident with precisely one curve of $\mathfrak{S}$ has a long tradition (see e.g. [17]). In particular since the second half of the previous century the systems $\mathfrak{S}$ has been studied intensively as natural generalisations of the real affine plane and the 2-dimensional hyperbolic geometry. These geometries, now called $\mathbb{R}^2$ -planes, are classified if they admit an at least 3-dimensional Lie group of automorphisms [15, Chapter 3].

Although already E. Beltrami has shown that a differentiable curve is a local geodesic with respect to an affine connection ∇ precisely if it is a solution of an Abelian differential equation having as coefficients expressions in Christoffel symbols associated with ∇ , the use of differential geometry for study of $\mathbb{R}^2$ -planes having differentiable curves as lines started only 2000 by G. GERLICH [5], [6], [7].

Mathematics Subject Classification: Primary: 51H20; Secondary: 53B05, 53C22.

Key words and phrases: shift space, extended translation shell of a curve, affine connection, geodesic.

The paper was supported by grant P201/11/0356 of The Czech Science Foundation and by the Department Mathematik der Universität Erlangen-Nürnberg.

He asked for which $\mathbb{R}^2$ -planes A with differentiable lines there exists an affine connection ∇ generating the lines of A and for affine planes A with an at least three-dimensional collineation group he proved that ∇ exists if and only if A is either desarguesian or a Moulton plane. Moreover, in [13] it is shown that the differentiable lines of a generalized shift $\mathbb{R}^2$ -plane A are geodesics with respect to an affine connection ∇ precisely if A is either the Euclidean plane or a Grünwald model of the real affine plane (see [8]).

The extension of the investigation from $\mathbb{R}^2$ -planes to geometries on $\mathbb{R}^n$ having as lines a system $\mathfrak{S}$ of curves such that any two different points are incident with precisely one curve of $\mathfrak{S}$ surprisingly turns out to be difficult as one can see in the papers [1], [2], [3], where D. BETTEN created a theory of 3-dimensional topological incidence geometries.

If one tries to extend the characterization of differentiable shift spaces having as lines geodesics with respect to an affine connection starting with a Grünwald plane (cf. [13, § 4]), then one meets also great difficulties. Namely, we show that for at least 3-dimensional differentiable shift spaces $\mathcal{S}$ generalizing in a natural way the 2-dimensional shift spaces corresponding to Grünwald planes there exists no affine connection ∇ such that the lines of $\mathcal{S}$ are geodesics of ∇ . This is surprising since there exist n -dimensional shift spaces if the derivatives of their generating functions are homeomorphisms of $\mathbb{R}$ (Proposition 1).

In contrast to a shift space the set of all images of the system of curves arising by shifting the argument from a Grünwald curve $\mathcal{C}$ under the translation group of $\mathbb{R}^n$ is a system of geodesics with respect to a natural affine connection if and only if $\mathcal{C}$ is a curve corresponding to parabolas in a suitable coordinate system (Theorem 2). Moreover, ∇ is metrizable and for $n = 2$ we get the metric tensor (4.4) of [13].

2. Grünwald shift spaces

An n -dimensional line space $\mathcal{S} = (\mathbb{R}^n, \mathcal{L})$, $n \geq 2$, is an incidence geometry such that the point set is the Euclidean space $\mathbb{R}^n$, the set $\mathcal{L}$ of lines consists of closed subsets of $\mathbb{R}^n$ homeomorphic to $\mathbb{R}$ and any two different points are incident with precisely one line.

We call an n -dimensional line space $\mathcal{S}$ an n -dimensional shift space if there exist continuous functions $f_i^{(k)} : \mathbb{R} \rightarrow \mathbb{R}$, $k = 1, \dots, n-1$, $i = k+1, \dots, n$, such that

$$\begin{aligned} \ell_{(u_1, \dots, u_n, v_{k+2}, \dots, v_n)}^{(k)} = \{ & (u_1, \dots, u_{k-1}, t + u_k, f_{k+1}^{(k)}(t) + u_{k+1}, f_{k+2}^{(k)}(t + v_{k+2}) \\ & + u_{k+2}, \dots, f_n^{(k)}(t + v_n) + u_n), t \in \mathbb{R} \}, \quad (1) \end{aligned}$$

where $u_1, \dots, u_n, v_{k+2}, \dots, v_n \in \mathbb{R}$,
 and $\{(u_1, \dots, u_{n-1}, t), t \in \mathbb{R}\}$ with $u_1, \dots, u_{n-1} \in \mathbb{R}$

form the set of lines for the line space $S(f_i^{(k)})$. The functions $f_i^{(k)}$ we will call *generating functions* of the shift space $S(f_i^{(k)})$.

Clearly, the group T of translations of $\mathbb{R}^n$ is a group of collineations of the shift spaces $S(f_i^{(k)})$.

We call an n -dimensional line space S , respectively n -dimensional shift space $S(f_i^{(k)})$, *differentiable* if the lines of S , respectively of $S(f_i^{(k)})$, are two times differentiable curves.

Shift spaces of the following proposition give for $n = 2$ Grünwald planes if their lines are geodesics with respect to an affine connection (cf. [13, § 4]). For this reason we call the shift spaces of the following proposition *Grünwald shift spaces*.

Proposition 1. *Let $f_i^{(k)} : \mathbb{R} \rightarrow \mathbb{R}$, $k = 1, \dots, n-1$, $i = k+1, \dots, n$, be differentiable functions such that the derivatives $f_i^{(k) \prime}$ are homeomorphisms of $\mathbb{R}$ for all $2 \leq i \leq n$. Then the functions $f_i^{(k)}$ are generating functions for an n -dimensional shift space $S(f_i^{(k)})$.*

PROOF. The lines of $S(f_i^{(k)})$ are the sets of form (1). Let

$$a = (a_1, a_2, \dots, a_n) \quad \text{and} \quad b = (b_1, b_2, \dots, b_n)$$

be two different points of $\mathbb{R}^n$.

Let $a_r = b_r$ for $r \leq k-1 < n-1$ and $a_k \neq b_k$. For a line through a and b we have $u_p = a_p = b_p$, $p = 1, \dots, k-1$. Moreover, the coordinates $a_k, b_k, a_{k+1}, b_{k+1}$ satisfy the following system of equations:

$$\begin{aligned} a_k &= t_a + u_k, & a_{k+1} &= f_{k+1}^{(k)}(t_a) + u_{k+1}; \\ b_k &= t_b + u_k, & b_{k+1} &= f_{k+1}^{(k)}(t_b) + u_{k+1}. \end{aligned} \tag{2}$$

Since the derivative of the function $f_p^{(k)}$, $p = k+1, \dots, n$, is a homeomorphism of $\mathbb{R}$, the function

$$t \longmapsto f_p^{(k)}(t+d) - f_p^{(k)}(t), \tag{3}$$

is a homeomorphism of $\mathbb{R}$ for any fixed $d \in \mathbb{R} \setminus \{0\}$ (cf. [15, § 3, p. 161]).

Now, from (2) we obtain $t_b = t_a - (a_k - b_k)$ and

$$a_{k+1} = f_{k+1}^{(k)}(t_a) + u_{k+1}; \quad b_{k+1} = f_{k+1}^{(k)}(t_a + (b_k - a_k)) + u_{k+1}.$$

This yields

$$a_{k+1} - b_{k+1} = f_{k+1}^{(k)}(t_a) - f_{k+1}^{(k)}(t_a + (b_k - a_k)).$$

Because $a_k \neq b_k$ relation (3) gives that there exists precisely one solution t_a of the last equation. Then we have $u_k = a_k - t_a$, $t_b = b_k - u_k$, and $u_{k+1} = a_{k+1} - f(t_a)$.

For $p = k + 2, \dots, n$ the coordinates a_p and b_p fulfill the following system of equations

$$a_p = f_p^{(k)}(t_a + v_p) + u_p; \quad b_p = f_p^{(k)}(t_b + v_p) + u_p.$$

Since the function $f_p^{(k)}$ satisfy (3) this system has precisely one solution u_p, v_p .

If $a_r = b_r$ for $r \leq n - 1$, and $a_n \neq b_n$ then the line $\{(a_1, \dots, a_{n-1}, t), t \in \mathbb{R}\}$ is the unique line joining a and b , and the proposition is proved. $\square$

3. Riccati and Abelian differential equations

For a later use we consider the special Riccati differential equations with unknown function $y = y(x)$:

$$y' + a_1 y^2 + a_2 y + a_3 = 0, \tag{4}$$

where a_i are constants (cf. [9, A4.9] or [10, pp. 33 and 41]).

(4.1) If $a_i = 0$ for $i \in \{1, 2\}$, then $y = -a_3 x + c$ for $c \in \mathbb{R}$.

(4.2) If $a_1 = 0$ and $a_2 \neq 0$, then $y = -\frac{a_3}{a_2} + ce^{-a_2 x}$ for $c \in \mathbb{R}$.

(4.3) If $a_1 \neq 0$ and $a_2^2 = 4a_1 a_3$, then we have

$$y = -\frac{a_2}{2a_1} + \frac{c_1}{c_1 a_1 x + c_2} \quad \text{with } c_1, c_2 \in \mathbb{R} \quad \text{and } (c_1, c_2) \neq (0, 0).$$

(4.4) If $a_1 \neq 0$ and $\lambda^2 = 4a_1 a_3 - a_2^2 > 0$, then

$$y = -\frac{a_2}{2a_1} + \frac{\lambda}{2a_1} \cotan \frac{\lambda}{2}(x + c) \quad \text{with } c \in \mathbb{R}.$$

(4.5) If $a_1 \neq 0$ and $\lambda^2 = a_2^2 - 4a_1 a_3 > 0$, then

$$y = -\frac{a_2}{2a_1} + \frac{\lambda}{2a_1} \frac{c_1 e^{\frac{\lambda}{2}x} - c_2 e^{-\frac{\lambda}{2}x}}{c_1 e^{\frac{\lambda}{2}x} + c_2 e^{-\frac{\lambda}{2}x}} \quad \text{with } c_1, c_2 \in \mathbb{R} \quad \text{and } (c_1, c_2) \neq (0, 0).$$

Also in (4.3) and (4.5) the solution depends only on one parameter which is defined on the projective line.

Lemma 1. *If f is a differentiable function such that its derivative f' is a homeomorphism of $\mathbb{R}$ and a solution of an equation (4), then f has the form $f = -1/2 a_3 x^2 + cx + d$ with $a_3 \neq 0$, $c, d \in \mathbb{R}$.*

PROOF. Solutions (4.1) with $a_3 = 0$ and (4.2) are excluded since in this case f' is not a homeomorphism. Solutions (4.1) with $a_3 \neq 0$ give the functions in the assertion.

Since in case of solutions (4.3) and (4.4) the function f' is not a homeomorphism of $\mathbb{R}$ we have to consider solutions (4.5). But also in this case the function f' is not a homeomorphism of $\mathbb{R}$ since we have $\lim_{x \rightarrow \pm\infty} f' = -\frac{a_2 \pm \lambda}{2a_1}$. $\square$

We consider Abelian differential equations

$$y' = \alpha + \beta y + \gamma y^2 + \varepsilon y^3 \quad \text{with} \quad \varepsilon \neq 0, \quad (5)$$

where $\alpha, \beta, \gamma, \varepsilon \in \mathbb{R}$, and we are interested in real functions f such that $f' = y$ is a homeomorphism of $\mathbb{R}$.

To differential equation (5) is associated the cubic algebraic equation

$$\alpha + \beta y + \gamma y^2 + \varepsilon y^3 = 0. \quad (6)$$

Because $\varepsilon \neq 0$, the cubic equation (6) has a real solution $y = y_1$ and hence equation (5) has a solution $y(t) = y_1$ for all $t \in \mathbb{R}$. According to the existence and uniqueness Theorem applied to (5), any other solution $f' = y$ of equation (5) satisfies either $y(t) > y_1$ or $y(t) < y_1$ for all $t \in \mathbb{R}$. Hence it follows

Lemma 2. *There exists no real function f with $f' = y$ satisfying (5) such that f' is a homeomorphism of the real line $\mathbb{R}$.*

4. Affine connections

Since we apply results of differential geometry only for the n -dimensional space $\mathbb{R}^n$ there exist global coordinates and the components Γ_{ij}^h , $h, i, j \in \{1, 2, \dots, n\}$, of any affine connection ∇ can be written in a unique way in these coordinates.

An affine connection ∇ is called symmetric if $\nabla_X Y = \nabla_Y X - [X, Y]$, where $[X, Y]$ is the Lie bracket, i.e. if for its components Γ_{ij}^h one has $\Gamma_{ij}^h = \Gamma_{ji}^h$ for all $h, i, j \in \{1, 2, \dots, n\}$.

By a *geodesic* of ∇ we mean a piecewise C^2 -curve $\gamma: I \rightarrow \mathbb{R}^n$ satisfying $\nabla_{\dot{\gamma}} \dot{\gamma} = \varrho \cdot \dot{\gamma}$, where $\varrho: I \rightarrow \mathbb{R}$ is a continuous function, and $I \subset \mathbb{R}$ is an open interval (cf. [4, p. 3], [14, p. 122]).

Using the components of ∇ the system of differential equations for geodesics has the form (cf. [14, p. 144])

$$\ddot{\gamma}^h + \sum_{i,j=1}^n \Gamma_{ij}^h \dot{\gamma}^i \dot{\gamma}^j = \varrho(t) \dot{\gamma}^h, \quad h \in \{1, 2, \dots, n\}. \quad (7)$$

From this it follows that the geodesics depend only on the symmetric part of the connection ∇ . Hence we will always assume that ∇ is symmetric.

Let $\mathfrak{g}$ be a Lie algebra of a group G of diffeomorphisms and let $\nabla = \{\Gamma_{ij}^h\}$ be an affine connection. The Lie derivative $\mathcal{L}_\xi \nabla$ along an element $\xi (\neq 0) \in \mathfrak{g}$ is given with respect to components of ∇ by

$$\mathcal{L}_\xi \Gamma_{ij}^h \equiv \frac{\partial^2 \xi^h}{\partial x_i \partial x_j} + \sum_{\alpha=1}^n \left(\xi^\alpha \frac{\partial \Gamma_{ij}^h}{\partial x_\alpha} - \frac{\partial \xi^h}{\partial x_\alpha} \Gamma_{ij}^\alpha + \frac{\partial \xi^\alpha}{\partial x_i} \Gamma_{\alpha j}^h + \frac{\partial \xi^\alpha}{\partial x_j} \Gamma_{\alpha i}^h \right),$$

where $h, i, \dots = 1, 2, \dots, n$.

The group G preserves geodesics with respect to ∇ if and only if

$$\mathcal{L}_\xi \Gamma_{ij}^h = \delta_i^h \psi_j + \delta_j^h \psi_i, \quad (8)$$

where δ_i^h is the Kronecker symbol and ψ_i are differentiable functions [11], [12, p. 143], [18].

The group G consists of affine mappings with respect to ∇ precisely if $\mathcal{L}_\xi \Gamma_{ij}^h = 0$ or, equivalently, if and only if ψ_i vanishes. Moreover, if $\mathbb{R}^n$ is a (pseudo-) Riemannian space with respect to the metric tensor g , then the Lie group G is a group of isometries precisely if $\mathcal{L}_\xi g = 0$ (cf. [18, p. 43], [12, p. 100]).

Proposition 2. *Let S be a system of geodesics with respect to an affine connection ∇ . If the translation group T of $\mathbb{R}^n$ consists of geodesic maps for S , then the affine connection ∇ may be chosen in such a way that the components Γ_{ij}^h are constant. Moreover, the components $\Gamma_{\sigma\sigma}^\sigma$, $\sigma = 1, \dots, n$, are zero.*

PROOF. Since T consists of geodesic maps for the Lie derivative $\mathcal{L}_\xi \Gamma_{ij}^h$ along any element $\xi \neq 0$ of the Lie algebra of T one has (8). Taking in particular $\xi = (\delta_\sigma^h)_{h=1}^n$ one obtains

$$\mathcal{L}_\xi \Gamma_{ij}^h \equiv \frac{\partial \Gamma_{ij}^h}{\partial x_\sigma} = \delta_i^h \psi_j + \delta_j^h \psi_i.$$

Integrating these equations for any $\sigma = 1, \dots, n$, we get

$$\Gamma_{ij}^h = \overset{\circ}{\Gamma}_{ij}^h + \delta_i^h \Psi_j + \delta_j^h \Psi_i,$$

where $\overset{\circ}{\Gamma}_{ij}^h$ are constants, and $\Psi_j(x)$ are suitable differentiable functions.

If with respect to the affine connection ∇ having the components Γ_{ij}^h the system S consists of geodesics, then the same holds for any connection with the components $\bar{\Gamma}_{ij}^h$ satisfying the equations $\bar{\Gamma}_{ij}^h = \Gamma_{ij}^h + \delta_i^h \bar{\psi}_j + \delta_j^h \bar{\psi}_i$, where $\bar{\psi}_i$ are differentiable functions ([11], [12], [18]). Choosing the functions $\bar{\psi}_i$ in such a way that $\bar{\psi}_i = -\Psi_i - \frac{1}{2} \overset{\circ}{\Gamma}_{ii}^i$ we see that then $\bar{\Gamma}_{ij}^h$ are constants, and $\bar{\Gamma}_{\sigma\sigma}^\sigma = 0$, $\sigma = 1, \dots, n$. $\square$

As representatives of affine connections for which the lines of a shift space S are geodesics we will take henceforth affine connections ∇° having constant components such that $\bar{\Gamma}_{\sigma\sigma}^\sigma = 0$, $\sigma = 1, \dots, n$. We shall call such connections *natural connections of S* . With respect to a natural connection ∇° the translation group of $\mathbb{R}^n$ consists of affine transformations of S . Namely, for $\xi = (\delta_\sigma^h)_{h=1}^n$ one has $\mathcal{L}_\xi \Gamma_{ij}^h \equiv \frac{\partial}{\partial x_\sigma} \Gamma_{ij}^h = 0$.

If a connection ∇ has the components Γ_{ij}^h , $h, i, j \in \{1, \dots, n\}$, the components R_{ijk}^h , $h, i, j, k \in \{1, \dots, n\}$ of the curvature tensor R of ∇ are given by (cf. [4, p. 8], [16, p. 27])

$$R_{ijk}^h = \frac{\partial}{\partial x_j} \Gamma_{ik}^h - \frac{\partial}{\partial x_k} \Gamma_{ij}^h + \sum_{\alpha=1}^n (\Gamma_{ik}^\alpha \Gamma_{\alpha j}^h - \Gamma_{ij}^\alpha \Gamma_{\alpha k}^h). \quad (9)$$

The Ricci tensor belonging to R has components $R_{ij} = \sum_{\alpha=1}^n R_{i\alpha j}^\alpha$.

The curvature tensor R of ∇ is often called the Riemannian tensor of ∇ .

In particular, ∇ is the *Levi-Civita connection* of a (pseudo-) Riemannian space with the metric $g = (g_{ij})$ if $\nabla g = 0$, i.e.

$$\frac{\partial}{\partial x_k} g_{ij} = \sum_{\alpha=1}^n (g_{i\alpha} \Gamma_{jk}^\alpha + g_{j\alpha} \Gamma_{ik}^\alpha), \quad (10)$$

where the components Γ_{ij}^h (called Christoffel symbols) are given by

$$\Gamma_{ij}^h = \frac{1}{2} \sum_{\alpha=1}^n g^{h\alpha} \left(\frac{\partial}{\partial x_i} g_{j\alpha} + \frac{\partial}{\partial x_j} g_{i\alpha} - \frac{\partial}{\partial x_\alpha} g_{ij} \right); \quad (11)$$

thereby $(g^{h\alpha})$ denotes the inverse matrix of (g_{ij}) . For g then there exists a unique symmetric affine connection ∇ such $\nabla g = 0$.

The integrability conditions of (10) have the following form [4, p. 79]:

$$\sum_{\alpha=1}^n (g_{i\alpha} R_{jkl}^\alpha + g_{j\alpha} R_{ikl}^\alpha) = 0. \quad (12)$$

5. Geometry of Grünwald shift spaces

Theorem 1. *Let $S(f_i^{(k)})$ be an n -dimensional Grünwald shift space. If the set of lines of $S(f_i^{(k)})$ forms the set of geodesics with respect to a natural connection ∇° , then $S(f_i^{(k)})$ is a Grünwald plane.*

PROOF. For a line

$$x_{(k)} = (u_1, \dots, u_{k-1}, t + u_k, f_{k+1}^{(k)}(t) + u_{k+1}, f_{k+2}^{(k)}(t + v_{k+2}) + u_{k+2}, \dots, f_n^{(k)}(t + v_n) + u_n), \quad t \in \mathbb{R},$$

one has

$$\begin{aligned} \dot{x}_{(k)} &= (0, \dots, 0, 1, f_{k+1}^{(k)'}(t), f_{k+2}^{(k)'}(t + v_{k+2}), \dots, f_n^{(k)'}(t + v_n)), \\ \ddot{x}_{(k)} &= (0, \dots, 0, 0, f_{k+1}^{(k)''}(t), f_{k+2}^{(k)''}(t + v_{k+2}), \dots, f_n^{(k)''}(t + v_n)). \end{aligned}$$

This line is a geodesic if and only if relation (7) holds. We put in this relation $t_{k+1} \equiv t$ and $t_\lambda \equiv t + v_\lambda$ for $\lambda > k + 1$.

For $h = k = n - 1$ one has $\varrho(t_n) = 2\Gamma_{n-1\ n}^{n-1} f_n^{(n-1)'}(t_n) + \Gamma_{nn}^{n-1} (f_n^{(n-1)'}(t_n))^2$ and for $h = n, k = n - 1$ using $\Gamma_{nn}^n = 0$ (Proposition 2) we get

$$f_n^{(n-1)''} + \Gamma_{n-1\ n-1}^n + 2\Gamma_{n-1\ n}^n f_\sigma^{(k)'} = \varrho(t_n) f_n^{(n-1)'}. \quad (13)$$

Substituting $\varrho(t_n)$ into (13) we obtain

$$f_n^{(n-1)''} = -\Gamma_{n-1\ n-1}^n - 2\Gamma_{n-1\ n}^n f_\sigma^{(k)'} + 2\Gamma_{n-1\ n}^{n-1} (f_n^{(n-1)'})^2 + \Gamma_{nn}^{n-1} (f_n^{(n-1)'})^3. \quad (14)$$

This equation with constant coefficients is an Abelian differential equation with respect to $f_n^{(n-1)'}(t_n)$. By Lemma 1 and 2 it follows that $f_n^{(n-1)'}(t_n) = \alpha t_n^2 + \beta t_n + \gamma$ with constants $\alpha \neq 0$, β and γ . Putting this in (14) we get

$$\Gamma_{nn}^{n-1} = \Gamma_{n-1\ n}^{n-1} = \Gamma_{n-1\ n}^n = 0, \quad \text{but} \quad \Gamma_{n-1\ n-1}^n = -2\alpha \neq 0. \quad (15)$$

If $n > 2$ then from (7) for $h = k = n - 2$ one has

$$\begin{aligned} \varrho(t_{n-1}) &= 2\Gamma_{n-2\ n-1}^{n-2} f_{n-1}^{(n-2)'}(t_{n-1}) + 2\Gamma_{n-2\ n}^{n-2} f_n^{(n-2)'}(t_n) + \Gamma_{n-1\ n-1}^{n-2} (f_{n-1}^{(n-2)'}(t_{n-1}))^2 \\ &\quad + 2\Gamma_{n-1\ n}^{n-2} f_{n-1}^{(n-2)'}(t_{n-1}) f_n^{(n-2)'}(t_n) + \Gamma_{nn}^{n-2} (f_n^{(n-2)'}(t_n))^2 \end{aligned}$$

and for $h = n, k = n - 2$ using $\Gamma_{nn}^n = \Gamma_{n-1\ n}^n = 0$ we get

$$\begin{aligned} f_h^{(n-2)''}(t_n) + \Gamma_{n-2\ n-2}^n + 2\Gamma_{n-2\ n-1}^n f_{n-1}^{(n-2)'}(t_{n-1}) + 2\Gamma_{n-2\ n}^n f_n^{(n-2)'}(t_n) \\ + \Gamma_{n-1\ n-1}^n (f_{n-1}^{(n-2)'}(t_{n-1}))^2 = \varrho(t_{n-1}) f_n^{(n-2)'}(t_n). \end{aligned}$$

Substituting into this $\varrho(t_{n-1})$ we obtain

$$\begin{aligned} & (f_{n-1}^{(n-2)'}(t_{n-1}))^2 \cdot (\Gamma_{n-1\ n-1}^n - \Gamma_{n-1\ n-1}^{n-2} f_n^{(n-2)'}(t_n)) \\ & + f_{n-1}^{(n-2)'}(t_{n-1}) \cdot A(t_n) + B(t_n) = 0, \end{aligned}$$

where $A(t_n)$ and $B(t_n)$ are functions of the variable t_n . Since the variables t_{n-1} and t_n are independent and $f_{n-1}^{(n-2)'} \neq 0$ the coefficient functions $A(t_n)$ and $B(t_n)$ vanish and

$$\Gamma_{n-1\ n-1}^n - \Gamma_{n-1\ n-1}^{n-2} f_n^{(n-2)'}(t_n) = 0. \quad (16)$$

From (16) it follows $\Gamma_{n-1\ n-1}^{n-2} = 0$ and $\Gamma_{n-1\ n-1}^n = 0$. This contradicts relations (15). Hence n must be 2 and $S(f_i^{(k)})$ is a Grünwald plane (cf. [13]). $\square$

Remark. If $n = 2$ then the proof of Theorem 1 yields that $\Gamma_{11}^2 = -2\alpha \neq 0$ and all other components are zero. Hence this shift space is the Grünwald plane M_α having a metric tensor g with corresponds to the Levi-Civita connection ∇ of the form (4.4) in [13].

6. Translation shell of a Grünwald curve

Let $\mathcal{C}$ be a curve homeomorphic to $\mathbb{R}$ which is a closed subset of in $\mathbb{R}^n$, $n \geq 2$. The translation shell $\mathcal{C}^T$ of $\mathcal{C}$ is the set of all images of $\mathcal{C}$ under the translation group T of $\mathbb{R}^n$. We consider a curve of the form

$$\mathcal{C} = \{(t, f_2(t), f_3(t), \dots, f_n(t)), t \in \mathbb{R}\}, \quad (17)$$

where $f_i(t)$ are two times differentiable functions such that the derivatives $f_i'(t)$ are homeomorphisms of $\mathbb{R}$ for all $i = 2, \dots, n$. The translation shell of $\mathcal{C}$ is the set

$$\mathcal{C}^T = \{(t + u_1, f_2(t) + u_2, f_3(t) + u_3, \dots, f_n(t) + u_n), t \in \mathbb{R}\},$$

where $u_1, \dots, u_n \in \mathbb{R}$.

The extended translation shell $\hat{\mathcal{C}}^T$ is the set

$$\hat{\mathcal{C}}^T = \{(t + u_1, f_2(t) + u_2, f_3(t + v_3) + u_3, \dots, f_n(t + v_n) + u_n), t \in \mathbb{R}\},$$

$$\text{where } u_1, \dots, u_n, v_3, \dots, v_n \in \mathbb{R}. \quad (18)$$

We search for affine connections ∇ for which the extended translation shell $\hat{\mathcal{C}}^T$ or the translation shell $\mathcal{C}^T$ consists of geodesics with respect to ∇ . If $n = 2$ then the extended shell $\hat{\mathcal{C}}^T$ is a Grünwald plane if we adjoin to $\hat{\mathcal{C}}^T$ the lines $\{(u, t); t \in \mathbb{R}\}$, $u \in \mathbb{R}$. For this reason we call such curves $\mathcal{C}$ *Grünwald curves*.

Theorem 2. For a Grünwald curve $\mathcal{C}$ the extended translation shell $\hat{\mathcal{C}}^T$ consists of geodesics with respect to a natural affine connection ∇° with components Γ_{ij}^h if and only if the functions $f_i(t)$ may be chosen as $f_i(t) = -\frac{1}{2}\Gamma_{11}^i t^2 + \beta_i t$ with $\Gamma_{11}^i \neq 0$, $\beta_i \in \mathbb{R}$, $i = 2, 3, \dots, n$, whereas all other components of ∇° are zero.

PROOF. Let $x(t)$ be a curve in (18). Then we have

$$\begin{aligned}\dot{x}(t) &= (1, f'_2(t), f'_3(t + v_3), \dots, f'_n(t + v_n)), \\ \ddot{x}(t) &= (0, f''_2(t), f''_3(t + v_3), \dots, f''_n(t + v_n)).\end{aligned}$$

The curve $x(t)$ is a geodesic if and only if relation (7) holds. We put in this relation $t_2 \equiv t$, and $t_\lambda \equiv t + v_\lambda$ for $\lambda > 2$.

For $h = 1$ one has

$$\varrho(t_2) = 2 \sum_{\sigma=2}^n \Gamma_{1\sigma}^1 f'_\sigma(t_\sigma) + \sum_{\sigma,\tau=2}^n \Gamma_{\sigma\tau}^1 f'_\sigma(t_\sigma) f'_\tau(t_\tau),$$

and for $h > 1$ we get

$$f''_h(t_h) + \Gamma_{11}^h + 2 \sum_{\sigma=2}^n \Gamma_{1\sigma}^h f'_\sigma(t_\sigma) + \sum_{\sigma,\tau=2}^n \Gamma_{\sigma\tau}^h f'_\sigma(t_\sigma) f'_\tau(t_\tau) = \varrho(t_2) f'_h(t_h). \quad (19)$$

Putting $\varrho(t_2)$ into (19) and fixing all variables t_σ different from t_h we obtain with respect to function $f'_h(t_h)$ an Abelian differential equation with constant coefficients since ∇° is a natural affine connection. By Lemmas 1 and 2 it follows $f_h(t_h) = \alpha_h t_h^2 + \beta_h t_h + \gamma_h$, with constants $\alpha_h \neq 0$, β_h and γ_h .

Substituting $\varrho(t_2)$ and $f_h(t_h)$ in (19) we obtain

$$\begin{aligned}\alpha_h + \Gamma_{11}^h + 2 \sum_{\sigma=2}^n (\Gamma_{1\sigma}^h - \Gamma_{1\sigma}^1 (\alpha_h t_h + \beta_h)) \cdot (\alpha_\sigma t_\sigma + \beta_\sigma) \\ + \sum_{\sigma,\tau=2}^n (\Gamma_{\sigma\tau}^h - \Gamma_{\sigma\tau}^1 (\alpha_h t_h + \beta_h)) (\alpha_\sigma t_\sigma + \beta_\sigma) (\alpha_\tau t_\tau + \beta_\tau) \equiv 0,\end{aligned} \quad (20)$$

where $h = 2, 3, \dots, n$ and $t_2, t_3, \dots, t_n$ are independent variables.

Since (20) is a cubic polynomial the coefficients at monomials are zero. This yields $\alpha_h = -\Gamma_{11}^h \neq 0$ and all other components of ∇° are zero. $\square$

Remark. The metric tensor g with components

$$g_{11} = 1 + (x_1)^2 \cdot \sum_{\alpha=2}^n (\Gamma_{11}^\alpha)^2, \quad g_{1b} = \Gamma_{11}^b \cdot x_1, \quad g_{ab} = \delta_{ab}, \quad a, b \neq 1,$$

where δ_{ab} is the Kronecker symbol, determines (see (11)) the Levi-Civita connection ∇ with components as in Theorem 2 (having $\Gamma_{11}^h, h = 2, \dots, n$, as the only non zero components). Moreover, the Riemannian tensor vanishes, hence the space $(\mathbb{R}^n, \Gamma_{ij}^h)$ is locally Euclidean (cf. [14]).

If we strength the hypothesis on the Grünwald curve $\mathcal{C}$ we obtain the same system of functions $f_i(t)$ as in Theorem 2, but for a given system of functions $f_i(t)$ there are more natural affine connections having the curves of the translation shell $\mathcal{C}^T$ as geodesics.

Theorem 3. *Let $\mathcal{C}$ be a Grünwald curve such that the derivatives of all its functions $f_i(t)$ satisfy Abelian differential equations with constant coefficients. Then the translation shell $\mathcal{C}^T$ of $\mathcal{C}$ consists of geodesics with respect to a natural affine connection ∇° with components Γ_{ij}^h if and only if the functions $f_i(t)$ may be chosen as $f_i(t) = -\frac{1}{2}\Gamma_{11}^i t^2 + \beta_i t$ with $\Gamma_{11}^i \neq 0, \beta_i \in \mathbb{R}, i = 2, 3, \dots, n$, whereas all other components of ∇° are zero with exception of $\Gamma_{h\sigma}^h = \Gamma_{\sigma h}^h = \Gamma_{1\sigma}^1 = \Gamma_{\sigma 1}^1$ for $h > 1, \sigma > 1$.*

PROOF. Let $x(t)$ be a curve in $\mathcal{C}^T$. Since $f'_i(t), i = 2, \dots, n$, satisfy Abelian differential equations with constant coefficients it follows from Lemma 1 and 2 that $f_i(t) = \frac{1}{2}\alpha_i t^2 + \beta_i t + \gamma_i$ with constants $\alpha_i \neq 0, \beta_i$ and γ_i . Therefore for $x(t)$ we have

$$\dot{x}(t) = (1, \alpha_2 t + b_2, \alpha_3 t + b_3, \dots, \alpha_n t + b_n) \quad \text{and} \quad \ddot{x}_{(k)} = (0, \alpha_2, \alpha_3, \dots, \alpha_n).$$

The curve $x(t)$ is a geodesic if and only if relation (7) holds. For $h = 1$ in (7) one has

$$\varrho(t) = 2 \sum_{\sigma=2}^n \Gamma_{1\sigma}^1 (\alpha_\sigma t + \beta_\sigma) + \sum_{\sigma, \tau=2}^n \Gamma_{\sigma\tau}^1 (\alpha_\sigma t + \beta_\sigma) (\alpha_\tau t + \beta_\tau),$$

and for $h > 1$ we get

$$\begin{aligned} \alpha_h + \Gamma_{11}^h + 2 \sum_{\sigma=2}^n \Gamma_{1\sigma}^h (\alpha_\sigma t + \beta_\sigma) + \sum_{\sigma, \tau=2}^n \Gamma_{\sigma\tau}^h (\alpha_\sigma t + \beta_\sigma) (\alpha_\tau t + \beta_\tau) \\ = \varrho(t)(\alpha_h t + \beta_h). \end{aligned} \quad (21)$$

Putting $\varrho(t)$ into (21) we obtain a polynomial which is identically zero. It follows immediately that $\Gamma_{\sigma\tau}^1 = 0$ for $\sigma > 1, \tau > 1$ and $\alpha_h = -\Gamma_{11}^h$ as well as $\Gamma_{\sigma 1}^h = \Gamma_{1\sigma}^h = 0$ for $h > 1, \sigma > 1$. Finally, we have

$$\sum_{\sigma, \tau=2}^n \Gamma_{\sigma\tau}^h (\alpha_\sigma t + \beta_\sigma) (\alpha_\tau t + \beta_\tau) - 2(\alpha_h t + \beta_h) \cdot \sum_{\sigma=2}^n \Gamma_{1\sigma}^1 (\alpha_\sigma t + \beta_\sigma) = 0.$$

From this relation it follows that $\Gamma_{h\sigma}^h = \Gamma_{\sigma h}^h = \Gamma_{1\sigma}^1 = \Gamma_{\sigma 1}^1$ for $h > 1, \sigma > 1$ and all other components $\Gamma_{\sigma\tau}^h$ vanish. $\square$

References

- [1] D. BETTEN, Eine Charakterisierung des Reellen Halbraumes, *Rend. Circ. Mat. Palermo, II. Ser.* **32** (1983), 145–150.
- [2] D. BETTEN, Einige Klassen topologischer 3-Räume, *Result. Math.* **12** (1987), 37–61.
- [3] D. BETTEN, Topological shift spaces, *Adv. Geom.* **5** (2005), 107–118.
- [4] L. P. EISENHART, Non-Riemannian geometry, Vol. 8, *American Mathematical Society Colloquium Publications*, 2000.
- [5] G. GERLICH, Stable projective planes with Riemannian metrics, *Arch. Math.* **79** (2002), 317–320.
- [6] G. GERLICH, Topological affine planes with affine connections, *Adv. Geom.* **5** (2005), 265–278.
- [7] G. GERLICH, Cones for the Moulton planes, *J. Geom.* **88**, no. 1–2 (2008), 30–40.
- [8] J. GRÜNWARD, Über duale Zahlen und ihre Anwendung in der Geometrie, *Monatsh. Math. Phys.* **17** (1906), 81–136.
- [9] E. KAMKE, Differentialgleichungen, Lösungsmethoden und Lösungen, *Akademische Verlagsgesellschaft, Leipzig*, 1942.
- [10] E. KAMKE, Differentialgleichungen reeller Funktionen, *Akademische Verlagsgesellschaft, Leipzig*, 1956.
- [11] J. MIKEŠ, Geodesic mappings of affine-connected and Riemannian spaces, *J. Math. Sci.* **78** (1996), 311–333.
- [12] J. MIKEŠ, A. VANŽUROVÁ and I. HINTERLEITNER, Geodesic mappings and some generalizations, *Palacky University Press, Olomouc*, 2009.
- [13] J. MIKEŠ and K. STRAMBACH, Differentiable structures on elementary geometries, *Result. Math.* **53**, no. 1–2 (2009), 153–172.
- [14] A. P. NORDEN, Spaces of an Affine Connection, *Nauka, Moscow*, 1976 (in Russian).
- [15] H. SALZMANN, D. BETTEN, T. GRUNDHÖFER, H. HÄHL, R. LÖWEN and M. STROPPEL, Compact projective planes, *de Gruyter*, 1995.
- [16] N. S. SINYUKOV, Geodesic mappings of Riemannian spaces, *Moscow, Nauka*, 1979 (in Russian).
- [17] L. A. SKORNJAKOV, Systems of curves on a surface, *Trudy Moskov. Mat. Obsc.* **6** (1957), 135–164 (in Russian).
- [18] K. YANO and S. BOCHNER, Curvature and Betti numbers, *Annals of Mathematics Studies* **32**, Princeton: N. J.: Princeton University Press, 1953.

JOSEF MIKEŠ
 DEPARTMENT OF ALGEBRA AND GEOMETRY
 FACULTY OF SCIENCE
 PALACKY UNIVERSITY
 17. LISTOPADU 12
 77146 OLOMOUC
 CZECH REPUBLIC
E-mail: josef.mikes@upol.cz

KARL STRAMBACH
 DEPARTMENT MATHEMATIK
 DER UNIVERSITÄT ERLANGEN-NÜRNBERG
 CAUERSTR. 11,
 D-91052 ERLANGEN
 GERMANY
E-mail: strambach@mi.uni-erlangen.de

(Received April 11, 2012; revised November 29, 2012)