

Balancing numbers which are products of consecutive integers

By SZABOLCS TENGELY (Debrecen)

Abstract. In 1999 A. BEHERA and G. K. PANDA defined balancing numbers as follows. A positive integer n is called a balancing number if $1 + 2 + \cdots + (n - 1) = (n + 1) + (n + 2) + \cdots + (n + k)$ for some $k \in \mathbb{N}$. The sequence of balancing numbers is denoted by B_m for $m \in \mathbb{N}$. In this paper we show that the Diophantine equation $B_m = x(x + 1)(x + 2)(x + 3)(x + 4)$ has no solution with $m \geq 0$ and $x \in \mathbb{Z}$. We follow the ideas described in [13], that is we combine Baker’s method and the so-called Mordell–Weil sieve to obtain all solutions.

1. Introduction

A positive integer n is called a balancing number if

$$1 + 2 + \cdots + (n - 1) = (n + 1) + (n + 2) + \cdots + (n + k)$$

for some $k \in \mathbb{N}$. The sequence of balancing numbers is denoted by B_m for $m \in \mathbb{N}$. We note that usually the initial values $B_0 = 0$, $B_1 = 1$ of the balancing sequence are used. BEHERA and PANDA [3] proved many interesting results related to the sequence B_m . They showed that the balancing numbers fulfill the following recurrence relation

$$B_{m+1} = 6B_m - B_{m-1} \quad (m \geq 1)$$

where $B_0 = 0$ and $B_1 = 1$. Later several authors investigated balancing numbers and their various generalizations. In [21] LIPTAI proved that there are no Fibonacci balancing numbers and in [22] he showed that there are no Lucas balancing

Mathematics Subject Classification: Primary: 11D61; Secondary: 11Y50.

Key words and phrases: Diophantine equations.

Research supported in part by the OTKA grants K100339, NK104208 and PD75264 and János Bolyai Research Scholarship of the Hungarian Academy of Sciences.

numbers. He used a method by BAKER and DAVENPORT [2]. SZALAY [29] obtained the same results by using different techniques. In [25] PANDA introduced the sequence balancing numbers. Let $\{s_m\}_{m=1}^{\infty}$ be a sequence such that $s_m \in \mathbb{R}$. An element s_m of this sequence is called a sequence balancing number if

$$s_1 + s_2 + \cdots + s_{m-1} = s_{m+1} + s_{m+2} + \cdots + s_{m+k}$$

for some $k \in \mathbb{N}$. Further generalization in this direction is due to BÉRCZES, LIPTAI and PINK [4]. Now let a, b two non-negative coprime integers and recall the following definition of [19]. A positive integer $an + b$ is called (a, b) -type balancing number if

$$(a + b) + (2a + b) + \cdots + (a(n - 1) + b) = (a(n + 1) + b) + \cdots + (a(n + k) + b)$$

for some $k \in \mathbb{N}$. Denote by $B_m^{(a,b)}$ the m -th positive integer $an + b$ among the (a, b) -type balancing numbers. KOVÁCS, LIPTAI and OLAJOS [19] proved some general finiteness results concerning the equation

$$B_m^{(a,b)} = f(x),$$

where f is a monic polynomial with integral coefficients. They also resolved some related Diophantine equations. LIPTAI, LUCA, PINTÉR and SZALAY [23] introduced the concept of (k, l) -power numerical center as follows. Let y, k, l be fixed positive integers with $y \geq 2$. A positive integer x with $x \leq y - 2$ is called a (k, l) -power numerical center for y if

$$1^k + \cdots + (x - 1)^k = (x + 1)^l + \cdots + (y - 1)^l.$$

The authors of [23] obtained certain effective and ineffective finiteness results for (k, l) -power numerical centers.

For positive integers k, x let

$$\Pi_k(x) = x(x + 1) \cdots (x + k - 1).$$

That is, $\Pi_k(x)$ is a polynomial in x of degree k . In [19] it was proved that the equation

$$B_m = \Pi_k(x)$$

for fixed $k \geq 2$ has only finitely many solutions and for $k \in \{2, 3, 4\}$ all solutions were determined. We note that in [19] the “small” solutions of the above equation with $k \in \{6, 8\}$ were also computed.

In this paper we deal with the case $k = 5$. That is we consider the equation

$$B_m = x(x + 1)(x + 2)(x + 3)(x + 4).$$

We prove the following theorem.

Theorem 1. *The Diophantine equation*

$$B_m = x(x+1)(x+2)(x+3)(x+4) \quad m \geq 1, x \in \mathbb{Z}$$

has no solution.

2. Auxiliary results

Consider the hyperelliptic curve

$$\mathcal{C}: y^2 = F(x) := x^5 + b_4x^4 + b_3x^3 + b_2x^2 + b_1x + b_0, \quad (1)$$

where $b_i \in \mathbb{Z}$. Let α be a root of F and $J(\mathbb{Q})$ be the Jacobian of the curve $\mathcal{C}$. We have that

$$x - \alpha = \kappa \xi^2$$

where $\kappa, \xi \in K = \mathbb{Q}(\alpha)$ and κ comes from a finite set. By knowing the Mordell–Weil group of the curve $\mathcal{C}$ it is possible to provide a method to compute such a finite set. To each coset representative $\sum_{i=1}^m (P_i - \infty)$ of $J(\mathbb{Q})/2J(\mathbb{Q})$ we associate

$$\kappa = \prod_{i=1}^m (\gamma_i - \alpha d_i^2),$$

where the set $\{P_1, \dots, P_m\}$ is stable under Galois action, all $y(P_i)$ are non-zero and $x(P_i) = \gamma_i/d_i^2$ where γ_i is an algebraic integer and $d_i \in \mathbb{Z}_{\geq 1}$. If P_i, P_j are conjugate, then we may suppose that $d_i = d_j$ and so γ_i, γ_j are conjugate. We have the following lemma (Lemma 3.1 in [13]).

Lemma 1. *Let $\mathcal{K}$ be a set of κ values associated as above to a complete set of coset representatives of $J(\mathbb{Q})/2J(\mathbb{Q})$. Then $\mathcal{K}$ is a finite subset of $\mathcal{O}_K$ and if (x, y) is an integral point on the curve (1) then $x - \alpha = \kappa \xi^2$ for some $\kappa \in \mathcal{K}$ and $\xi \in K$.*

As an application of his theory of lower bounds for linear forms in logarithms, BAKER [1] gave an explicit upper bound for the size of integral solutions of hyperelliptic curves. This result has been improved by many authors (see e.g. [5], [7], [10], [28] and [30]).

In [13] an improved completely explicit upper bound was proved by combining ideas from [10], [11], [12], [20], [24], [26], [30], [31]. Now we will state the theorem which gives the improved bound. We introduce some notation. Let K be a

number field of degree d and let r_K be its unit rank further R_K its regulator. For $\alpha \in K$ we denote by $h(\alpha)$ the logarithmic height of the element α . Let

$$\partial_K = \begin{cases} \frac{\log 2}{d} & \text{if } d = 1, 2, \\ \frac{1}{4} \left(\frac{\log \log d}{\log d} \right)^3 & \text{if } d \geq 3 \end{cases}$$

and

$$\partial'_K = \left(1 + \frac{\pi^2}{\partial_K^2} \right)^{1/2}.$$

Define the constants

$$\begin{aligned} c_1(K) &= \frac{(r_K!)^2}{2^{r_K-1} d^{r_K}}, & c_2(K) &= c_1(K) \left(\frac{d}{\partial_K} \right)^{r_K-1}, \\ c_3(K) &= c_1(K) \frac{d^{r_K}}{\partial_K}, & c_4(K) &= r_K d c_3(K), \\ c_5(K) &= \frac{r_K^{r_K+1}}{2 \partial_K^{r_K-1}}. \end{aligned}$$

Let

$$\partial_{L/K} = \max \left\{ [L : \mathbb{Q}], [K : \mathbb{Q}] \partial'_K, \frac{0.16[K : \mathbb{Q}]}{\partial_K} \right\},$$

where $K \subseteq L$ are number fields. Define

$$C(K, n) := 3 \cdot 30^{n+4} \cdot (n+1)^{5.5} d^2 (1 + \log d).$$

The following theorem will be used to get an upper bound for the size of the integral solutions of our equation. It is Theorem 3 in [13].

Theorem 2. *Let α be an algebraic integer of degree at least 3 and κ be an integer belonging to K . Denote by $\alpha_1, \alpha_2, \alpha_3$ distinct conjugates of α and by $\kappa_1, \kappa_2, \kappa_3$ the corresponding conjugates of κ . Let*

$$K_1 = \mathbb{Q}(\alpha_1, \alpha_2, \sqrt{\kappa_1 \kappa_2}), \quad K_2 = \mathbb{Q}(\alpha_1, \alpha_3, \sqrt{\kappa_1 \kappa_3}), \quad K_3 = \mathbb{Q}(\alpha_2, \alpha_3, \sqrt{\kappa_2 \kappa_3}),$$

and

$$L = \mathbb{Q}(\alpha_1, \alpha_2, \alpha_3, \sqrt{\kappa_1 \kappa_2}, \sqrt{\kappa_1 \kappa_3}).$$

In what follows R stands for an upper bound for the regulators of K_1, K_2 and K_3 and r denotes the maximum of the unit ranks of K_1, K_2, K_3 . Let

$$c_j^* = \max_{1 \leq i \leq 3} c_j(K_i), \quad N = \max_{1 \leq i, j \leq 3} \left| \text{Norm}_{\mathbb{Q}(\alpha_i, \alpha_j)/\mathbb{Q}}(\kappa_i(\alpha_i - \alpha_j)) \right|^2,$$

and

$$H^* = c_5^* R + \frac{\log N}{\min_{1 \leq i \leq 3} [K_i : \mathbb{Q}]} + h(\kappa).$$

Define

$$A_1^* = 2H^* \cdot C(L, 2r+1) \cdot (c_1^*)^2 \partial_{L/L} \cdot \left(\max_{1 \leq i \leq 3} \partial_{L/K_i} \right)^{2r} \cdot R^2,$$

and

$$A_2^* = 2H^* + A_1^* + A_1^* \log\{(2r+1) \cdot \max\{c_4^*, 1\}\}.$$

If $x \in \mathbb{Z} \setminus \{0\}$ satisfies $x - \alpha = \kappa \xi^2$ for some $\xi \in K$ then

$$\log|x| \leq 8A_1^* \log(4A_1^*) + 8A_2^* + H^* + 20 \log 2 + 13 h(\kappa) + 19 h(\alpha).$$

To obtain a lower bound for the possible unknown integer solutions we are going to use the so-called Mordell–Weil sieve. The Mordell–Weil sieve has been successfully applied to prove the non-existence of rational points on curves (see e.g. [8], [9], [17] and [27]).

Let $C/\mathbb{Q}$ be a smooth projective curve (in our case a hyperelliptic curve) of genus $g \geq 2$. Let J be its Jacobian. We assume the knowledge of some rational points on C , so let D be a fixed rational point on C and let j be the corresponding Abel–Jacobi map:

$$j : C \rightarrow J, \quad P \mapsto [P - D].$$

Let W be the image in J of the known rational points on C and $D_1, \dots, D_r$ generators for the free part of $J(\mathbb{Q})$. By using the Mordell–Weil sieve we are going to obtain a very large and smooth integer B such that

$$j(C(\mathbb{Q})) \subseteq W + BJ(\mathbb{Q}).$$

Let

$$\phi : \mathbb{Z}^r \rightarrow J(\mathbb{Q}), \quad \phi(a_1, \dots, a_r) = \sum a_i D_i,$$

so that the image of ϕ is the free part of $J(\mathbb{Q})$. The variant of the Mordell–Weil sieve explained in [13] provides a method to obtain a very long decreasing sequence of lattices in $\mathbb{Z}^r$

$$B\mathbb{Z}^r = L_0 \supsetneq L_1 \supsetneq L_2 \supsetneq \dots \supsetneq L_k$$

such that

$$j(C(\mathbb{Q})) \subset W + \phi(L_j)$$

for $j = 1, \dots, k$.

The next lemma [13, Lemma 12.1] gives a lower bound for the size of rational points whose image are not in the set W . Let h be the logarithmic height on J and $\hat{h}$ be the canonical height on J .

Lemma 2. *Let W be a finite subset of $J(\mathbb{Q})$ and L be a sublattice of $\mathbb{Z}^r$. Suppose that $j(C(\mathbb{Q})) \subset W + \phi(L)$. Let μ_1 be a lower bound for $h - \hat{h}$ and*

$$\mu_2 = \max \left\{ \sqrt{\hat{h}(w)} : w \in W \right\}.$$

Denote by M the height-pairing matrix for the Mordell–Weil basis $D_1, \dots, D_r$ and let $\lambda_1, \dots, \lambda_r$ be its eigenvalues. Let

$$\mu_3 = \min \left\{ \sqrt{\lambda_j} : j = 1, \dots, r \right\}$$

and $m(L)$ the Euclidean norm of the shortest non-zero vector of L . Then, for any $P \in C(\mathbb{Q})$, either $j(P) \in W$ or

$$h(j(P)) \geq (\mu_3 m(L) - \mu_2)^2 + \mu_1.$$

3. Proof of Theorem 1

It was shown by Liptai that the integers B_m satisfy the following equation

$$z^2 - 8y^2 = 1$$

for some integer z . So one has to determine all solution of the equation

$$z^2 = 8(x(x+1)(x+2)(x+3)(x+4))^2 + 1.$$

Rewrite the latter equation as follows

$$z^2 = 8(x^2 + 4x)^2(x^2 + 4x + 3)^2(x^2 + 4x + 4) + 1.$$

Let $X = 2x^2 + 8x$. We obtain that

$$\mathcal{C} : \quad Y^2 = X^2(X+6)^2(X+8) + 4, \quad (2)$$

where $Y = 2z$. It remains to find all integral points on $\mathcal{C}$. The rank of the Jacobian of $\mathcal{C}$ is 3, so classical Chabauty's method [14], [15], [16] cannot be applied. In this paper we combine Baker's method and the so-called Mordell–Weil sieve to obtain all integral solutions of equation (2).

Lemma 3. *The only integral solutions to the equation (2) are*

$$(0, \pm 2), (-6, \pm 2), (-8, \pm 2).$$

PROOF. Let $J(\mathbb{Q})$ be the Jacobian of the genus two curve (2). Using MAGMA [6] we determine a Mordell–Weil basis which is given by

$$\begin{aligned} D_1 &= (0, 2) - \infty, \\ D_2 &= (-6, 2) - \infty, \\ D_3 &= (\omega, -\omega - 10) + (\bar{\omega}, -\bar{\omega} - 10) - 2\infty, \end{aligned}$$

where ω is a root of the polynomial $x^2 + 7x + 4$. Let $f = x^2(x+6)^2(x+8) + 4$ and α be a root of f . We will choose for coset representatives of $J(\mathbb{Q})/2J(\mathbb{Q})$ the linear combinations $\sum_{i=1}^3 n_i D_i$, where $n_i \in \{0, 1\}$. We have

$$x - \alpha = \kappa \xi^2,$$

where κ belongs to a finite set (having 8 elements). This set can be constructed as described in Lemma 1. We apply Theorem 2 to get a large upper bound for $\log |x|$. A MAGMA code was written by GALLEGOS–RUIZ [18] to obtain bounds for such equations. We used the above MAGMA functions to compute an upper bound for $\log |x|$, the results are summarized in the following table

κ	bound for $\log x $
1	$4.17 \cdot 10^{204}$
$-\alpha$	$1.59 \cdot 10^{411}$
$-6 - \alpha$	$3.11 \cdot 10^{430}$
$4 + 7\alpha + \alpha^2$	$1.59 \cdot 10^{411}$
$-8 + 6\alpha + \alpha^2$	$3.11 \cdot 10^{430}$
$13 + 9\alpha + \alpha^2$	$1.59 \cdot 10^{411}$
$6\alpha + \alpha^2$	$3.11 \cdot 10^{430}$
$-10 + 5\alpha + \alpha^2$	$3.11 \cdot 10^{430}$

The set of known rational points on the curve (2) is $\{\infty, (0, \pm 2), (-6, \pm 2), (-8, \pm 2)\}$. Let W be the image of this set in $J(\mathbb{Q})$. Applying the Mordell–Weil sieve implemented by BRUIN and STOLL we obtain that

$$J(C(\mathbb{Q})) \subseteq W + BJ(\mathbb{Q}),$$

where

$$B = 2^6 \cdot 3^4 \cdot 5^3 \cdot 7^3 \cdot 11^2 \cdot 13^2 \cdot 17 \cdot 19^2 \cdot 29 \cdot 31 \cdot 41 \cdot 43 \cdot 47 \cdot 61 \cdot 67 \cdot 73 \cdot 79 \cdot 83 \cdot 89 \cdot 97 \cdot 107 \cdot 109 \cdot 113,$$

that is

$$B = 46247720065121846143591520774334300410472000.$$

Now we use an extension of the Mordell–Weil sieve due to SAMIR SIKSEK to obtain a very long decreasing sequence of lattices in $\mathbb{Z}^3$. After that we apply Lemma 2 to obtain a lower bound for possible unknown rational points. We get that if (x, y) is an unknown integral point, then

$$\log |x| \geq 1.03 \times 10^{580}.$$

This contradicts the bound for $\log |x|$ we obtained by Baker’s method. $\square$

The statement of the Theorem now easily follows. It is enough to find the values of $X = 2x^2 + 8x$. Afterwards the values for x and m are recovered immediately.

ACKNOWLEDGEMENT. The authors would like to thank the referee for the helpful comments and suggestions. The work was supported by the TÁMOP 4.2.1./B-09/1/KONV-2010-0007 project. The project is implemented through the New Hungary Development Plan, co-financed by the European Social Fund and the European Regional Development Fund.

References

- [1] A. BAKER, Bounds for the solutions of the hyperelliptic equation, *Proc. Cambridge Philos. Soc.* **65** (1969), 439–444.
- [2] A. BAKER and H. DAVENPORT, The equations $3x^2 - 2 = y^2$ and $8x^2 - 7 = z^2$, *Quart. J. Math. Oxford Ser. (2)* **20** (1969), 129–137.
- [3] A. BEHERA and G. K. PANDA, On the square roots of triangular numbers, *Fibonacci Quart.* **37**(2) (1999), 98–105.
- [4] A. BÉRCZES, K. LIPTAI and I. PINK, On generalized balancing sequences, *Fibonacci Quart.* **48**(2) (2010), 121–128.
- [5] Y. F. BILU and G. HANROT, Solving superelliptic Diophantine equations by Baker’s method, *Compositio Math.* **112**(3) (1998), 273–312.
- [6] W. BOSMA, J. CANNON and C. PLAYOUST, The Magma algebra system, I, The user language, *J. Symbolic Comput.* **24**3–4 (1997), 235–265, Computational algebra and number theory (London, 1993).
- [7] B. BRINDZA, On S -integral solutions of the equation $y^m = f(x)$, *Acta Math. Hungar.* **44**(1–2) (1984), 133–139.
- [8] N. BRUIN and M. STOLL, Deciding existence of rational points on curves: an experiment, *Experiment. Math.* **17**(2) (2008), 181–189.
- [9] N. BRUIN and M. STOLL, The Mordell–Weil sieve: proving non-existence of rational points on curves, *LMS J. Comput. Math.* **13** (2010), 272–306.
- [10] Y. BUGEAUD, Bounds for the solutions of superelliptic equations, *Compositio Math.* **107**(2) (1997), 187–219.
- [11] Y. BUGEAUD and K. GYÖRY, Bounds for the solutions of unit equations, *Acta Arith.* **74**(1) (1996), 67–80.

- [12] Y. BUGEAUD, M. MIGNOTTE and S. SIKSEK, Classical and modular approaches to exponential Diophantine equations, I, Fibonacci and Lucas perfect powers, *Ann. of Math.* (2) **163**(3) (2006), 969–1018.
- [13] Y. BUGEAUD, M. MIGNOTTE, S. SIKSEK, M. STOLL and SZ. TENGELY, Integral points on hyperelliptic curves, *Algebra Number Theory* **2**(8) (2008), 859–885.
- [14] J. W. S. CASSELS and E. V. FLYNN, Prolegomena to a middlebrow arithmetic of curves of genus 2, Vol. 230, London Mathematical Society Lecture Note Series, *Cambridge University Press, Cambridge*, 1996.
- [15] C. CHABAUTY, Sur les points rationnels des courbes algébriques de genre supérieur à l'unité, *C. R. Acad. Sci. Paris* **212** (1941), 882–885.
- [16] R. F. COLEMAN, Effective Chabauty, *Duke Math. J.* **52**(3) (1985), 765–770.
- [17] E. V. FLYNN, The Hasse principle and the Brauer–Manin obstruction for curves, *Manuscripta Math.* **115**(4) (2004), 437–466.
- [18] H. R. GALLEGOS-RUIZ, S -integral points on hyperelliptic curves, *Int. J. Number Theory* **7**(3) (2011), 803–824.
- [19] T. KOVÁCS, K. LIPTAI and P. OLAJOS, On (a, b) -balancing numbers, *Publ. Math. Debrecen* **77** (2010), 485–498.
- [20] E. LANDAU, Verallgemeinerung eines Pólyaschen Satzes auf algebraische Zahlkörper, 1918 (in *German*).
- [21] K. LIPTAI, Fibonacci balancing numbers, *Fibonacci Quart.* **42**(4) (2004), 330–340.
- [22] K. LIPTAI, Lucas balancing numbers, *Acta Math. Univ. Ostrav.* **14**(1) (2006), 43–47.
- [23] K. LIPTAI, F. LUCA, Á. PINTÉR and L. SZALAY, Generalized balancing numbers, *Indag. Math. (N.S.)* **20**(1) (2009), 87–100.
- [24] E. M. MATVEEV, An explicit lower bound for a homogeneous rational linear form in logarithms of algebraic numbers, II, *Izv. Ross. Akad. Nauk Ser. Mat.* **64**(6) (2000), 125–180.
- [25] G. K. PANDA, Sequence balancing and cobalancing numbers, *Fibonacci Quart.* **45**(3) (2007), 265–271 (2008).
- [26] A. PETHŐ and B. M. M. DE WEGER, Products of prime powers in binary recurrence sequences, I, The hyperbolic case, with an application to the generalized Ramanujan–Nagell equation, *Math. Comp.* **47**(176) (1986), 713–727.
- [27] V. SCHARASCHKIN, Local-global problems and the Brauer–Manin obstruction, PhD thesis, *University of Michigan*, 1999.
- [28] V. G. SPRINDŽUK, The arithmetic structure of integer polynomials and class numbers, Analytic number theory, mathematical analysis and their applications (dedicated to I. M. Vinogradov on his 85th birthday), *Trudy Mat. Inst. Steklov.* **143** (1977), 152–174, 210.
- [29] L. SZALAY, On the resolution of simultaneous Pell equations, *Ann. Math. Inform.* **34** (2007), 77–87.
- [30] P. M. VOUTIER, An upper bound for the size of integral solutions to $Y^m = f(X)$, *J. Number Theory* **53**(2) (1995), 247–271.
- [31] P. M. VOUTIER, An effective lower bound for the height of algebraic numbers, *Acta Arith.* **74**(1) (1996), 81–95.

SZABOLCS TENGELY
 MATHEMATICAL INSTITUTE
 UNIVERSITY OF DEBRECEN
 4010 DEBRECEN, P.O.BOX 12
 HUNGARY

E-mail: tengely@science.unideb.hu

(Received September 10, 2012; revised December 13, 2012)