

On the Banach algebra $(w_\infty(\Lambda), w_\infty(\Lambda))$ and applications to the solvability of matrix equations in $w_\infty(\Lambda)$

By BRUNO DE MALAFOSSE (Le Havre) and EBERHARD MALKOWSKY (Giessen)

Abstract. We apply the characterisation of the class $(w_\infty(\Lambda), w_\infty(\Sigma))$ and the fact that this is a Banach algebra to study the solvability in $w_\infty(\Lambda)$ of matrix equations of the form $\Delta_\rho^+ X = B$ and $\Delta_\rho X = B$, where Δ_ρ^+ and Δ_ρ are upper and lower triangular matrices. Finally, we obtain some results on infinite tridiagonal matrices considered as operators from $w_\infty(\Lambda)$ into itself, and study the solvability in $w_\infty(\Lambda)$ of matrix equations for tridiagonal matrices.

1. Introduction and known results

Let ω denote the set of all sequences $x = (x_k)_{k=0}^\infty$, and ℓ_∞ , c_0 and ϕ be the sets of all bounded, null and finite sequences, respectively. We write e and $e^{(n)}$ ($n = 0, 1, \dots$) for the sequences with $e_k = 1$ for all k , and $e_n^{(n)} = 1$ and $e_k^{(n)} = 0$ for $k \neq n$.

A *BK space* X is a Banach sequence space with continuous coordinates $P_k : X \rightarrow \mathbb{C}$ where $P_n(x) = x_k$ ($k = 0, 1, \dots$) for all $x \in X$. A *BK space* $X \supset \phi$ is said to have *AK* if $x = \sum_{k=0}^\infty x_k e^{(k)}$ for every sequence $x = (x_k)_{k=0}^\infty \in X$.

Let X be a subset of ω . Then the set $X^\beta = \{a \in \omega : \sum_{k=0}^\infty a_k x_k \text{ converges for all } x \in X\}$ is called the β -dual of X .

Let $A = (a_{nk})_{k=0}^\infty$ be an infinite matrix of complex numbers and $x = (x_k)_{k=0}^\infty \in \omega$. Then we write $A_n = (a_{nk})_{k=0}^\infty$ ($n = 0, 1, \dots$) and $A^k = (a_{nk})_{n=0}^\infty$ ($k = 0, 1, \dots$) for

Mathematics Subject Classification: Primary: 45H05; Secondary: 46A45.

Key words and phrases: summability, BK spaces, spaces of strongly bounded sequences, matrix transformations, infinite linear systems.

Work of the second author supported by the research project #174025 of the Serbian Ministry of Science, Technology and Environment.

the sequences in the n -th row and the k -th column of A , and $A_n(x) = \sum_{k=0}^{\infty} a_{nk}x_k$ provided the series converges. Given any subsets X and Y of ω , we write (X, Y) for the class of all infinite matrices A that map X into Y , that is $A_n \in X^\beta$ for all n , and $A(x) = (A_n(x))_{n=0}^{\infty} \in Y$.

Let X and Y be Banach spaces, and $B_X = \{x \in X : \|x\| \leq 1\}$ denote the unit ball in X . Then $\mathcal{B}(X, Y)$ denotes the Banach space of all bounded linear operators $L : X \rightarrow Y$ with the operator norm $\|L\| = \sup_{x \in B_X} \|L(x)\|$; $X^* = \mathcal{B}(X, \mathbb{C})$ is the *continuous dual* of X with the norm $\|f\| = \sup_{x \in B_X} |f(x)|$ for all $f \in X^*$. It is well known ([12, Theorem 4.2.8]) that if X and Y are *BK* spaces then $(X, Y) \subset \mathcal{B}(X, Y)$, that is, if $A \in (X, Y)$, then $L_A \in \mathcal{B}(X, Y)$ where $L_A(x) = A(x)$ for all $x \in X$.

Let $(\mu_n)_{n=0}^{\infty}$ be a non-decreasing sequence of positive reals tending to infinity. The set

$$\tilde{w}_\infty(\mu) = \left\{ x \in \omega : \sup_n \frac{1}{\mu_n} \sum_{k=0}^n |x_k| < \infty \right\}$$

was defined and studied in [9], where the concept of *exponentially bounded sequences* was introduced. When $\mu_n = n + 1$ ($n = 0, 1, \dots$) then this set reduces to w_∞ , the set of all sequences that are strongly bounded by the Cesàro method of order 1 ([1]). A non-decreasing sequence $\Lambda = (\lambda_n)_{n=0}^{\infty}$ of positive is called *exponentially bounded* if there is an integer $m \geq 2$ such that for all non-negative integers ν there is at least one term λ_n in the interval $I_m^{(\nu)} = [m^\nu, m^{\nu+1} - 1]$. It was shown ([9, Lemma 1]) that a non-decreasing sequence $\Lambda = (\lambda_n)_{n=0}^{\infty}$ is exponentially bounded, if and only if there are reals $s \leq t$ such that for some subsequence $(\lambda_{n(\nu)})_{\nu=0}^{\infty}$

$$0 < s \leq \frac{\lambda_{n(\nu)}}{\lambda_{n(\nu+1)}} \leq t < 1 \quad \text{for all } \nu = 0, 1, \dots;$$

such a subsequence is called an *associated subsequence*.

If $\Lambda = (\lambda_n)_{n=0}^{\infty}$ is an exponentially bounded sequence, and $(\lambda_{n(\nu)})_{\nu=0}^{\infty}$ is an associated subsequence with $\lambda_{n(0)} = \lambda_0$, then we write K_ν ($\nu = 0, 1, \dots$) for the set of all integers k with $n(\nu) \leq k \leq n(\nu+1) - 1$, and define the sets

$$w_0(\Lambda) = \left\{ x \in \omega : \lim_{\nu \rightarrow \infty} \frac{1}{\lambda_{n(\nu+1)}} \sum_{k \in K_\nu} |x_k| = 0 \right\}$$

and

$$w_\infty(\Lambda) = \left\{ x \in \omega : \sup_\nu \frac{1}{\lambda_{n(\nu+1)}} \sum_{k \in K_\nu} |x_k| < \infty \right\}.$$

The next result is well known.

Theorem 1.1 ([9, Theorem 1 (a), (b)]). *Let $(\mu_n)_{n=0}^\infty$ be a non-decreasing sequence of positive reals tending to infinity, $\Lambda = (\lambda_n)_{n=0}^\infty$ be an exponentially bounded sequence and $(\lambda_{n(\nu)})_{n=0}^\infty$ be an associated subsequence. Then $\tilde{w}_\infty(\mu)$ is a BK space with the norm $\|\cdot\|_\mu^-$ defined by*

$$\|x\|_\mu^- = \sup_n \frac{1}{\mu_n} \sum_{k=0}^n |x_k|. \quad (1.1)$$

Moreover, we have $\tilde{w}_\infty(\Lambda) = w_\infty(\Lambda)$, and the norms $\|\cdot\|_\Lambda^-$ and $\|\cdot\|_\Lambda$ are equivalent on $w_\infty(\Lambda)$, where

$$\|x\|_\Lambda = \sup_\nu \frac{1}{\lambda_{n(\nu+1)}} \sum_{k \in K_\nu} |x_k|. \quad (1.2)$$

Thus, in view of the previous result, we always assume, unless explicitly stated otherwise, that Λ is an exponentially bounded sequence and $(\lambda_{n(\nu)})_{n=0}^\infty$ is an associated subsequence, and we consider the spaces $w_0(\Lambda)$ and $w_\infty(\Lambda)$ with the norm $\|\cdot\| = \|\cdot\|_\Lambda$.

First we give the characterisation of the class $(w_\infty(\Lambda), w_\infty(\Sigma))$. Let $\Lambda = (\lambda_k)_{k=0}^\infty$ and $\Sigma = (\sigma_m)_{m=0}^\infty$ be exponentially bounded sequences and $(\lambda_{k(\nu)})_{\nu=0}^\infty$ and $(\sigma_{m(\mu)})_{\mu=0}^\infty$ be associated subsequences. Furthermore, let K_ν ($\nu = 0, 1, \dots$) and M_μ ($\mu = 0, 1, \dots$) be the sets of all integers k and m with $k(\nu) \leq k \leq k(\nu+1) - 1$ and $m(\mu) \leq m \leq m(\mu+1) - 1$. We obtain the characterisation of the class $(w_\infty(\Lambda), w_\infty(\Sigma))$ as the special case $p = 1$ of [10, Theorem 3.1].

Theorem 1.2. *We have $A \in (w_\infty(\Lambda), w_\infty(\Sigma))$ if and only if*

$$\begin{aligned} & \|A\|_{(w_\infty(\Lambda), w_\infty(\Sigma))} \\ &= \sup_\mu \left(\frac{1}{\sigma_{m(\mu+1)}} \max_{M(\mu) \subset M_\mu} \sum_{\nu=0}^\infty \lambda_{k(\nu+1)} \max_{k \in K_\nu} \left| \sum_{m \in M(\mu)} a_{mk} \right| \right) < \infty. \end{aligned} \quad (1.3)$$

If $A \in (w_\infty(\Lambda), w_\infty(\Sigma))$, then

$$\|A\|_{(w_\infty(\Lambda), w_\infty(\Sigma))} \leq \|L_A\| \leq 4 \cdot \|A\|_{(w_\infty(\Lambda), w_\infty(\Sigma))}. \quad (1.4)$$

Finally we need the following known result.

Theorem 1.3 ([10, Theorem 4.2]). *The class $(w_\infty(\Lambda), w_\infty(\Lambda))$ is a Banach algebra with respect to the norm*

$$\|A\|_{(\Lambda, \Lambda)} = \sup\{\|A\|_\Lambda : \|x\|_\Lambda \leq 1\} = \|L_A\| \quad \text{for all } A \in (w_\infty(\Lambda), w_\infty(\Lambda)).$$

In this paper, we use the characterisation of the class $(w_\infty(\Lambda), w_\infty(\Lambda))$ in Theorem 1.2 and the fact that this class is a Banach algebra to study the solvability in $w_\infty(\Lambda)$ of matrix equations of the form $\Delta_\rho^+ X = B$ and $\Delta_\rho X = B$, where Δ_ρ^+ and Δ_ρ are upper and lower triangular matrices. Finally, we obtain some results on infinite tridiagonal matrices considered as operators from $w_\infty(\Lambda)$ into itself, and study the solvability in $w_\infty(\Lambda)$ of matrix equations for tridiagonal matrices; these results are new. Our results on the Banach algebra and the deduced solvability in $w_\infty(\Lambda)$ of matrix equations are also useful for statistical convergence, and for A -statistical convergence, where A is an operator represented by an infinite matrix. They can also be applied in spectral theory and the study of operators generators of analytic semi-group, and be used in the intuitionistic fuzzy normed space (IFNS).

2. The sets s_τ^0 , $s_\tau^{(c)}$ and s_τ

We are going to study the solvability of certain matrix equations $AX = B$ in $w_\infty(\Lambda)$, where B belongs to the set $w_\infty(\Lambda)$ and A is an infinite matrix.

First, we recall some results on the sets s_τ^0 , $s_\tau^{(c)}$ and s_τ which are needed in the sequel. These sets are closely related to the sets c_0 , c and ℓ_∞ . We also define and study the sets $\widehat{C}$, $\widehat{\Gamma}$, Γ and $\widehat{C}_1$.

We slightly change our notations to match those normally used in the theory of infinite systems of linear equations, in particular, we consider sequences x as one-column matrices X . It is also more convenient for the indices to run from 1 to ∞ instead of from 0 to ∞ .

Thus, for a given infinite matrix $A = (a_{nk})_{n,k \geq 1}$, we consider the operators A_n for any integer $n \geq 1$, defined by

$$A_n(X) = \sum_{k=1}^{\infty} a_{nk} x_k \quad (2.1)$$

where $X = (x_n)_{n \geq 1}$ is a one-column matrix of complex numbers, and the series are assumed to be convergent. So we are led to the study of the infinite linear system

$$A_n(X) = b_n \quad \text{for } n = 1, 2, \dots \quad (2.2)$$

where $B = (b_n)_{n \geq 1}$ is a one-column matrix and X the unknown ([2], [4], [5], [6], [7], [8]). The equations in (2.2) can be written in the form $AX = B$, where $AX = (A_n(X))_{n \geq 1}$. We also consider A as an operator from a sequence space into another sequence space.

By cs , we denote the set of all convergent series.

2.1. The sets $D_\tau E$, where E is any of the sets c_0 , c , or ℓ_∞ . Throughout, we use the set

$$U^+ = \{(u_n)_{n \geq 1} \in \omega : u_n > 0 \text{ for all } n\}.$$

For a given sequence $\tau = (\tau_n)_{n \geq 1} \in U^+$, we define the infinite diagonal matrix $D_\tau = (\tau_n \delta_{nk})_{n,k \geq 1}$; D_τ can be considered as the operator $X = (x_n)_{n \geq 1} \mapsto D_\tau X = (\tau_n x_n)_{n \geq 1}$ from ω to itself. For any subset E of ω , we write

$$D_\tau E = \left\{ (x_n)_{n \geq 1} \in \omega : \left(\frac{x_n}{\tau_n} \right)_n \in E \right\},$$

in particular,

$$D_\tau E = \begin{cases} s_\tau^0 & \text{if } E = c_0, \\ s_\tau^{(c)} & \text{if } E = c, \\ s_\tau & \text{if } E = \ell_\infty \end{cases}$$

([3]). Each of the spaces $D_\tau E$, where $E \in \{c_0, c, \ell_\infty\}$, is a *BK space* normed by

$$\|X\|_{s_\tau} = \sup_{n \geq 1} \left(\frac{|x_n|}{\tau_n} \right), \quad (2.3)$$

and s_τ^0 has *AK*.

Now let $\tau = (\tau_n)_{n \geq 1}, \nu = (\nu_n)_{n \geq 1} \in U^+$. We write $S_{\tau, \nu}$ for the set of infinite matrices $A = (a_{nm})_{n,m \geq 1}$ such that $\sup_{n \geq 1} \left(\sum_{k=1}^{\infty} |a_{nk}| \tau_k / \nu_n \right) < \infty$. The set $S_{\tau, \nu}$ is a Banach space with the norm

$$\|A\|_{S_{\tau, \nu}} = \sup_{n \geq 1} \left(\frac{1}{\nu_n} \sum_{k=1}^{\infty} |a_{nk}| \tau_k \right).$$

It was proved in [8] that $A \in (s_\tau, s_\nu)$ if and only if $A \in S_{\tau, \nu}$, that is, $(s_\tau, s_\nu) = S_{\tau, \nu}$.

When $s_\tau = s_\nu$ then $S_\tau = S_{\tau, \nu}$ is a *Banach algebra with identity*, normed by $\|A\|_{S_\tau} = \|A\|_{S_{\tau, \tau}}$ ([3, 5]).

If $\tau = (r^n)_{n \geq 1}$, we write S_r, s_r, s_r^0 and $s_r^{(c)}$ for S_τ, s_τ, s_τ^0 and $s_\tau^{(c)}$, respectively. When $r = 1$, we obtain $s_1 = \ell_\infty, s_1^0 = c_0$ and $s_1^{(c)} = c$. Thus we have $S_1 = S_e$. It is well known ([12, Example 8.4.5A]) that

$$(s_1, s_1) = (c_0, s_1) = (c, s_1) = S_1.$$

For any subset E of ω , we put

$$AE = \{Y \in \omega : Y = AX \text{ for some } X \in E\}. \quad (2.4)$$

If F is a subset of ω , we write

$$F(A) = F_A = \{X \in \omega : Y = AX \in F\}. \quad (2.5)$$

2.2. Some properties of the sequence $C(X)X$. Here we deal with the operators represented by $C(\Lambda)$ and $\Delta(\Lambda)$. Let U be the set of all sequences $(u_n)_{n \geq 1}$ with $u_n \neq 0$ for all n . We define $C(\Lambda) = (c_{nk})_{n,k \geq 1}$ for $\Lambda = (\lambda_n)_{n \geq 1} \in U$, by

$$c_{nk} = \begin{cases} \frac{1}{\lambda_n} & \text{if } k \leq n \\ 0 & \text{otherwise.} \end{cases}$$

We write $C(\Lambda)^T = C^+(\Lambda)$, $C(e) = \Sigma$ and $\Sigma^+ = \Sigma^T$. The matrix $\Delta(\Lambda) = (c'_{nk})_{n,k \geq 1}$ with

$$c'_{nk} = \begin{cases} \lambda_n & \text{if } k = n \\ -\lambda_{n-1} & \text{if } k = n-1 \text{ and } n \geq 2 \\ 0 & \text{otherwise} \end{cases}$$

is the inverse of $C(\Lambda)$ ([3], [4]). We need to recall some results given in [3], [7]. For this, we consider the following sets

$$\begin{aligned} \widehat{C}_1 &= \left\{ X = (x_n)_{n \geq 1} \in U^+ : \sup_n \left[\frac{1}{x_n} \left(\sum_{k=1}^n x_k \right) \right] < \infty \right\}, \\ \widehat{C} &= \left\{ X = (x_n)_{n \geq 1} \in U^+ : \left(\frac{1}{x_n} \left(\sum_{k=1}^n x_k \right) \right)_{n \geq 1} \in c \right\}, \\ \widehat{C}_1^+ &= \left\{ X \in U^+ \cap cs : \sup_n \left[\frac{1}{x_n} \left(\sum_{k=n}^\infty x_k \right) \right] < \infty \right\}, \\ \Gamma &= \left\{ X \in U^+ : \overline{\lim}_{n \rightarrow \infty} \left(\frac{x_{n-1}}{x_n} \right) < 1 \right\}, \\ \widehat{\Gamma} &= \left\{ X \in U^+ : \lim_{n \rightarrow \infty} \left(\frac{x_{n-1}}{x_n} \right) < 1 \right\}, \\ \Gamma^+ &= \left\{ X \in U^+ : \overline{\lim}_{n \rightarrow \infty} \left(\frac{x_{n+1}}{x_n} \right) < 1 \right\}. \end{aligned}$$

Note that $X \in \Gamma^+$ if and only if $1/X \in \Gamma$. We will see in Lemma 2.1 that if $X \in \widehat{C}_1$, then $x_n \rightarrow \infty$ ($n \rightarrow \infty$). Furthermore, $X \in \Gamma$ if and only if there is an integer $q \geq 1$ such that

$$\gamma_q(X) = \sup_{n \geq q+1} \left(\frac{x_{n-1}}{x_n} \right) < 1.$$

Writing

$$[C(X)X]_n = \frac{1}{x_n} \left(\sum_{k=1}^n x_k \right),$$

we obtain the following the following result from [5, Proposition 2.1].

Lemma 2.1. *Let $\tau \in U^+$. Then*

- i) $(\tau_{n-1}/\tau_n)_{n \geq 1} \rightarrow 0$ ($n \rightarrow \infty$) if and only if $[C(\tau)\tau]_n \rightarrow 1$ ($n \rightarrow \infty$).
- ii) a) $\tau \in \widehat{C}$ if and only if $(\tau_{n-1}/\tau_n)_{n=0}^\infty \in c$,
b) $[C(\tau)\tau]_n \rightarrow l$ ($n \rightarrow \infty$) if and only if $\tau_{n-1}/\tau_n \rightarrow 1 - 1/l$ ($n \rightarrow \infty$).
- iii) If $\tau \in \widehat{C}_1$ then there are $K > 0$ and $\gamma > 1$ such that

$$\tau_n \geq K\gamma^n \quad \text{for all } n. \quad (2.6)$$

- iv) The condition $\tau \in \Gamma$ implies $\tau \in \widehat{C}_1$ and there is a real $b > 0$ such that

$$[C(\tau)\tau]_n \leq \frac{1}{1 - \gamma_q(\tau)} + b\gamma_q(\tau)^n \quad \text{for } n \geq q + 1. \quad (2.7)$$

- v) The condition $\tau \in \Gamma^+$ implies $\tau \in \widehat{C}_1^+$.

It was also shown in [7] that

$$\widehat{C} = \widehat{\Gamma} \subset \Gamma \subset \widehat{C}_1.$$

3. The Characterizations of some operators mapping $w_\infty(\Lambda)$ into itself

In this section, we apply Theorem 1.3 to infinite matrices such as Δ_ρ^+ , Δ_ρ considered as operators from $w_\infty(\Lambda)$ into $D_a w_\infty(\Lambda)$.

Throughout, we assume that $\Lambda = (\lambda_n)_{n \geq 1}$ is an exponentially bounded sequence, and $(\lambda_{n_i})_{i \geq 1}$ is an associated subsequence.

As an immediate consequence of Theorem 1.3, we see that $(w_\infty(\Lambda), w_\infty(\Lambda))$ is a Banach algebra with the norm

$$\|A\|_{(\Lambda, \Lambda)}^- = \sup_{X \neq 0} \left(\frac{\|AX\|_{\Lambda}^-}{\|X\|_{\Lambda}^-} \right). \quad (3.1)$$

where $\|\cdot\|^-$ is the norm defined in (1.1).

Since the sequence Λ with $\lambda_n = n$ ($n = 1, 2, \dots$) is exponentially bounded, we obtain in particular that (w_∞, w_∞) is a Banach algebra with the norm in (3.1) where

$$\|X\|^- = \sup_n \left(\frac{1}{n} \sum_{k=1}^n |x_k| \right).$$

3.1. Some properties of the matrix map Δ_ρ^+ .

3.1.1. *The map Δ_ρ^+ considered as an operator from $w_\infty(\Lambda)$ into itself. Let $\rho = (\rho_n)_{n \geq 1}$ and consider the infinite matrix $\Delta_\rho^+ = \{(\Delta_\rho^+)_{nk}\}$ defined by*

$$(\Delta_\rho^+)_{nk} = \begin{cases} 1 & (k = n) \\ -\rho_n & (k = n + 1) \\ 0 & (\text{otherwise}) \end{cases} \quad \text{for all } n, k.$$

First we see that if ρ and $(\lambda_{n+1}/\lambda_n)_{n \geq 1} \in \ell_\infty$ then $\Delta_\rho^+ \in (w_\infty(\Lambda), w_\infty(\Lambda))$. Indeed, we have

$$\Delta_\rho^+ x_n = x_n - \rho_n x_{n+1} \quad \text{for all } n$$

and

$$\begin{aligned} \frac{1}{\lambda_n} \sum_{k=1}^n |x_k - \rho_k x_{k+1}| &\leq \frac{1}{\lambda_n} \sum_{k=1}^n |x_k| + \frac{\lambda_{n+1}}{\lambda_n} \sup_n |\rho_n| \frac{1}{\lambda_{n+1}} \sum_{k=2}^{n+1} |x_k| \\ &\leq \left(1 + \sup_n \left(\frac{\lambda_{n+1}}{\lambda_n} \right) \sup_n |\rho_n| \right) \|X\|_{w_\infty(\Lambda)}^-. \end{aligned}$$

for all $X \in w_\infty(\Lambda)$ and for all n . This shows that $\Delta_\rho^+ X \in w_\infty(\Lambda)$ for all $X \in w_\infty(\Lambda)$.

More precisely we have the following result.

Theorem 3.1. *Let $\Lambda \in U^+$ be an exponentially bounded sequence and assume*

$$\overline{\lim}_{n \rightarrow \infty} \left(\frac{\lambda_{n+1}}{\lambda_n} \right) < \infty \quad \text{and} \quad \overline{\lim}_{n \rightarrow \infty} |\rho_n| < \frac{1}{\overline{\lim}_{n \rightarrow \infty} \left(\frac{\lambda_{n+1}}{\lambda_n} \right)}. \quad (3.2)$$

For given $B \in w_\infty(\Lambda)$ the equation $\Delta_\rho^+ X = B$ has a unique solution in $w_\infty(\Lambda)$ given by

$$x_n = b_n + \sum_{i=n+1}^{\infty} \left(\prod_{j=n}^{i-1} \rho_j \right) b_i \quad \text{for all } n. \quad (3.3)$$

PROOF. Let

$$\Sigma_\rho^{+(N)} = \begin{pmatrix} [\Delta_\rho^{+(N)}]^{-1} & 0 \\ 0 & 1 \\ & & . \end{pmatrix},$$

where $\Delta_\rho^{+(N)}$ is the finite matrix whose elements are those of the N first rows and columns of Δ_ρ^+ . The finite matrix $\Delta_\rho^{+(N)}$ is invertible, since it is an upper

triangle. We get $\Delta_\rho^+ \Sigma_\rho^{+(N)} = (a_{nk})_{n,k \geq 1}^\infty$, with $a_{nn} = 1$ for all n ; $a_{n,n+1} = -\rho_n$ for all $n \geq N$; and $a_{nk} = 0$ otherwise. For any given $X \in w_\infty(\Lambda)$, we have $(I - \Delta_\rho^+ \Sigma_\rho^{+(N)})X = (\xi_n(X))_{n \geq 1}$ with $\xi_n(X) = 0$ for all $n \leq N-1$ and $\xi_n(X) = \rho_n x_{n+1}$ for all $n \geq N$. Now we put $K_N = \sup_{n \geq N} (\lambda_{n+1}/\lambda_n) \sup_{k \geq N} |\rho_k|$ and show that (3.2) implies $K_N < 1$ for N large enough. For this let $\varepsilon > 0$ be given. From (3.2), we have $\overline{\lim}_{n \rightarrow \infty} |\rho_n| = l < \infty$, since $\overline{\lim}_{n \rightarrow \infty} \lambda_{n+1}/\lambda_n \geq 1$. Then there is an integer N_1 such that

$$\sup_{n \geq N_1} |\rho_n| < l + \varepsilon.$$

Now since $\overline{\lim}_{n \rightarrow \infty} \lambda_{n+1}/\lambda_n < \infty$, (3.2) implies $\overline{\lim}_{n \rightarrow \infty} \lambda_{n+1}/\lambda_n = L < 1/l$, and as above there is an integer N_2 such that

$$\sup_{n \geq N_2} \left(\frac{\lambda_{n+1}}{\lambda_n} \right) < L + \varepsilon.$$

Since $lL < 1$, for ε small enough, taking $N = \max\{N_1, N_2\}$, we then have

$$K_N \leq (l + \varepsilon)(L + \varepsilon) = lL + l\varepsilon + L\varepsilon + \varepsilon^2 < 1.$$

Now we obtain

$$\begin{aligned} \|(I - \Delta_\rho^+ \Sigma_\rho^{+(N)})X\|_{w_\infty(\Lambda)}^- &= \sup_{n \geq N} \left(\frac{1}{\lambda_n} \sum_{k=N}^n |\rho_k x_{k+1}| \right) \\ &\leq \sup_{n \geq N} \left[\left(\sup_{k \geq N} |\rho_k| \right) \frac{1}{\lambda_n} \sum_{k=N+1}^{n+1} |x_k| \right] \\ &\leq \sup_{n \geq N} \left(\frac{\lambda_{n+1}}{\lambda_n} \right) \sup_{k \geq N} |\rho_k| \sup_{n \geq N} \left(\frac{1}{\lambda_{n+1}} \sum_{k=N+1}^{n+1} |x_k| \right) \\ &\leq \left[\sup_{n \geq N} \left(\frac{\lambda_{n+1}}{\lambda_n} \right) \sup_{k \geq N} |\rho_k| \right] \|X\|_{w_\infty(\Lambda)}^-. \end{aligned}$$

Then $\|(I - \Delta_\rho^+ \Sigma_\rho^{+(N)})X\|_{w_\infty(\Lambda)}^- \leq K_N \|X\|_{w_\infty(\Lambda)}^-$ and we conclude

$$\|I - \Delta_\rho^+ \Sigma_\rho^{+(N)}\|_{(\Lambda, \Lambda)}^- = \sup_{X \neq 0} \left(\frac{\|(I - \Delta_\rho^+ \Sigma_\rho^{+(N)})X\|_{w_\infty(\Lambda)}^-}{\|X\|_{w_\infty(\Lambda)}^-} \right) \leq K_N < 1.$$

Then (3.2) implies $\|I - \Delta_\rho^+ \Sigma_\rho^{+(N)}\|_{(\Lambda, \Lambda)}^- \leq K_N < 1$ and $\Delta_\rho^+ \Sigma_\rho^{+(N)}$ has a unique inverse in the Banach algebra $(w_\infty(\Lambda), w_\infty(\Lambda))$. Since obviously $\Sigma_\rho^{+(N)}$ is

bijjective from $w_\infty(\Lambda)$ into itself, the operators defined by $\Delta_\rho^+ \Sigma_\rho^{+(N)}$ and $\Delta_\rho^+ = (\Delta_\rho^+ \Sigma_\rho^{+(N)})(\Sigma_\rho^{+(N)})^{-1}$ are bijective from $w_\infty(\Lambda)$ into itself. So, for any given $B \in w_\infty(\Lambda)$, the equation $\Delta_\rho^+ X = B$ has a unique solution in $w_\infty(\Lambda)$. Finally $(\Delta_\rho^+)^{-1} = \sum_{i=0}^\infty (I - \Delta_\rho^+)^i = [[(\Delta_\rho^+)^T]^{-1}]^T$ and an elementary calculation shows that

$$(\Delta_\rho^+)^{-1} = [[(\Delta_\rho^+)^T]^{-1}]^T = \begin{pmatrix} 1 & \rho_1 & \rho_1 \rho_2 & \cdot & \cdot \\ & 1 & \rho_2 & \rho_2 \rho_3 & \cdot \\ & & \cdot & \cdot & \cdot \\ 0 & & & 1 & \rho_n \\ & & & & \cdot \end{pmatrix}.$$

We then obtain (3.3). This completes the proof. $\square$

Proposition 3.2. *For given $B \in w_\infty(\Lambda)$, the equation $\Delta_\rho^+ X = B$ has a unique solution in $w_\infty(\Lambda)$ given by (3.3) in the following cases*

- i) $\Lambda \in \widehat{C}_1$ and $\overline{\lim}_{n \rightarrow \infty} (\lambda_{n+1}/\lambda_n) |\rho_n| < 1$;
- ii) $\Lambda \notin \widehat{C}_1$ and condition (3.2) holds.

PROOF. i) The condition $\Lambda \in \widehat{C}_1$ implies $w_\infty(\Lambda) = s_\Lambda$. Furthermore, the condition $\overline{\lim}_{n \rightarrow \infty} (\lambda_{n+1}/\lambda_n) |\rho_n| < 1$ implies that there is an integer N such that

$$\sup_{n \geq N} \left(\frac{\lambda_{n+1}}{\lambda_n} |\rho_n| \right) < 1.$$

Now since $(w_\infty(\Lambda), w_\infty(\Lambda)) = (s_\Lambda, s_\Lambda) = S_\Lambda$, we deduce

$$\left\| I - \Delta_\rho^+ \Sigma_\rho^{+(N)} \right\|_{(\Lambda, \Lambda)}^- = \left\| I - \Delta_\rho^+ \Sigma_\rho^{+(N)} \right\|_{S_\Lambda} = \sup_{n \geq N} \left(\frac{\lambda_{n+1}}{\lambda_n} |\rho_n| \right) < 1.$$

So we have shown i).

- ii) This is a direct consequence of Theorem 3.1. $\square$

Remark 3.3. Note that since

$\overline{\lim}_{n \rightarrow \infty} (\lambda_{n+1}/\lambda_n) |\rho_n| \leq \overline{\lim}_{n \rightarrow \infty} (\lambda_{n+1}/\lambda_n) \overline{\lim}_{n \rightarrow \infty} |\rho_n|$, condition (3.2) is weaker than the condition $\overline{\lim}_{n \rightarrow \infty} (\lambda_{n+1}/\lambda_n) |\rho_n| < 1$ in Proposition 3.2 ii).

Putting $\Lambda^- = (1, \lambda_1, \dots, \lambda_{n-1}, \dots)$, we easily deduce the following result.

Corollary 3.4. *We assume $\Lambda \in \widehat{C}_1$. Then we have*

- i) $w_\infty(\Lambda) = s_\Lambda$;

- ii) $w_\infty(\Lambda)(\Delta^+) = w_\infty(\Lambda^-) = s_{\Lambda^-}$ and, for any given $B \in w_\infty(\Lambda)$, the equation

$$\Delta^+ X = B \quad (3.4)$$

has infinitely many solutions in $w_\infty(\Lambda^-)$ given by $x_n = u - \sum_{k=1}^{n-1} b_k$, where u is an arbitrary scalar.

PROOF. i) Part (i) comes from [4].

ii) It can easily be seen that $\Lambda \in \widehat{C_1}$ implies $\Lambda^- \in \widehat{C_1}$ and since $w_\infty(\Lambda) = s_\Lambda$, we only have to show that $s_\Lambda(\Delta^+) = s_{\Lambda^-}$. From the inequality

$$\frac{\lambda_{n-1}}{\lambda_n} \leq \sup_n \left(\frac{\sum_{k=1}^n \lambda_k}{\lambda_n} \right) < \infty \quad \text{for all } n,$$

we deduce that $\sup_n (\lambda_{n-1}/\lambda_n) < \infty$ and $\Delta^+ \in (s_{\Lambda^-}, s_\Lambda)$. Then, for any given $B \in s_\Lambda$, the solutions of the equation $\Delta^+ X = B$ are given by $x_1 = -u$, and

$$x_n = u + \sum_{k=1}^{n-1} b_k \quad \text{for } n \geq 2, \quad \text{where } u \text{ is an arbitrary scalar.} \quad (3.5)$$

So there exists a real $K > 0$, such that

$$\frac{|x_n|}{\lambda_{n-1}} = \frac{|u + \sum_{k=1}^{n-1} b_k|}{\lambda_{n-1}} \leq \sup_n \left(\frac{|u| + K(\sum_{k=1}^{n-1} \lambda_k)}{\lambda_{n-1}} \right) < \infty,$$

since $\sup_n (|u|/\lambda_{n-1}) < \infty$. So $X \in s_{\Lambda^-}$ and we conclude that Δ^+ is surjective from s_{Λ^-} into s_Λ . Then $\Lambda \in \widehat{C_1}$ implies

$$s_\Lambda(\Delta^+) = s_{\Lambda^-}.$$

The previous argument shows that the equation $\Delta^+ X = B$ has infinitely many solutions in $w_\infty(\Lambda^-) = s_{\Lambda^-}$ given by $x_n = u - \sum_{k=1}^{n-1} b_k$. This completes the proof. $\square$

3.1.2. *Properties of $\Delta^+(\mu)$ considered as an operator from $D_\tau w_\infty(\Lambda)$ to $D_{\tau\mu} w_\infty(\Lambda)$.* Here we consider the set

$$D_\tau w_\infty(\Lambda) = \left\{ X = (x_n)_{n \geq 1} : \sup_n \left(\frac{1}{\lambda_n} \sum_{k=1}^n \left| \frac{x_k}{\tau_k} \right| \right) < \infty \right\}.$$

We need the following lemma.

Lemma 3.5. *Let E and F be linear subspaces of ω and assume $A \in (E, F)$. If A is a surjective map from E to F and $\text{Ker } A = \{0\}$, then*

$$F(A) = E.$$

PROOF. Let $X \in F(A)$. Then $Y = AX \in F$. Now we show that $AX \in F$ implies $X \in E$. For this, we assume that $AX \in F$ and $X \notin E$. Since A is surjective

from E to F , there is $X_0 \in E$ such that $Y = AX = AX_0$ and $X - X_0 \in \text{Ker } A$. So $X = X_0 \in E$ which gives a contradiction. Thus we have shown $F(A) \subset E$. The inclusion $E \subset F(A)$ is immediate. This shows the lemma. $\square$

Note, for instance, that for $0 < r < 1$, we have $\|I - \Delta^+\|_{s_r} = r < 1$. This means that Δ^+ is bijective from s_r to itself. Since $e \in \text{Ker } \Delta^+ \setminus \{\tau\}$, we have $\Delta^+e = 0 \in s_r$ and $e \in s_r(\Delta^+) \setminus s_r$. This shows $s_r(\Delta^+) \neq s_r$.

We have as a direct consequence of Theorem 3.1

Corollary 3.6. *Let $\Lambda, \mu, \tau \in U^+$ and assume*

$$\overline{\lim}_{n \rightarrow \infty} \frac{\tau_{n+1}}{\tau_n} < \frac{1}{\overline{\lim}_{n \rightarrow \infty} \left(\frac{\lambda_{n+1}}{\lambda_n} \right)}. \quad (3.6)$$

- i) *Then Δ^+ is bijective from $D_\tau w_\infty(\Lambda)$ into itself and $\Delta^+(\mu)$ is bijective from $D_\tau w_\infty(\Lambda)$ into $D_{\tau\mu} w_\infty(\Lambda)$.*
- ii) *$(D_\tau w_\infty(\Lambda))(C^+(\mu)) = D_{\tau\mu} w_\infty(\Lambda)$.*

PROOF. i) Here we have that $\Delta^+X = B$ with $B, X \in D_\tau w_\infty(\Lambda)$ (X being the unknown) is equivalent to $(D_{1/\tau} \Delta^+ D_\tau)X' = D_{1/\tau} B$ where $D_{1/\tau} B \in w_\infty(\Lambda)$ and $X' = D_{1/\tau} X \in w_\infty(\Lambda)$. So $\rho_n = \tau_{n+1}/\tau_n$ for all n and, by Theorem 3.1, the operator Δ^+ is bijective from $D_\tau w_\infty(\Lambda)$ into itself. Now since $\Delta^+(\mu) = D_\mu \Delta^+$, we easily conclude that Δ^+ is bijective from $D_\tau w_\infty(\Lambda)$ into itself and D_μ is bijective from $D_\tau w_\infty(\Lambda)$ into $D_{\tau\mu} w_\infty(\Lambda)$ and $\Delta^+(\mu)$ is bijective from $D_\tau w_\infty(\Lambda)$ into $D_{\tau\mu} w_\infty(\Lambda)$.

ii) Since $\Delta^+(\mu)$ is bijective from $D_\tau w_\infty(\Lambda)$ into $D_{\tau\mu} w_\infty(\Lambda)$, $C^+(\mu) = \Sigma^+ D_{1/\mu} = (\Delta^+(\mu))^{-1}$ is bijective from $D_{\tau\mu} w_\infty(\Lambda)$ into $D_\tau w_\infty(\Lambda)$. Now the equation $C^+(\mu)X = 0$ is equivalent to $\Sigma^+ Y = 0$ with $Y = D_{1/\mu} X = (y_n)_{n \geq 1}$ and $Y \in cs$. Then we have

$$\sum_{k=n}^{\infty} y_k = 0 \quad \text{for } n = 1, 2, \dots$$

We conclude $y_1 = y_2 = \dots = y_n = 0$ for all n and $\text{Ker } \Sigma^+ = \text{Ker } C^+(\mu) = \{0\}$. By Lemma 3.5, we conclude $(D_\tau w_\infty(\Lambda))(C^+(\mu)) = D_{\tau\mu} w_\infty(\Lambda)$. $\square$

Corollary 3.7. *If $\tau \in \Gamma^+$, then the equation $\Delta^+X = B$ has a unique solution in $D_\tau w_\infty$ for any given $B \in D_\tau w_\infty$.*

PROOF. Indeed by Corollary 3.6, the condition $\tau \in \Gamma^+$ implies that (3.2) is satisfied for $\lambda_n = n$, that is,

$$\overline{\lim}_{n \rightarrow \infty} \frac{\tau_{n+1}}{\tau_n} < \frac{1}{\overline{\lim}_{n \rightarrow \infty} \left(\frac{n+1}{n} \right)} = 1. \quad \square$$

These results lead to the next remarks.

Remark 3.8. Note that if $\Lambda \in \Gamma^+$ then (3.6) yields $1 < 1/\overline{\lim}_{n \rightarrow \infty} (\lambda_{n+1}/\lambda_n)$ and Δ^+ is bijective from $w_\infty(\Lambda)$ into itself.

Remark 3.9. Using the inverse of the infinite matrix $(\Delta_\rho^+)^T$ ([2]), we see that, under (3.2), we also have that $\sum_{k=1}^n |x_k| = O(\lambda_n)$ ($n \rightarrow \infty$) implies

$$\sum_{k=1}^n \left| x_k + \sum_{i=k+1}^{\infty} \left(\prod_{j=k}^{i-1} \rho_j \right) x_i \right| = O(\lambda_n) \quad (n \rightarrow \infty) \quad \text{for all } X \in \omega.$$

Now we consider an example.

Example 3.10. We choose $\lambda_n = n$ and $\rho_n = a_n - 1/n^\alpha$ for all n , where $(a_n)_n \in c$ is a positive sequence with $a_n \rightarrow l$ for some $l < 1$ ($n \rightarrow \infty$), and $\alpha > 0$. We see that if $\sum_{k=1}^n |x_k| = O(n)$ ($n \rightarrow \infty$) then

$$\sum_{k=1}^n \left| x_k + \sum_{i=k+1}^{\infty} \left[\prod_{j=k}^{i-1} \left(a_j - \frac{1}{j^\alpha} \right) \right] x_i \right| = O(n) \quad (n \rightarrow \infty)$$

for all $X \in \omega$.

Remark 3.11. Note that, by Corollary 3.4 ii), we have $\Lambda \in \widehat{C}_1$ if and only if $w_\infty(\Lambda)(\Delta^+) = w_\infty(\Lambda^-)$. Indeed, since $w_\infty(\Lambda) = s_\Lambda$, it is enough to assume that $s_\Lambda(\Delta^+) = s_{\Lambda^-}$. If we take $B = \Lambda \in s_\Lambda$, then the solutions of the equation $\Delta^+ X = B$ belong to s_{Λ^-} , and are given by $x_n = x_1 - \sum_{k=1}^{n-1} \lambda_k$, where x_1 is an arbitrary scalar and

$$\frac{x_n}{\lambda_{n-1}} = \frac{x_1}{\lambda_{n-1}} - \frac{1}{\lambda_{n-1}} \left(\sum_{k=1}^{n-1} \lambda_k \right) = O(1) \quad (n \rightarrow \infty).$$

Putting $x_1 = 0$, we conclude that $\Lambda \in \widehat{C}_1$.

3.2. Some properties of the matrix map Δ_ρ .

3.2.1. *On the operator Δ_ρ mapping $w_\infty(\Lambda)$ into itself.* Now we study the triangle $\Delta_\rho = (\Delta_\rho^+)^T$, that is,

$$\Delta_\rho = \begin{pmatrix} 1 & & & & \\ -\rho_1 & 1 & & & 0 \\ & \cdot & \cdot & & \\ & & -\rho_{n-1} & 1 & \cdot \\ & & & \cdot & \cdot \end{pmatrix}.$$

considered as an operator that maps $w_\infty(\Lambda)$ to itself.

As in Subsection 3.1.1, we easily see that $\Delta_\rho \in (w_\infty(\Lambda), w_\infty(\Lambda))$, if $(\lambda_{n-1}/\lambda_n)_{n \geq 2} \in \ell_\infty$.

Theorem 3.12. *Let $\Lambda \in U^+$ be an exponentially bounded sequence. We assume*

$$\overline{\lim}_{n \rightarrow \infty} |\rho_n| < \frac{1}{\overline{\lim}_{n \rightarrow \infty} \left(\frac{\lambda_{n-1}}{\lambda_n} \right)}. \quad (3.7)$$

Then, for any given $B \in w_\infty(\Lambda)$, the equation $\Delta_\rho X = B$ has a unique solution in $w_\infty(\Lambda)$ given by $x_1 = b_1$ and

$$x_n = b_n + \sum_{k=1}^{n-1} \left(\prod_{j=k}^{n-1} \rho_j \right) b_k \quad \text{for all } n \geq 2. \quad (3.8)$$

PROOF. Let $\Sigma_\rho^{(N)}$ be defined by

$$\begin{pmatrix} [\Delta_\rho^{(N)}]^{-1} & 0 \\ & 1 \\ 0 & . \end{pmatrix},$$

where $\Delta_\rho^{(N)}$ is the finite matrix whose elements are those of the N first rows and columns of Δ_ρ . The finite matrix $\Delta_\rho^{(N)}$ is invertible, since it is a triangle. We get $\Sigma_\rho^{(N)} \Delta_\rho = (a_{nk})_{n,k \geq 1}$, with $a_{nn} = 1$ for all n ; $a_{n,n-1} = -\rho_{n-1}$ for all $n \geq N+1$; and $a_{nm} = 0$ otherwise. For any given $X \in w_\infty(\Lambda)$, we have $(I - \Sigma_\rho^{(N)} \Delta_\rho)X = (\xi_n(X))_{n \geq 1}$ with $\xi_n(X) = 0$ for all $n \leq N$ and $\xi_n(X) = \rho_{n-1}x_{n-1}$ for all $n \geq N+1$. Now put $K'_N = \sup_{n \geq N+1} (\lambda_{n-1}/\lambda_n) \sup_{k \geq N+1} |\rho_{k-1}|$. Then we have as in Theorem 3.1

$$\sup_{n \geq N+1} \left[\left(\sup_{N \leq k \leq n} |\rho_k| \right) \frac{\lambda_{n-1}}{\lambda_n} \right] \leq K'_N < 1 \quad \text{for all } n \geq N. \quad (3.9)$$

Furthermore, we obtain

$$\begin{aligned} \left\| (I - \Sigma_\rho^{(N)} \Delta_\rho)X \right\|_{w_\infty(\Lambda)}^- &= \sup_{n \geq N+1} \left(\frac{1}{\lambda_n} \sum_{k=N+1}^n |\rho_{k-1}x_{k-1}| \right) \\ &\leq \sup_{n \geq N+1} \left[\left(\sup_{N+1 \leq k \leq n} |\rho_{k-1}| \right) \frac{\lambda_{n-1}}{\lambda_n} \right] \sup_{n \geq N} \left(\frac{1}{\lambda_{n-1}} \sum_{k=N}^{n-1} |x_k| \right) \leq K'_N \|X\|_{w_\infty(\Lambda)}^-. \end{aligned}$$

Finally, using (3.9), we get

$$\left\| I - \Sigma_\rho^{(N)} \Delta_\rho \right\|_{(\Lambda, \Lambda)}^- \leq K'_N < 1,$$

and we conclude, reasoning as Theorem 3.1. $\square$

Remark 3.13. Note that since Λ is a non-decreasing sequence, we do not need to assume $\overline{\lim}_{n \rightarrow \infty} (\lambda_{n-1}/\lambda_n) < \infty$ in Theorem 3.12.

We immediately get

Corollary 3.14. *Under (3.7), we have $w_\infty(\Lambda)(\Delta_\rho) = w_\infty(\Lambda)$.*

Corollary 3.15. i) *We assume that (3.7) holds. Then*

$$\sup_n \left(\frac{1}{\lambda_n} \sum_{k=1}^n |x_k| \right) < \infty$$

implies

$$\sup_n \left\{ \frac{1}{\lambda_n} \sum_{k=2}^n \left| x_k + \sum_{j=1}^{k-1} \left(\prod_{i=j}^{k-1} \rho_i \right) x_j \right| \right\} < \infty \quad \text{for all } X.$$

ii) *If $\Lambda \in \Gamma$ and $\overline{\lim}_{n \rightarrow \infty} |\rho_n| < 1$, then*

$$\sup_n \left(\frac{|x_k|}{\lambda_n} \right) < \infty$$

implies

$$\sup_n \left\{ \frac{1}{\lambda_n} \left| x_n + \sum_{j=1}^{n-1} \left(\prod_{i=j}^{n-1} \rho_i \right) x_j \right| \right\} < \infty \quad \text{for all } X.$$

PROOF. i) Part i) is an immediate consequence of Theorem 3.12.

ii) Since $\Lambda \in \Gamma$, that is, $\overline{\lim}_{n \rightarrow \infty} (\lambda_{n-1}/\lambda_n) < 1$, we have

$$\overline{\lim}_{n \rightarrow \infty} |\rho_n| < 1 < \frac{1}{\overline{\lim}_{n \rightarrow \infty} (\lambda_{n-1}/\lambda_n)}$$

and condition (3.7) follows. Now, by Corollary 3.4, we have $w_\infty(\Lambda) = s_\Lambda$ and, by Theorem 3.12, Δ_ρ is bijective from s_Λ into itself. We conclude, using identity (3.8). $\square$

Remark 3.16. Putting $\Delta_1 = \Delta$, we see that if $\Lambda \in \Gamma$ then $w_\infty(\Lambda)(\Delta) = w_\infty(\Lambda)$. Indeed by Theorem 3.12, condition (3.7) implies $\overline{\lim}_{n \rightarrow \infty} (\lambda_{n-1}/\lambda_n) < 1$, that is, $\Lambda \in \Gamma$. Since $\Gamma \subset \widehat{C}_1$, applying Corollary 3.4, we conclude $w_\infty(\Lambda) = s_\Lambda$ and $s_\Lambda(\Delta) = s_\Lambda$.

These results lead to consider the sets $[C(\Lambda), C(\mu)]_\infty$ and $[C(\Lambda), \Delta_\rho(\mu)]_\infty$, ([5]). In this way we obtain the following results.

3.2.2. *On the sets $[C(\Lambda), C(\mu)]_\infty$ and $[C(\Lambda), \Delta_\rho(\mu)]_\infty$.* Now we can give some applications of the previous results. For this, we define for $\Lambda, \mu \in U^+$ the set

$$[C(\Lambda), C(\mu)]_\infty = \{X \in \omega : C(\Lambda)(|C(\mu)X|) \in \ell_\infty\}.$$

Then we have

$$[C(\Lambda), C(\mu)]_\infty = \left\{ X \in \omega : \sup_n \left(\frac{1}{\lambda_n} \sum_{k=1}^n \frac{1}{\mu_k} \left| \sum_{i=1}^k x_i \right| \right) < \infty \right\}.$$

It was shown in [5, Theorem 3.1] that if $\Lambda, \Lambda\mu \in \widehat{C}_1$ then

$$[C(\Lambda), C(\mu)]_\infty = s_{\Lambda\mu}.$$

This result can be improved here as follows. We put $C_1 = C((n)_{n \geq 1})$. The following result holds.

Proposition 3.17. *Let $\mu, \Lambda \in U^+$.*

i) *If*

$$\overline{\lim}_{n \rightarrow \infty} \frac{\mu_{n-1}}{\mu_n} < \frac{1}{\overline{\lim}_{n \rightarrow \infty} \frac{\lambda_{n-1}}{\lambda_n}},$$

then $[C(\Lambda), C(\mu)]_\infty = D_\mu w_\infty(\Lambda)$.

ii) *If $\mu \in \Gamma$, then $A \in ([C_1, C(\mu)]_\infty, s_\eta)$ if and only if*

$$\sum_{i=1}^{\infty} 2^i \max_{2^i \leq m \leq 2^{i+1}-1} |a_{nm}| \mu_m = \eta_n O(1) \quad (n \rightarrow \infty).$$

PROOF. i) We have $\Delta_\rho = D_{1/\mu} \Delta D_\mu$ with $\rho_n = \mu_{n-1}/\mu_n$ for all n , and $\Delta : D_\mu w_\infty(\Lambda) \mapsto D_\mu w_\infty(\Lambda)$ is bijective, so $D_\mu w_\infty(\Lambda)(\Delta) = D_\mu w_\infty(\Lambda)$. We have $X \in [C(\Lambda), C(\mu)]_\infty$ if and only if $C(\mu)X \in w_\infty(\Lambda)$, that is

$$X \in C(\mu)^{-1} w_\infty(\Lambda);$$

and since $C(\mu)^{-1} = \Delta(\mu) = \Delta D_\mu$, we have $X \in \Delta D_\mu w_\infty(\Lambda) = D_\mu w_\infty(\Lambda)$. Thus we have shown Part i).

ii) Here $\lambda_n = n$ for all n , so $\overline{\lim}_{n \rightarrow \infty} (\lambda_{n-1}/\lambda_n) = 1$ and condition (3.7) is satisfied, since $\mu \in \Gamma$. By Part i), we have $[C_1, C(\mu)]_\infty = D_\mu w_\infty$ and $A \in ([C_1, C(\mu)]_\infty, s_\eta)$ if and only if $D_{1/\eta} A D_\mu \in (w_\infty(\Lambda), \ell_\infty)$. We conclude, using the characterization of $(w_\infty(\Lambda), \ell_\infty)$, given in [11, Remark, p. 33]. $\square$

Now we write $\Delta_\rho(\mu) = \Delta_\rho D_\mu$ and

$$[C(\Lambda), \Delta_\rho(\mu)]_\infty = \{X \in \omega : C(\Lambda)(|\Delta_\rho(\mu)X|) \in \ell_\infty\}.$$

Then we have

$$[C(\Lambda), \Delta_\rho(\mu)]_\infty = \left\{ X \in \omega : \sup_n \left(\frac{1}{\lambda_n} \sum_{k=1}^n |\mu_k x_k - \mu_{k-1} \rho_{k-1} x_{k-1}| \right) < \infty \right\}.$$

We obtain the following result.

Proposition 3.18. *Let $\Lambda, \mu, \rho \in U^+$.*

- i) *We assume that (3.7) holds. Then $[C(\Lambda), \Delta_\rho(\mu)]_\infty = D_{1/\mu} w_\infty(\Lambda)$;*
- ii) *$A \in ([C(\Lambda), \Delta_\rho(\mu)]_\infty, s_\eta)$ if and only if*

$$\sum_{i=1}^{\infty} 2^i \max_{2^i \leq m \leq 2^{i+1}-1} \frac{|a_{nm}|}{\mu_n} = \eta_n O(1) \quad (n \rightarrow \infty).$$

PROOF. Here $C(\Lambda)(|\Delta_\rho(\mu)X|) \in \ell_\infty$ if and only if $D_\mu X \in w_\infty(\Lambda)(\Delta_\rho)$ and, by (3.7), we get $w_\infty(\Lambda)(\Delta_\rho) = w_\infty(\Lambda)$.

(ii) We obtain Part ii), reasoning as in Proposition 3.17 ii). $\square$

4. On an infinite tridiagonal matrix considered as an operator from $w_\infty(\Lambda)$ into itself and applications

In this section, we give some results on the infinite tridiagonal matrix $M(\gamma, a, \eta)$, considered as an operator from χ into $D_a \chi$ where χ is any of the sets $w_\infty(\Lambda)$, $s_\Lambda, s_\Lambda^0, s_\Lambda^{(c)}$, or $\ell_p(\Lambda)$.

Note that the previous results on Δ_ρ^+ and Δ_ρ cannot be considered as a consequence of the next ones.

Let $\gamma = (\gamma_n)_{n \geq 1}$, $\eta = (\eta_n)_{n \geq 1}$ and $a = (a_n)_{n \geq 1}$ be given sequences, and $a \in U$. We consider the infinite tridiagonal matrix

$$M(\gamma, a, \eta) = \begin{pmatrix} a_1 & \eta_1 & & & \\ \gamma_1 & a_2 & \eta_2 & & \mathbf{O} \\ & \cdot & \cdot & \cdot & \\ \mathbf{O} & & \gamma_{n-1} & a_n & \eta_n \\ & & & \cdot & \cdot \end{pmatrix}.$$

In the following we use the sets

$$\ell_p = \left\{ X = (x_n)_{n \geq 1} : \sum_{n=1}^{\infty} |x_n|^p < \infty \right\} \quad \text{for } 1 \leq p < \infty$$

and $\ell_p(\tau) = D_\tau \ell_p$ for $\tau \in U^+$.

We notice that $D_a s_\tau = D_{|a|} s_\tau = s_{|a|\tau}$, $D_a s_\tau^0 = D_{|a|} s_\tau^0 = s_{|a|\tau}^0$ and $D_a \ell_p(\tau) = \ell_p(|a|\tau)$ for $\tau \in U^+$.

We suppose that $K_1 = \sup_{n \geq 2} |\gamma_{n-1}/a_n| < 1$ and $K_2 = \sup_n |\eta_n/a_n| > 0$.

The following theorem holds.

Theorem 4.1. *Let $\Lambda \in U^+$ be an exponentially bounded sequence and assume that*

$$\sup_n \left(\frac{\lambda_{n+1}}{\lambda_n} \right) < \frac{1 - K_1}{K_2}. \quad (4.1)$$

Then, for any given $B \in D_a w_\infty(\Lambda)$, the equation $M(\gamma, a, \eta)X = B$ has a unique solution in $w_\infty(\Lambda)$ given by $X = \sum_{i=0}^{\infty} (I - M(\gamma, a, \eta))^i B$.

PROOF. We consider $D_{1/a} M(\gamma, a, \eta) = M(\zeta, e, \rho)$ with $\zeta_{n-1} = \gamma_{n-1}/a_n$ for $n \geq 2$ and $\rho_n = \eta_n/a_n$ for all n . Then we have

$$M(\zeta, e, \rho) = M(\zeta, \rho) = \begin{pmatrix} 1 & \rho_1 & & & \\ \zeta_1 & 1 & \rho_2 & & \mathbf{0} \\ & \cdot & \cdot & \cdot & \\ \mathbf{0} & & \zeta_{n-1} & 1 & \rho_n \\ & & & \cdot & \cdot \end{pmatrix}.$$

(Note that $M(\zeta, \rho) = (\Delta_{-2\zeta} + \Delta_{-2\rho}^+)/2$). Here we have

$$(I - M(\zeta, \rho))X = -(\rho_1 x_2, \zeta_1 x_1 + \rho_2 x_3, \dots, \zeta_{n-1} x_{n-1} + \rho_n x_{n+1}, \dots)^T.$$

Then

$$\|(I - M(\zeta, \rho))X\|_{w_\infty(\Lambda)}^- = \sup_n \left(\frac{1}{\lambda_n} \sum_{k=1}^n |\zeta_{k-1} x_{k-1} + \rho_k x_{k+1}| \right)$$

with $x_0 = 0$ and

$$\begin{aligned} \frac{1}{\lambda_n} \sum_{k=1}^n |\zeta_{k-1} x_{k-1} + \rho_k x_{k+1}| &\leq \frac{\lambda_{n-1}}{\lambda_n} \sup_{k \leq n-1} |\zeta_k| \frac{1}{\lambda_{n-1}} \sum_{k=1}^{n-1} |x_k| \\ + \frac{\lambda_{n+1}}{\lambda_n} \sup_{k \leq n} |\rho_k| \frac{1}{\lambda_{n+1}} \sum_{k=2}^{n+1} |x_k| &\leq \left[\sup_n \left(\frac{\lambda_{n-1}}{\lambda_n} \right) K_1 + \sup_n \left(\frac{\lambda_{n+1}}{\lambda_n} \right) K_2 \right] \|X\|_{w_\infty(\Lambda)}^-. \end{aligned}$$

Since Λ is a non-decreasing sequence, $\sup_n(\lambda_{n-1}/\lambda_n) \leq 1$, and using (4.1), we easily conclude that

$$\begin{aligned} \|I - M(\zeta, \rho)\|_{(w_\infty(\Lambda), w_\infty(\Lambda))}^- &= \sup_{X \neq 0} \frac{\|(I - M(\zeta, \rho))X\|_{w_\infty(\Lambda)}^-}{\|X\|_{w_\infty(\Lambda)}^-} \\ &\leq K_1 + \sup_n \left(\frac{\lambda_{n+1}}{\lambda_n} \right) K_2 < 1. \end{aligned}$$

Finally, for any given $B \in w_\infty(\Lambda)$, the equation $M(\gamma, a, \eta)X = B$ is equivalent to $M(\zeta, \rho)X = D_{1/a}B$. So, for any given $B \in D_a w_\infty(\Lambda)$, the last equation has a unique solution in $w_\infty(\Lambda)$. $\square$

Corollary 4.2. *Let $\lambda \in U^+$ be an exponentially bounded sequence and assume that (4.1) holds. Then*

- i) $M(\gamma, a, \eta)$ is bijective from $w_\infty(\Lambda)$ to $D_a w_\infty(\Lambda)$ and $M(\gamma, a, \eta)^{-1} \in (D_a w_\infty(\Lambda), w_\infty(\Lambda))$;
- ii) $M(\gamma, a, \eta)$ is bijective from s_Λ into $s_{|a|\Lambda}$ and $M(\gamma, a, \eta)^{-1} \in (s_{|a|\Lambda}, s_\Lambda)$;
- iii) $M(\gamma, a, \eta)$ is bijective from s_Λ° into $s_{|a|\Lambda}^\circ$ and $M(\gamma, a, \eta)^{-1} \in (s_{|a|\Lambda}^\circ, s_\Lambda^\circ)$;
- iv) for $a \in U^+$, the condition $\lim_{n \rightarrow \infty} (\gamma_{n-1}\lambda_{n-1} + \eta_n\lambda_{n+1})\lambda_n^{-1}a_n^{-1} = l \neq 0$, implies that $M(\gamma, a, \eta)$ is bijective from $s_\Lambda^{(c)}$ into $s_{a\Lambda}^{(c)}$, and $M(\gamma, a, \eta)^{-1} \in (s_{a\Lambda}^{(c)}, s_\Lambda^{(c)})$.
- v) Let $p \geq 1$ be a real. If $\tilde{K}_{p,\Lambda} = K_1 + K'_2 < 1$ with $K_1 = \sup_n(|\gamma_{n-1}/a_n|)$ and $K'_2 = \sup_n(|\eta_n/a_n|\lambda_{n+1}/\lambda_n)$, then $M(\gamma, a, \eta)$ is bijective from $\ell_p(\Lambda)$ into $\ell_p(|a|\Lambda)$, and $M(\gamma, a, \eta)^{-1} \in (\ell_p(|a|\Lambda), \ell_p(\Lambda))$.

PROOF. By [5, Theorem 6.1] 4.1 and Condition (4.1), we have

$$\xi(\Lambda) = \sup_{n \geq 1} \left[\frac{1}{|a_n|} \left(|\gamma_{n-1}| \frac{\lambda_{n-1}}{\lambda_n} + |\eta_n| \frac{\lambda_{n+1}}{\lambda_n} \right) \right] < 1,$$

and $\tilde{K}_{p,\Lambda} = K_1 + K'_2 < 1$, since

$$\xi(\Lambda) \leq \tilde{K}_{p,\Lambda} \leq K_1 + \sup_n \left(\frac{\lambda_{n+1}}{\lambda_n} \right) K_2 < 1. \quad \square$$

We can also explicitly calculate the solution of an infinite tridiagonal system when $\gamma_n = \gamma$ and $\eta_n = \eta$ for all n . Indeed the next result was shown in [6], where

$$M(\zeta, \rho) = M(\zeta, e, \rho).$$

Proposition 4.3. *Let ζ, ρ be positive reals with $0 < \zeta + \rho < 1$. Then*

- i) $M(\zeta, \rho) : X \mapsto M(\zeta, \rho)X$ is bijective from χ into itself, for $\chi \in \{s_1, c_0, c\}$.
- ii) a) Let χ be any of the sets s_1 , or c_0 , or c , and put

$$u = \left(1 - \sqrt{1 - 4\zeta\rho}\right) / 2\zeta \quad \text{and} \quad v = \left(1 - \sqrt{1 - 4\zeta\rho}\right) / 2\zeta.$$

Then, for any given $B \in \chi$, the equation $M(\zeta, \rho)X = B$ has a unique solution $X^\circ = (x_n^\circ)_{n \geq 1}$ in χ given by

$$x_n^\circ = \left(\frac{uv + 1}{uv - 1}\right) (-1)^n v^n \sum_{m=1}^{\infty} [1 - (uv)^{-l}] (-1)^m u^m b_m \quad \text{for all } n,$$

with $l = \min\{n, m\}$.

- b) The inverse $[M(\zeta, \rho)]^{-1} = (a'_{nm})_{n, m \geq 1}$ is given by

$$a'_{nm} = \left(\frac{uv + 1}{uv - 1}\right) (-v)^{n-m} [(uv)^l - 1] \quad \text{for all } n, m \geq 1 \text{ and } l = \min\{n, m\}.$$

References

- [1] I. J. MADDOX, On Kuttner's theorem, *J. London Math. Soc.* **43** (1968), 285–290.
- [2] B. DE MALAFOSSE, Sets of sequences that are strongly τ -bounded and matrix transformations between these sets, *Demonstr. Math.* **36** (2003), 155–171.
- [3] B. DE MALAFOSSE, On some BK space, *Int. J. Math. Math. Sci.* **28** (2003), 1783–1801.
- [4] B. DE MALAFOSSE, On the set of sequences that are strongly α -bounded and α -convergent to naught with index p , *Semin. Mat. Univ. Politec. Torino* **61** (2003), 13–32.
- [5] B. DE MALAFOSSE, Calculations on some sequence spaces, *Int. J. Math. Math. Sci.* **31** (2004), 1653–1670.
- [6] B. DE MALAFOSSE, The Banach algebra $\mathcal{B}(X)$, where X is a BK space and applications, *Mat. Vesnik* **57**, (2005), 41–60.
- [7] B. DE MALAFOSSE and E. MALKOWSKY, Matrix transformations in the sets $\chi(\overline{N}_p \overline{N}_q)$ where χ is in the form s_ξ , or s_ξ° , or $s_\xi^{(c)}$, *Filomat* **17** (2003), 85–106.
- [8] B. DE MALAFOSSE and E. MALKOWSKY, Sequence spaces and inverse of an infinite matrix, *Rend. del Circ. Mat. di Palermo, Serie II.* **51** (2002), 277–294.
- [9] E. MALKOWSKY, The continuous duals of the spaces $c_0(\Lambda)$ and $c(\Lambda)$ for exponentially bounded sequences Λ , *Acta Sci. Math. (Szeged)* **61** (1995), 241–250.
- [10] E. MALKOWSKY, Banach algebras of matrix transformations between spaces of strongly bounded and summable sequences, *Adv. Dyn. Syst. Appl.* **6** (2011), 91–109.
- [11] E. MALKOWSKY and V. RAKOČEVIĆ, The measure of noncompactness of linear operators between spaces of strongly C_1 summable and bounded sequences, *Acta Math. Hungar.* **89** (2000), 29–45.

On the Banach algebra $(w_\infty(\Lambda), w_\infty(\Lambda))$ and applications to the solvability... 217

- [12] A. WILANSKY, Summability through Functional Analysis, Mathematics Studies, **85**, North-Holland, Amsterdam, 1984.

BRUNO DE MALAFOSSE
I.U.T UNIVERSITÉ DU HAVRE
76600 LE HAVRE
FRANCE

E-mail: bdemalaf@wanadoo.fr

EBERHARD MALKOWSKY
DEPARTMENT OF MATHEMATICS
UNIVERSITY OF GIESSEN
ARNDTSTRASSE 2
D-35392 GIESSEN
GERMANY
C/O DEPARTMENT OF MATHEMATICS
FACULTY OF ARTS AND SCIENCES
FATİH UNIVERSITY
34500 BÜYÜKÇEKMECE
İSTANBUL
TURKEY

E-mail: Eberhard.Malkowsky@Math.uni-giessen.de,
ema@BankerInter.net

(Received July 3, 2013; revised October 23, 2013)