*Publ. Math. Debrecen*
**93/3-4** (2018), 413–424
DOI: 10.5486/PMD.2018.8130

# Random power series near the endpoint of the convergence interval

By BALÁZS MAGA (Budapest) and PÉTER MAGA (Budapest)

**Abstract.** In this paper, we are going to consider power series

$$\sum_{n=1}^{\infty} a_n x^n,$$

where the coefficients $a_n$ are chosen independently at random from a finite set with uniform distribution. We prove that if the expected value of the coefficients is 0, then

$$\limsup_{x\to 1-} \sum_{n=1}^{\infty} a_n x^n = \infty, \qquad \liminf_{x\to 1-} \sum_{n=1}^{\infty} a_n x^n = -\infty,$$

with probability 1. We investigate the analogous question in terms of Baire categories.

## 1. Introduction

In complex analysis, the behaviour of random power series near the radius of convergence has been thorougly examined, partly due to the following classical problem: if we consider the Taylor series $f(z) = \sum_{n=0}^{\infty} a_n z^n$, what properties

*Mathematics Subject Classification:* Primary: 60F20; Secondary: 11A63, 54E52.
*Key words and phrases:* real random power series, boundary behaviour, zero-one laws, residuality.
The first author was supported by the ÚNKP-17-2 New National Excellence Program of the Hungarian Ministry of Human Capacities and by the Hungarian National Research, Development and Innovation Office (NKFIH), Grant No. 124003. The second author was supported by the Postdoctoral Fellowship of the Hungarian Academy of Sciences and by NKFIH grants NK 104183 and ERC_HU_15 118946.

of the sequence $(a_n)_{n=0}^{\infty}$ imply that $f$ has its radius of convergence as a natural boundary, that is all of the points on its radius of convergence are singular? It turned out that random power series form a large family of such functions: it was proven in [6] that if $f$ has a finite radius of convergence and $(a_n)_{n=0}^{\infty}$ are independent, identically distributed random variables with uniform distribution on $\{|z| = 1\}$, then for almost every choice, $f$ has a natural boundary on the radius of convergence. Later, somewhat stronger and more specific results were obtained, even in the recent years (see, e.g., [1]).

These theorems showed that random power series in the complex plane tend to behave rather chaotically near the radius of convergence. In this paper, we investigate a similar question on the real line, motivated by a problem raised in [2]. Although the results are somewhat natural and are easy to formulate, we did not manage to find them in the literature.

Let $D = \{d_1, \dots, d_k\}$ be a finite set of real numbers. Then we may consider the random power series with coefficients from $D$, i.e.,

$$f(x) = \sum_{n=1}^{\infty} a_n x^n,$$

where each $a_n$ equals $d_j$ (for $1 \leqslant j \leqslant k$) with probability $1/k$, independently in $n$. To exclude trivialities, assume from now on that $k \geqslant 2$.[1]

To make this more rigorous, define the probability space $(D, \mathcal{A}_n, \mathrm{P}_n)$, where $D$ is the fixed set above, $\mathcal{A}_n$ is the discrete topology on $D$, and for each $D' \subseteq D$,

$$\mathrm{P}_n(D') = \#D'/\#D = \#D'/k,$$

with $\#$ standing for the cardinality.

Then set $(\Omega, \mathcal{A}, \mathrm{P})$ for the product probability space, i.e., $\Omega = \prod_{n\in\mathbf{N}} D$, $\mathcal{A}$ is the set of Borel sets of $\Omega$ (in the product topology $\prod_{n\in\mathbf{N}} \mathcal{A}_n$), $\mathrm{P} = \prod_{n\in\mathbf{N}} \mathrm{P}_n$.

We will denote a general element of $\Omega$ by $(a_n)$, and by $a_n$ its $n$-th coordinate (i.e., $a_n \in D$, $(a_n) \in \Omega$). To any $(a_n) \in \Omega$, we may associate the power series

$$f_{(a_n)}(x) = \sum_{n=1}^{\infty} a_n x^n.$$

In most cases below, there will be a single sequence $(a_n)$ and a resulting power series $f_{(a_n)}$, therefore we write simply $f$ in place of $f_{(a_n)}$. Of course, when there is any chance for confusion, we return to the longer (and less loose) notation.

[1] By a slight change of notation, we start the power series with the first-order term, in order to index the random variables by $\mathbf{N}$.

It is easy to see that the convergence radius of $f(x)$ is 1 for almost all coefficient sequences $(a_n)$ (except for the trivial case $D = \{0\}$, which is already excluded by our assumption $k \geqslant 2$). In this paper, we investigate the behaviour of $f$, as $x$ tends to 1 from below. To this end, we introduce the following notation:

$$L_+((a_n)) = \limsup_{x\to 1-} a_n x^n, \qquad L_-((a_n)) = \liminf_{x\to 1-} a_n x^n.$$

Then our result is the following.

**Theorem 1.** *If $\sum_{d\in D} d = 0$, then*

$$\mathrm{P}((a_n) \in \Omega : L_+((a_n)) = \infty,\ L_-((a_n)) = -\infty) = 1.$$

*Remark 1.* If $\sum_{d\in D} d > 0$, then

$$\mathrm{P}((a_n) \in \Omega : L_+((a_n)) = L_-((a_n)) = \infty) = 1;$$

while if $\sum_{d\in D} d < 0$, then

$$\mathrm{P}((a_n) \in \Omega : L_+((a_n)) = L_-((a_n)) = -\infty) = 1.$$

This can be seen via Abel's theorem combined with the law of large numbers. Indeed, assume $\sum_{d\in D} d > 0$, then by the law of large numbers, $\sum_{n=1}^{\infty} a_n = \infty$ for almost all $(a_n) \in \Omega$, and if $\sum_{n=1}^{\infty} a_n = \infty$, then $\lim_{x\to 1-} \sum_{n=1}^{\infty} a_n x^n = \infty$, as it follows by a simple partial summation, see, e.g., [5, Theorem 8.2].

In Section 4, we investigate the same properties of generic power series in the Baire categorial sense (see [4, pp. 40–41]). The corresponding statements are summarized as follows.

**Theorem 2.** *If each element of $D$ is nonnegative (resp. nonpositive), then*

$$\{(a_n) \in \Omega : L_+((a_n)) = L_-((a_n)) = \infty\}$$
$$\text{(resp. } \{(a_n) \in \Omega : L_+((a_n)) = L_-((a_n)) = -\infty\})$$

*is residual.*

*If $D$ contains positive and negative elements simultaneously, then*

$$\{(a_n) \in \Omega : L_+((a_n)) = \infty,\ L_-((a_n)) = -\infty\}$$

*is residual.*

Now Theorems 1 and 2 have the following simple consequence via Bolzano's theorem on continuous functions, answering a question in [2].

**Corollary 1.** *If $\#D \geqslant 2$, and $\sum_{d\in D} d = 0$, then for almost all and residually many sequences $(a_n) \in \Omega$, the following holds. For any real number $y$, there are infinitely many numbers $0 < x < 1$ satisfying*

$$y = \sum_{n=1}^{\infty} a_n x^n.$$

For the sake of completeness, we record that using the facts that the function $\Omega \to \mathbf{R}$, $(a_n) \mapsto f_{(a_n)}(x)$ is continuous for all $x \in [0,1)$ and that the supremum over countably many continuous functions is measurable, one may easily see that the functions $L_+$ and $L_-$ are random variables, i.e., it makes sense to speak about the probabilities in Theorem 1 and Remark 1.

## 2. Preparatory results

The main goal of this section is to prove the following.

**Proposition 1.** *Under the conditions of the introduction, we have*

$$\mathrm{P}((a_n) \in \Omega : L_+((a_n)), L_-((a_n)) \in \{\pm\infty\}) = 1.$$

Of course, $L_-((a_n)) \leqslant L_+((a_n))$, so in fact three possibilities are covered in Proposition 1. Then the statements in Theorem 1 and Remark 1 can be thought of as identifying which one of these three holds almost surely.

First of all, define the following Borel measures on $\mathbf{R}$:

$$\mu_+(B) = \mathrm{P}\left(L_+^{-1}(B)\right), \qquad \mu_-(B) = \mathrm{P}\left(L_-^{-1}(B)\right).$$

In other words, these are the distributions of the random variables $L_+$ and $L_-$, in particular, both of them are finite. One may easily see that Proposition 1 is equivalent to the fact that both $\mu_+$ and $\mu_-$ are the constant 0 measures on $\mathbf{R}$.

We start with a concept of combinatorial nature. For any $N \in \mathbf{N}$, define the function $g_N^\sharp$ between two subsets of $D^N$ satisfying the following conditions:

(i) $g_N^\sharp$ is a bijection between its domain and range;

(ii) if $g_N^\sharp((a_1, \dots, a_N)) = (b_1, \dots, b_N)$, then

$$\sum_{n=1}^{N} b_n = (d_2 - d_1) + \sum_{n=1}^{N} a_n.$$

It is easy to see that, in general, we cannot define $g_N^\sharp$ on the whole set $D^N$. However, as the following lemma points it out, it can be defined on a considerably large subset.

**Lemma 1.** *The map $g_N^\sharp$ can be defined such that*

$$\#\operatorname{dom} g_N^\sharp = k^N(1-o(1)),$$

*as $N \to \infty$.*

Proof. First of all, split up the set $D^N$ as follows. Take any $0 \leqslant l \leqslant N$, and any numbers $1 \leqslant c_1 < \cdots < c_l \leqslant N$. Set then $\mathbf{c} = \{c_1, \dots, c_l\}$ and $\mathbf{c}' = \{1, \dots, N\} \setminus \{c_1, \dots, c_l\}$. Further, let $\mathbf{s} : \mathbf{c}' \to D \setminus \{d_1, d_2\}$. Attached to this data, set

$$D^N_{l,\mathbf{c},\mathbf{s}} = \{(a_1, \dots, a_N) \in D^N \mid \forall c_j \in \mathbf{c} : a_{c_j} \in \{d_1, d_2\} \text{ and } \forall c' \in \mathbf{c}' : a_{c'} = \mathbf{s}(c')\},$$

i.e., $D^N_{l,\mathbf{c},\mathbf{s}}$ stands for those sequences which contain $d_1$'s and $d_2$'s in positions indexed by $\mathbf{c}$, while outside of $\mathbf{c}$, there is a fixed sequence $\mathbf{s}$ made of coefficients other than $d_1, d_2$.

Decompose $D^N_{l,\mathbf{c},\mathbf{s}}$ as

$$D^N_{l,\mathbf{c},\mathbf{s}} = \bigcup_{l_1=0}^{l} D^N_{l,l_1,\mathbf{c},\mathbf{s}},$$

where $D^N_{l,l_1,\mathbf{c},\mathbf{s}}$ is the subset of $D^N_{l,\mathbf{c},\mathbf{s}}$ which consists of sequences containing exactly $l_1$ many $d_1$'s. Obviously, $g_N^\sharp$ can be defined on a set of size

$$\#D^N_{l,\mathbf{c},\mathbf{s}} - \sum_{l_1=0}^{l} \max(0, \#D^N_{l,l_1,\mathbf{c},\mathbf{s}} - \#D^N_{l,l_1-1,\mathbf{c},\mathbf{s}}),$$

namely, $g_N^\sharp$ maps a sequence in $D^N_{l,\mathbf{c},\mathbf{s}}$ to another one which contains one more copy of $d_2$ and one less copy of $d_1$. Clearly, $\#D^N_{l,\mathbf{c},\mathbf{s}} = 2^l$ and $\#D^N_{l,l_1,\mathbf{c},\mathbf{s}} = \binom{l}{l_1}$. We claim that, for any fixed $\varepsilon > 0$,

$$\sum_{l_1=0}^{l} \max(0, \#D^N_{l,l_1,\mathbf{c},\mathbf{s}} - \#D^N_{l,l_1-1,\mathbf{c},\mathbf{s}}) \leqslant 2\varepsilon 2^l + o(2^l), \tag{1}$$

as $l \to \infty$. Split this summation up according to $l_1 \geqslant l(1/2-\varepsilon)$ and $l_1 < l(1/2-\varepsilon)$. As for the latter, even

$$\sum_{l_1 < l(1/2-\varepsilon)} \#D^N_{l,l_1,\mathbf{c},\mathbf{s}} = o(2^l),$$

as $l \to \infty$, follows simply from Chebyshev's inequality applied to the random walk of length $l$. Therefore, apart from $o(2^l)$ sequences in $D^N_{l,\mathbf{c},\mathbf{s}}$, we have $l_1 \geqslant l(1/2-\varepsilon)$. In this part of the summation,

$$\sum_{l_1 \geqslant l(1/2-\varepsilon)} \max(0, \#D^N_{l,l_1,\mathbf{c},\mathbf{s}} - \#D^N_{l,l_1-1,\mathbf{c},\mathbf{s}})$$
$$\leqslant 2^l \max_{l(1/2-\varepsilon) \leqslant l_1 \leqslant l/2} \left(1 - \frac{\#D^N_{l,l_1-1,\mathbf{c},\mathbf{s}}}{\#D^N_{l,l_1,\mathbf{c},\mathbf{s}}}\right) \leqslant 2^l(2\varepsilon + o(1)),$$

hence (1) is established.

It is easy to see that for any fixed $L$, as $N \to \infty$,

$$\sum_{l \leqslant L,\mathbf{c},\mathbf{s}} \#D^N_{l,\mathbf{c},\mathbf{s}} = o(k^N), \qquad \sum_{l > L,\mathbf{c},\mathbf{s}} \#D^N_{l,\mathbf{c},\mathbf{s}} = k^N - o(k^N).$$

Now let $\delta > 0$ be arbitrary. Choose $L$ such that (1) can be continued as $2\varepsilon 2^l + o(2^l) < 3\varepsilon 2^l$ for any $N \geqslant l > L$. Then, with this fixed $L$, if $N$ is large enough, at least $(1-\delta)k^N$ sequences in $D^N$ satisfies $l > L$ (with $l$ standing for the total number of $d_1$'s and $d_2$'s). This altogether yields that $g^\sharp_N$ can be defined on a set of size at least $k^N(1-\delta)(1-3\varepsilon)$. Since $\delta > 0$ and $\varepsilon > 0$ are arbitrary, this completes the proof. □

Similarly, we can define the functions $g^\flat_N$ which map sequences $(a_1, \dots, a_N)$ to $(b_1, \dots, b_N)$ such that

$$\sum_{n=1}^N b_n = (d_1 - d_2) + \sum_{n=1}^N a_n,$$

and $g^\flat_N$ are bijections between their domain and range. The same argument as that in the proof of Lemma 1 gives

$$\#\operatorname{dom} g^\flat_N = k^N(1 - o(1)),$$

as $N \to \infty$, for a well-chosen function $g^\flat_N$.

From now on, fix two sequences of such functions $g^\sharp_N$ and $g^\flat_N$ (with domains of size $k^N(1-o(1))$).

**Lemma 2.** *Both $\mu_+$ and $\mu_-$ are invariant under translations by $d_2 - d_1$, i.e., for any Borel set $B \subseteq \mathbf{R}$,*

$$\mu_+(B + d_2 - d_1) = \mu_+(B), \qquad \mu_-(B + d_2 - d_1) = \mu_-(B).$$

PROOF. Let $\mu = \mu_+$, the argument for $\mu_-$ is literally the same, writing lim inf's in place of lim sup's and $L_-$'s in place of $L_+$'s. Fix first $\varepsilon > 0$.

On certain sequences $(a_n) \in \Omega$, apply the following sequence of operations: if $(a_1, \dots, a_N) \in \operatorname{dom} g_N^\sharp$, then let

$$G_N^\sharp((a_n)) = (b_n),$$

where $g_N^\sharp((a_1, \dots, a_N)) = (b_1, \dots, b_N)$, and $b_n = a_n$ for all $n > N$. Now observe that

$$\begin{aligned}\limsup_{x\to 1-} \sum_{n=1}^{\infty} b_n x^n &= \sum_{n=1}^{N} b_n + \limsup_{x\to 1-} \sum_{n=N+1}^{\infty} b_n x^n \\ &= d_2 - d_1 + \sum_{n=1}^{N} a_n + \limsup_{x\to 1-} \sum_{n=N+1}^{\infty} a_n x^n \\ &= d_2 - d_1 + \limsup_{x\to 1-} \sum_{n=1}^{\infty} a_n x^n.\end{aligned}$$

This altogether means that if $(a_n) \in \operatorname{dom} G_N^\sharp \cap L_+^{-1}(B)$, then $G_N^\sharp((a_n)) \in L_+^{-1}(B + d_2 - d_1)$ for any Borel set $B \subseteq \mathbf{R}$.

By Lemma 1, if $N$ is large enough, $\mathrm{P}(\operatorname{dom} G_N^\sharp) \geqslant 1-\varepsilon$, implying $\mathrm{P}(\operatorname{dom} G_N^\sharp \cap L_+^{-1}(B)) \geqslant \mu(B) - \varepsilon$. Then, using the simple fact that $G_N^\sharp$ preserves P on its domain, we see

$$\mu(B + d_2 - d_1) = \mathrm{P}(L_+^{-1}(B + d_2 - d_1)) \geqslant \mathrm{P}(G_N^\sharp(\operatorname{dom} G_N^\sharp \cap L_+^{-1}(B))) \geqslant \mu(B) - \varepsilon.$$

Since this holds for all $\varepsilon > 0$, we obtain

$$\mu(B + d_2 - d_1) \geqslant \mu(B).$$

The same way we obtain $\mu(B - d_2 + d_1) \geqslant \mu(B)$ (this time applying $g_N^\flat$), which yields the statement. □

It is well-known (and simple) that there are no nontrivial finite Borel measures on $\mathbf{R}$ which are invariant under a nontrivial translation, implying $\mu_+(\mathbf{R}) = \mu_-(\mathbf{R}) = 0$. As it was mentioned in the introduction of this section, this completes the proof of Proposition 1.

Another tool we need is a standard application of Kolmogorov's 0-1 law.

**Lemma 3.** *We have*

$$\mathrm{P}((a_n) \in \Omega : L_+((a_n)) = L_-((a_n)) \in \{\pm\infty\}) \in \{0, 1\}.$$

PROOF. It is easy to see that the events

$$A_{\pm} = \left\{ (a_n) \in \Omega : \lim_{x \to 1-} \sum_{n=1}^{\infty} a_n x^n = \pm\infty \right\}$$

are tail events in the sense of [3, Section 16.3], since

$$\lim_{x \to 1-} \sum_{n=1}^{\infty} a_n x^n = \pm\infty \quad \text{if and only if} \quad \lim_{x \to 1-} \sum_{n=N+1}^{\infty} a_n x^n = \pm\infty$$

holds for any $N \in \mathbf{N}$, implying that the events $A_{\pm}$ are independent of the first few coefficients $a_1, \dots, a_N$. Tail events have probability 0 or 1 by Kolmogorov's 0-1 law, see [3, Theorem 16.3 B]. □

## 3. Proof of Theorem 1

In this section, we are going to prove Theorem 1, so assume $\sum_{j=1}^{k} d_j = 0$. We introduce the following permutation $p$ on $D$: $p(d_j) = d_{j+1}$ for $1 \leqslant j \leqslant k-1$, and $p(d_k) = d_1$. This gives rise to a permutation $\mathbf{p}$ on $\Omega$: $\mathbf{p}((a_n)) = (b_n)$, where $b_n = p(a_n)$ for each $n \in \mathbf{N}$. Now for any $0 < x < 1$, by absolute convergence,

$$\sum_{j=0}^{k-1} f_{\mathbf{p}^j((a_n))}(x) = \sum_{j=0}^{k-1} \sum_{n=1}^{\infty} p^j(a_n) x^n = \sum_{n=1}^{\infty} \left( \sum_{j=0}^{k-1} p^j(a_n) \right) x^n = \sum_{n=1}^{\infty} \left( \sum_{j=1}^{k} d_j \right) x^n = 0.$$

Consequently, for any $(a_n) \in \Omega$ and any $0 < x < 1$, among the numbers $f_{(a_n)}(x), f_{\mathbf{p}((a_n))}(x), \dots, f_{\mathbf{p}^{k-1}((a_n))}(x)$ there is at least one nonnegative and at least one nonpositive one. By the pigeon-hole principle, this means that for any $(a_n) \in \Omega$, there exist $0 \leqslant j_+, j_- \leqslant k-1$ such that

$$L_+(\mathbf{p}^{j_+}(a_n)) \geqslant 0, \qquad L_-(\mathbf{p}^{j_-}(a_n)) \leqslant 0.$$

This yields that

$$\bigcap_{j=0}^{k-1} ((a_n) \in \Omega : L_+(\mathbf{p}^j((a_n))) = L_-(\mathbf{p}^j((a_n))) = \infty) = \emptyset,$$

$$\bigcap_{j=0}^{k-1} ((a_n) \in \Omega : L_+(\mathbf{p}^j((a_n))) = L_-(\mathbf{p}^j((a_n))) = -\infty) = \emptyset.$$

Also, it is easy to see that $\mathbf{p}$ is P-preserving, which implies that for some $0 \leqslant a_+, a_- \leqslant 1$,

$$\mathrm{P}((a_n) \in \Omega : L_+(\mathbf{p}^j((a_n))) = L_-(\mathbf{p}^j((a_n))) = \infty) = a_+, \text{ for all } 0 \leqslant j \leqslant k-1,$$
$$\mathrm{P}((a_n) \in \Omega : L_+(\mathbf{p}^j((a_n))) = L_-(\mathbf{p}^j((a_n))) = -\infty) = a_-, \text{ for all } 0 \leqslant j \leqslant k-1.$$

The above two displays clearly imply that $a_+, a_- \leqslant 1 - 1/k$, i.e.,

$$\mathrm{P}((a_n) \in \Omega : L_+((a_n)) = L_-((a_n)) \in \{\pm\infty\}) \leqslant 1 - 1/k.$$

This, combined with Lemma 3, gives

$$\mathrm{P}((a_n) \in \Omega : L_+((a_n)) = L_-((a_n)) \in \{\pm\infty\}) = 0.$$

Now Theorem 1 follows from Proposition 1.

## 4. About residuality: the proof of Theorem 2

In this section, we prove Theorem 2. First assume that each element of $D$ is nonnegative. In this case, we have $\lim_{x\to 1-} f(x) \neq \infty$ if and only if the sequence of the coefficients contains only finitely many nonzero elements. However, the set $E$ of these sequences is of first category. Indeed, write $E = \bigcup_{m=1}^{\infty} E_m$, where $E_m$ denotes the set of sequences for which $a_n = 0$ holds for $n \geqslant m$. It suffices to see that $E_m$ is nowhere dense in $\Omega$ (for each $m \in \mathbf{N}$). Given any nonempty open set $U$, it has a nonempty open subset

$$V = \{(a_n) \in \Omega \mid a_1 = b_1, a_2 = b_2, \ldots, a_j = b_j\},$$

where $j \in \mathbf{N}$ and $b_1, \ldots, b_j \in D$. Now define

$$W = \{(a_n) \in \Omega \mid a_1 = b_1, a_2 = b_2, \ldots, a_j = b_j, a_{\max(j,m)+1} = b\},$$

where we choose $b \in D$ to be nonzero. Clearly, $W \subseteq U$ is nonempty, open, and $W \cap E_m = \emptyset$, therefore the proof of the first statement is complete (the case when each element of $D$ is nonpositive follows by symmetry).

Now let us consider the case in which $D$ contains positive and negative elements simultaneously. One can easily see that $\limsup_{x\to 1-} f(x) = \infty$ holds if and only if $\sup_{x\in(0,1)} f(x) = \infty$, and $\liminf_{x\to 1-} f(x) = -\infty$ holds if and only if $\inf_{x\in(0,1)} f(x) = -\infty$. Thus it suffices to prove that $\sup_{x\in(0,1)} f(x) \neq +\infty$ or

$\inf_{x\in(0,1)} f(x) \neq -\infty$ hold only in a set of first category. By symmetry, we can focus on the set $F$ where $\sup_{x\in(0,1)} f(x) \neq \infty$ holds. Write it as a countable union $F = \bigcup_{n=1}^{\infty} F_m$, where $F_m$ contains the sequences for which $\sup_{x\in(0,1)} f(x) \leqslant m$. It suffices to see that $F_m$ is nowhere dense in $\Omega$ (for each $m \in \mathbf{N}$).

Given any nonempty open set $U$, it has a nonempty open subset

$$V = \{(a_n) \in \Omega \mid a_1 = b_1, a_2 = b_2, \dots, a_j = b_j\},$$

where $j \in \mathbf{N}$ and $b_1, \dots, b_j \in D$. Set

$$R = \inf_{x\in(0,1)} \sum_{n=1}^{j} b_n x^n.$$

Choose an integer $M > j$ satisfying also $M > (m + 1 - R)/(\max D) + j$, then

$$R + \max D \sum_{n=j+1}^{M} 1 > m + 1.$$

Now fix $x < 1$ close enough to 1 such that

$$R + \max D \sum_{n=j+1}^{M} x^n > m + 1.$$

Then choose $N > M$ large enough such that $|\min D \sum_{n=N+1}^{\infty} x^n| < 1$. Taking

$$W = \{(a_n) \in \Omega \mid a_1 = b_1, a_2 = b_2, \dots, a_j = b_j, a_{j+1} = \cdots = a_N = \max D\},$$

we have, for $(a_n) \in W$,

$$\begin{aligned}\sum_{n=1}^{\infty} a_n x^n &\geqslant \sum_{n=1}^{j} b_n x^n + \sum_{n=j+1}^{N} \max D \cdot x^n + \sum_{n=N+1}^{\infty} \min D \cdot x^n \\ &\geqslant R + \max D \sum_{i=j+1}^{N} x^n + \min D \sum_{n=N+1}^{\infty} x^n > m + 1 - 1 = m.\end{aligned}$$

Therefore, $W \cap F_m = \emptyset$, and since $W \subseteq U$ is nonempty and open, the proof of Theorem 2 is complete.

## 5. Concluding remarks

It would be interesting to investigate the question of non-uniform distributions, i.e., when $D = \{d_1, \dots, d_k\}$ and the positive numbers $p_1, \dots, p_k$ are given such that $\sum_{j=1}^{k} p_j = 1$, and each coefficient takes the value $d_j$ with probability $p_j$ for $1 \leqslant j \leqslant k$.

Proposition 1 can be proved similarly, apart from the following subtlety. Assuming $p_1 \leqslant p_2$, take the function $g_N^\sharp$ with the same properties as above. Then the resulting function $G_N^\sharp$ in the proof of Lemma 2 does not preserve P for $p_2 > p_1$, but increases it. (Similarly, $G_N^\flat$ is P-decreasing.) All in all, although our measure $\mu = \mu_\pm$ will not be invariant any more under the translation by $d_2 - d_1$, we still have

$$\mu(B + d_2 - d_1) \geqslant \mu(B)$$

for any Borel set $B \subseteq \mathbf{R}$, and there is no such finite measure on $\mathbf{R}$ other than the trivial one.

It is also clear that the argument in Remark 1 goes through independently of whether the underlying distribution is uniform or not. However, it is clear that uniformity is heavily used in Section 3. At the moment, we do not see any obvious modification of the argument which would tell us the typical behaviour of $\lim_{x\to 1-} \sum_{n=1} a_n x^n$, where each $a_n$ takes, e.g., the value $-1$ with probability $\sqrt{2}/(1+\sqrt{2})$ and the value $\sqrt{2}$ with probability $1/(1+\sqrt{2})$, jointly independently in $n$.

Acknowledgements. We thank the referees whose careful reading and detailed comments helped us to improve the exposition.

## References

[1] J. Breuer and B. Simon, Natural boundaries and spectral theory, *Adv. Math.* **226** (2011), 4902–4920.

[2] V. Komornik, M. Pedicini and A. Pethő, Multiple common expansions in non-integer bases, *Acta Sci. Math. (Szeged)* **83** (2017), 51–60.

[3] M. Loève, Probability Theory. I. Fourth Edition, Graduate Texts in Mathematics, Vol. **45**, *Springer-Verlag, New York – Heidelberg*, 1977.

[4] J. C. Oxtoby, Measure and Category. A Survey of the Analogies Between Topological and Measure Spaces. Second Edition, Graduate Texts in Mathematics, Vol. **2**, *Springer-Verlag, New York – Berlin*, 1980.

[5] W. Rudin, Principles of Mathematical Analysis. Third Edition, International Series in Pure and Applied Mathematics, *McGraw-Hill Book Co., New York – Auckland – Düsseldorf*, 1976.

[6] H. Steinhaus, Über die Wahrscheinlichkeit dafür, daß der Konvergenzkreis einer Potenzreihe ihre natürliche Grenze ist, *Math. Z.* **31** (1930), 408–416.

BALÁZS MAGA
EÖTVÖS LORÁND UNIVERSITY
PÁZMÁNY PÉTER SÉTÁNY 1/C
BUDAPEST, H-1117
HUNGARY

*E-mail:* magab@cs.elte.hu

PÉTER MAGA
LENDÜLET AUTOMORPHIC RESEARCH GROUP
MTA ALFRÉD RÉNYI INSTITUTE
OF MATHEMATICS
POB 127
BUDAPEST H-1364
HUNGARY

*E-mail:* magapeter@gmail.com

*(Received September 12, 2017; revised April 5, 2018)*