

Interiors of continuous images of self-similar sets with overlaps

By LIFENG XI (Ningbo), KAN JIANG (Ningbo), JIALI ZHU (Ningbo)
 and QIYANG PEI (Ningbo)

Abstract. Let K be the attractor of the following iterated function system

$$\{S_1(x) = \lambda x, S_2(x) = \lambda x + c - \lambda, S_3(x) = \lambda x + 1 - \lambda\},$$

where $S_1(I) \cap S_2(I) \neq \emptyset$, $(S_1(I) \cup S_2(I)) \cap S_3(I) = \emptyset$, and $I = [0, 1]$ is the convex hull of K . Let $d_1 = \frac{1-c-\lambda}{\lambda} < \frac{1}{1-c-\lambda} = d_2$. Suppose that f is a continuous function defined on an open set $U \subset \mathbb{R}^2$. Denote the image

$$f_U(K, K) = \{f(x, y) : (x, y) \in (K \times K) \cap U\}.$$

If $\partial_x f, \partial_y f$ are continuous on U , and there is a point $(x_0, y_0) \in (K \times K) \cap U$ such that

$$\left| \frac{\partial_y f|_{(x_0, y_0)}}{\partial_x f|_{(x_0, y_0)}} \right| \in (d_1, d_2) \quad \text{or} \quad \left| \frac{\partial_x f|_{(x_0, y_0)}}{\partial_y f|_{(x_0, y_0)}} \right| \in (d_1, d_2),$$

then $f_U(K, K)$ contains an interval. As a result, we let $c = \lambda = \frac{1}{3}$, and if

$$f(x, y) = x^\alpha y^\beta (\alpha\beta \neq 0), \quad x^\alpha \pm y^\alpha (\alpha \neq 0), \quad \sin(x) \cos(y), \quad \text{or } x \sin(xy),$$

then $f_U(C, C)$ contains an interval, where C is the middle-third Cantor set.

Mathematics Subject Classification: Primary: 28A80.

Key words and phrases: fractal, self-similar set, overlap, arithmetic representation, interior.

The work is supported by the National Natural Science Foundation of China (Nos. 11831007, 11771226, 11701302, 11371329, 11471124, 11671147). The work is also supported by K. C. Wong Magna Fund in Ningbo University.

The first author is the corresponding author.

1. Introduction

Representation of real numbers is an important area in number theory. There are many ways of expressing numbers, for instance, the β -expansions ([1], [3], [6], [7], [13], [14], [15], [26]), the continued fractions ([11], [12]), the Lüroth expansions ([5]), and so on. These representations are popular in number theory. There are, however, some other approaches that can represent real numbers ([2], [24], [37], [39]). First, let us introduce some basic definitions. Given two non-empty sets $A, B \subset \mathbb{R}$, we define

$$A * B = \{x * y : x \in A, y \in B\},$$

where $*$ is $+$, $-$, $\cdot$ or $\div$ (when $*$ is $\div$, $y \neq 0$). We call $u = x * y$ an arithmetic representation in terms of A and B . Steinhaus proved that

$$C - C = \{x - y : x, y \in C\} = [-1, 1],$$

where C is the middle-third Cantor set. This result also implies that $C + C = [0, 2]$ due to the equation $C = 1 - C$. In [2], ATHREYA, REZNICK and TYSON considered the multiplication on C , and proved that

$$17/21 \leq \mathcal{L}(C \cdot C) \leq 8/9,$$

where $\mathcal{L}$ denotes the Lebesgue measure. Moreover, they also considered the division on C . Due to these two results, it is natural to ask that for the general self-similar sets K_1 and K_2 , how large $K_1 * K_2$ is in the sense of Hausdorff dimension or Lebesgue measure, where $*$ is $+$ or $\cdot$. This question was mainly addressed by PERES and SHMERKIN [31], HOCHMAN and SHMERKIN [19], who proved the following results. Let K_1 and K_2 be two self-similar sets with iterated function systems (IFS's)

$$\{f_i(x) = r_i x + a_i\}_{i=1}^n \quad \text{and} \quad \{g_j(x) = r'_j x + b_j\}_{j=1}^m,$$

respectively, if there are some r_i, r'_j such that $\log |r_i| / \log |r'_j| \notin \mathbb{Q}$, then

$$\dim_H(K_1 + K_2) = \min\{\dim_H(K_1) + \dim_H(K_2), 1\}.$$

The condition in the above result is called the irrational condition. In [34], SHMERKIN stated that

$$\dim_H(K_1 \cdot K_2) = \min\{\dim_H(K_1) + \dim_H(K_2), 1\}$$

under the irrational condition. It is natural to consider the Hausdorff dimension of $K_1 * K_2$, where $*$ = + or $\cdot$, without the irrational condition. In [22], JIANG proved that without the irrational condition, $K_1 + K_2$ is either a self-similar set or an attractor of some infinite iterated function system. Moreover, in [23], JIANG considered the sum $K_1 + K_2$ from the projectional perspective, and calculated in certain cases the Hausdorff dimension of the projection of $K_1 \times K_2$ for some special angles. If $\dim_H(K_1 + K_2) = 1$ or $\dim_H(K_1 \cdot K_2) = 1$, then one may ask whether $K_1 + K_2$ or $K_1 \cdot K_2$ contains an interval. For $K_1 + K_2$, there is a celebrated conjecture posed by PALIS [30], i.e. whether it is true (at least generically) that the arithmetic sum of dynamically defined Cantor sets either has measure zero or contains an interval. This conjecture was solved in [28]. However, for the general self-similar sets this conjecture is still open. Without the irrational condition, there are few results for $K_1 \cdot K_2$. In [38], YUKI considered the general Cantor sets, and proved under some conditions that $K_1 \cdot K_2$ contains an interval.

Suppose that $K = K_{\lambda,c}$ is the self-similar set generated by the following IFS,

$$\{S_1(x) = \lambda x, S_2(x) = \lambda x + c - \lambda, S_3(x) = \lambda x + 1 - \lambda\},$$

where $S_1(I) \cap S_2(I) \neq \emptyset$, $(S_1(I) \cup S_2(I)) \cap S_3(I) = \emptyset$, and $I = [0, 1]$ is the convex hull of K . The class $\{K_{\lambda,c}\}_{\lambda,c}$, which is an important class of self-similar sets with overlaps [21], was investigated by many scholars. It was investigated from different aspects, see [9], [10], [16], [20], [25], [29], [32], [39]. For example, HOCHMAN [20] proved the Furstenberg's conjecture which states $\dim_H K_{1/3,c} = 1$ for any $c \notin \mathbb{Q}$, Keyon discussed the necessary and sufficient condition for the open set condition satisfied on $K_{1/3,c}$, RAO and WEN [32] obtained the graph-directed construction when λ^{-1} is a P.V. number and $c \in \mathbb{Q}$.

In [39], TIAN *et al.* proved that $K \cdot K = [0, 1]$ if and only if $c \geq (1 - \lambda)^2$. In this paper, we shall analyze the continuous image of K . Let f be a continuous function defined on an open set $U \subset \mathbb{R}^2$. We call

$$f_U(K, K) = \{f(x, y) : (x, y) \in (K \times K) \cap U\}$$

the continuous image of K . For convenience, we write $f(K, K) = f_{\mathbb{R}^2}(K, K)$.

In this paper, we shall give a sufficient condition such that $f_U(K, K)$ contains an interval. In fact, whether a fractal contains an interval is a crucial problem in fractal geometry and dynamical systems. SCHIEF [33], BANDT and GRAF [4] showed the relations among the open set condition, positive Hausdorff measure and non-empty interiors. DAJANI *et al.* [8], HARE and SIDOROV [17], [18] found that the existence of an interior of a class of self-affine sets implies the existence of

the simultaneous expansions. In [27], LAGARIAS and WANG showed that for some self-affine tiles, the existence of interiors is equivalent to the positive Lebesgue measure of the tiles. More results can be found in [27], [29] and references therein.

Now we state the main results of this paper.

Theorem 1. *Let K be the attractor of the following IFS*

$$\{S_1(x) = \lambda x, S_2(x) = \lambda x + c - \lambda, S_3(x) = \lambda x + 1 - \lambda\},$$

where $S_1(I) \cap S_2(I) \neq \emptyset$, $(S_1(I) \cup S_2(I)) \cap S_3(I) = \emptyset$ and $I = [0, 1]$ is the convex hull of K . Let $d_1 = \frac{1-c-\lambda}{\lambda} < \frac{1}{1-c-\lambda} = d_2$. Suppose that f is a continuous function defined on an open set $U \subset \mathbb{R}^2$. If $\partial_x f, \partial_y f$ are continuous on U , and there is a point $(x_0, y_0) \in (K \times K) \cap U$ such that

$$\left| \frac{\partial_y f|_{(x_0, y_0)}}{\partial_x f|_{(x_0, y_0)}} \right| \in (d_1, d_2) \quad \text{or} \quad \left| \frac{\partial_x f|_{(x_0, y_0)}}{\partial_y f|_{(x_0, y_0)}} \right| \in (d_1, d_2),$$

then $f_U(K, K)$ contains an interval.

Remark 1. We point that it is not easy to prove this result in terms of the Newhouse thickness theorem [36], as the IFS of K consists of very complicated overlaps.

Corollary 1. *Suppose $d_1 < d_2$, and let*

$$f(x, y) = x^\alpha \pm y^\beta \quad \text{with } \alpha\beta \neq 0 \text{ and } \frac{\alpha-1}{\beta-1} \notin \mathbb{Q}.$$

Then $f_U(K, K)$ contains an interval.

PROOF. Note that $\left| \frac{\partial_x f|_{(x_0, y_0)}}{\partial_y f|_{(x_0, y_0)}} \right| = \left| \frac{\alpha}{\beta} \right| \cdot \left| \frac{x^{\alpha-1}}{y^{\beta-1}} \right|$. Take $x_0 = \lambda^{k_1}$ and $y_0 = \lambda^{k_2}$, we have $\left| \frac{\partial_x f|_{(x_0, y_0)}}{\partial_y f|_{(x_0, y_0)}} \right| = \left| \frac{\alpha}{\beta} \right| \cdot \left| \lambda^{k_1(\frac{\alpha-1}{\beta-1}) - k_2} \right|^{(\beta-1)}$. Since $\frac{\alpha-1}{\beta-1} \notin \mathbb{Q}$, we can take integers k_1 and k_2 such that

$$k_1 \left(\frac{\alpha-1}{\beta-1} \right) - k_2 \text{ is so close to } \frac{1}{\beta-1} \log_\lambda \left| \frac{\beta}{\alpha} \cdot \frac{d_1 + d_2}{2} \right|,$$

hence $\left| \frac{\partial_x f|_{(x_0, y_0)}}{\partial_y f|_{(x_0, y_0)}} \right|$ is close enough to $\frac{d_1 + d_2}{2}$, which implies $\left| \frac{\partial_x f|_{(x_0, y_0)}}{\partial_y f|_{(x_0, y_0)}} \right| \in (d_1, d_2)$. Therefore, $f_U(K, K)$ contains an interval. $\square$

Note that if $c = \lambda = 1/3$, then K is the middle-third Cantor set. We have the following results.

Corollary 2. *Let C be the middle-third Cantor sets. If $\partial_x f, \partial_y f$ are continuous on U , and there is a point $(x_0, y_0) \in (C \times C) \cap U$ such that*

$$1 < \left| \frac{\partial_x f|_{(x_0, y_0)}}{\partial_y f|_{(x_0, y_0)}} \right| < 3 \text{ or } 1 < \left| \frac{\partial_y f|_{(x_0, y_0)}}{\partial_x f|_{(x_0, y_0)}} \right| < 3,$$

then $f_U(C, C)$ contains an interval.

Corollary 3. *Let C be the middle-third Cantor set. If $f(x, y)$ is one of the following functions,*

$$x^\alpha y^\beta (\alpha\beta \neq 0), \quad x^\alpha \pm y^\alpha (\alpha \neq 0), \quad x \pm y^2, \quad \sin(x) \cos(y), \quad x \sin(xy),$$

then $f_U(C, C)$ contains an interval.

This paper is organized as follows. In Section 2, we will give proofs of Theorem 1 and Corollary 3.

2. Proofs of the main results

In this section, we assume that $f : U \rightarrow \mathbb{R}$ is a function such that $\partial_x f, \partial_y f$ are continuous on U .

Fix $\lambda \in (0, 1)$ and $2\lambda \geq c \geq \lambda$ with $c + \lambda < 1$. Given $J = [x_1, x_2]$, we let $g_J(t) = x_1 + (x_2 - x_1)t$ with $g_J([0, 1]) = J$, and

$$\tilde{J} = J^{(1)} \cup J^{(2)} \cup J^{(3)},$$

where

$$J^{(i)} = g_J(S_i([0, 1])).$$

Suppose $G = \cup_{J \in \Lambda} J$ is the union of some closed intervals pairwise disjoint, we let $\tilde{G} = \cup_{J \in \Lambda} \tilde{J}$.

Lemma 1. *Suppose $I = [a, a + t]$ and $J = [b, b + t]$. Then*

$$\bigcap_{1 \leq i, j \leq 2} (I^{(i)} \times J^{(j)}), \quad \bigcap_{1 \leq i \leq 2} (I^{(i)} \times J^{(3)}), \quad \bigcap_{1 \leq j \leq 2} (I^{(3)} \times J^{(j)})$$

are non-empty.

PROOF. We note that

$$\begin{aligned} [a + (c - \lambda)t, a + \lambda t] \times [b + (c - \lambda)t, b + \lambda t] &\subset \bigcap_{1 \leq i, j \leq 2} (I^{(i)} \times J^{(j)}), \\ [a + (c - \lambda)t, a + \lambda t] \times [b + (1 - \lambda)t, b + t] &= \bigcap_{1 \leq i \leq 2} (I^{(i)} \times J^{(3)}), \\ [a + (1 - \lambda)t, a + t] \times [b + (c - \lambda)t, b + \lambda t] &= \bigcap_{1 \leq j \leq 2} (I^{(3)} \times J^{(j)}). \quad \square \end{aligned}$$

Let $H = [0, 1]$. For any $i_1 \cdots i_n \in \{1, 2, 3\}^n$, we call $f_{i_1 \cdots i_n}(H) = (f_{i_1} \circ \cdots \circ f_{i_n})(H)$ a basic interval of rank n , or n -basic interval, which has length λ^n . Denote by H_n the collection of all these basic intervals of rank n . We say that $I \times J$ is a basic square of $K \times K$, if I and J are basic intervals of the same rank. Suppose A and B are the left and right endpoints of some basic intervals in H_k for some $k \geq 1$, respectively. Denote by G_n the union of basic intervals of rank n contained in $[A, B]$.

The following Lemma 2 comes from [2] and [39], here we give its proof just for the self-containedness of the paper.

Lemma 2. *Let $F : U \rightarrow \mathbb{R}$ be a continuous function. Suppose A and B (M and N) are the left and right endpoints of some basic intervals in H_{k_0} for some $k_0 \geq 1$, respectively, such that $[A, B] \times [M, N] \subset U$. Then $K \cap [A, B] = \bigcap_{n=k_0}^{\infty} G_n$, and $K \cap [M, N] = \bigcap_{n=k_0}^{\infty} G'_n$. Moreover, if for any $n \geq k_0$ and any two n -basic intervals $I \subset G_n$, $J \subset G'_n$ such that*

$$F(I, J) = F(\tilde{I}, \tilde{J}),$$

then $F(K \cap [A, B], K \cap [M, N]) = F(G_{k_0}, G'_{k_0})$.

PROOF. By the construction of G_n (G'_n), i.e. $G_{n+1} \subset G_n$ ($G'_{n+1} \subset G'_n$) for any $n \geq k_0$, it follows that

$$K \cap [A, B] = \bigcap_{n=k_0}^{\infty} G_n \quad \text{and} \quad K \cap [M, N] = \bigcap_{n=k_0}^{\infty} G'_n.$$

The continuity of F yields that

$$F(K \cap [A, B], K \cap [M, N]) = \bigcap_{n=k_0}^{\infty} F(G_n, G'_n).$$

In terms of the relation $G_{n+1} = \tilde{G}_n$, $G'_{n+1} = \tilde{G}'_n$ and the condition in the lemma, it follows that

$$\begin{aligned} F(G_n, G'_n) &= \bigcup_{1 \leq i \leq t_n} \bigcup_{1 \leq j \leq t'_n} F(I_{n,i}, J_{n,j}) = \bigcup_{1 \leq i \leq t_n} \bigcup_{1 \leq j \leq t'_n} F(\tilde{I}_{n,i}, \tilde{J}_{n,j}) \\ &= F\left(\bigcup_{1 \leq i \leq t_n} \tilde{I}_{n,i}, \bigcup_{1 \leq j \leq t'_n} \tilde{J}_{n,j}\right) = F(G_{n+1}, G'_{n+1}). \end{aligned}$$

Therefore, $F(K \cap [A, B], K \cap [M, N]) = F(G_{k_0}, G'_{k_0})$. $\square$

Proposition 1. *If there is a point $(x_0, y_0) \in (K \times K) \cap U$ such that*

$$\begin{cases} \partial_x f|_{(x_0, y_0)} > 0 > \partial_y f|_{(x_0, y_0)}, \\ d_1 < \left| \frac{\partial_y f|_{(x_0, y_0)}}{\partial_x f|_{(x_0, y_0)}} \right| < d_2, \end{cases}$$

then $f_U(K, K)$ contains an interval.

PROOF. Let $I \times J (\subset U)$ be a basic square of $K \times K$ close enough to (x_0, y_0) , that means $I = [a, a + t]$, $J = [b, b + t]$ are basic intervals such that (a, b) is close enough to (x_0, y_0) and t is small enough. Then

$$\tilde{I} = [a, a + ct] \cup [a + (1 - d)t, a + t], \quad \tilde{J} = [b, b + ct] \cup [b + (1 - d)t, b + t],$$

where $d = \lambda$. In what follows, we let $\delta = 1 - c - \lambda$. In terms of Lemma 2, it suffices to prove

$$f(I, J) = f(\tilde{I}, \tilde{J})$$

if we want to prove that $f_U(K, K)$ contains an interval. Since

$$\partial_x f|_{(x_0, y_0)} > 0 > \partial_y f|_{(x_0, y_0)},$$

it follows that

$$f(I, J) = [f(a, b + t), f(a + t, b)].$$

Moreover, in terms of the conditions

$$\frac{\delta}{d} < \left| \frac{\partial_y f|_{(x_0, y_0)}}{\partial_x f|_{(x_0, y_0)}} \right| < \frac{1}{\delta} \quad \text{and} \quad c \geq d = \lambda,$$

we conclude that

$$h_1(x, y) = f(x - \delta t, y) - f(x, y + dt) = t(-d\partial_y f - \delta\partial_x f) + o(t) \geq 0,$$

$$h_2(x, y) = f(x, y + \delta t) - f(x - t, y) = t(\partial_x f + \delta\partial_y f) + o(t) \geq 0,$$

$$h_3(x, y) = f(x - \delta t, y) - f(x, y + ct) = t(-c\partial_y f - \delta\partial_x f) + o(t) \geq 0,$$

where $o(t)/t \rightarrow 0$ uniformly as $t \rightarrow 0$, i.e. $o(t)$ is independent of the choice of (x, y) as $\partial_x f$ and $\partial_y f$ are continuous at (x_0, y_0) . In fact,

$$f(\tilde{I}, \tilde{J}) = \bigcup_{1 \leq i, j \leq 3} f(I^{(i)}, J^{(j)}).$$

Using Lemma 1, we obtain that

$$f(\tilde{I}, \tilde{J}) = \bigcup_{1 \leq i, j \leq 3} f(I^{(i)}, J^{(j)}) = J_1 \cup J_2 \cup J_3 \cup J_4,$$

where

$$\begin{aligned} J_1 &= [f(A), f(B)], & J_2 &= [f(C), f(D)], \\ J_3 &= [f(E), f(F)], & J_4 &= [f(G), f(H)]. \end{aligned}$$




Figure 1. Basic square in Proposition 1

Hence, in order to prove $f(I, J) = f(\tilde{I}, \tilde{J})$, we only need to check

$$\begin{cases} f(B) - f(C) \geq 0, \\ f(D) - f(E) \geq 0, \\ f(F) - f(G) \geq 0. \end{cases}$$

However, the above inequalities are the direct consequences of

$$\begin{aligned} f(B) - f(C) &= h_1(P) \geq 0, & f(D) - f(E) &= h_2(Q) \geq 0, \\ f(F) - f(G) &= h_3(R) \geq 0. \end{aligned}$$

□

For the function $f(x, y)$ in Proposition 1, when considering the function $-f(x, y)$, we obtain the following Proposition 2.

Proposition 2. *If there is a point $(x_0, y_0) \in (K \times K) \cap U$ such that*

$$\begin{cases} \partial_x f|_{(x_0, y_0)} < 0 < \partial_y f|_{(x_0, y_0)}, \\ d_1 < \left| \frac{\partial_y f|_{(x_0, y_0)}}{\partial_x f|_{(x_0, y_0)}} \right| < d_2, \end{cases}$$

then $f_U(K, K)$ contains an interval.

By the symmetry of $K \times K$, we obtain the following result.

Proposition 3. *If there is a point $(x_0, y_0) \in (K \times K) \cap U$ such that*

$$\begin{cases} \partial_x f|_{(x_0, y_0)} \cdot \partial_y f|_{(x_0, y_0)} < 0, \\ d_1 < \left| \frac{\partial_x f|_{(x_0, y_0)}}{\partial_y f|_{(x_0, y_0)}} \right| < d_2, \end{cases}$$

then $f_U(K, K)$ contains an interval.

With a similar discussion, we are allowed to prove the following result.

Proposition 4. *If there is a point $(x_0, y_0) \in (K \times K) \cap U$ such that*

$$\begin{cases} \partial_x f|_{(x_0, y_0)} > 0, \partial_y f|_{(x_0, y_0)} > 0, \\ d_1 < \left| \frac{\partial_x f|_{(x_0, y_0)}}{\partial_y f|_{(x_0, y_0)}} \right| < d_2, \end{cases}$$

then $f_U(K, K)$ contains an interval.

PROOF. The proof is similar to Proposition 1. Let $d = \lambda$ and $\delta = 1 - c - \lambda$. Let $I \times J (\subset U)$ be a basic square of $K \times K$ close enough to (x_0, y_0) , that means $I = [a, a + t]$, $J = [b, b + t]$ are basic intervals such that (a, b) is close enough to (x_0, y_0) and t is small enough. Then

$$\tilde{I} = [a, a + ct] \cup [a + (1 - d)t, a + t], \quad \tilde{J} = [b, b + ct] \cup [b + (1 - d)t, b + t].$$

In terms of Lemma 2, it suffices to prove

$$f(I, J) = f(\tilde{I}, \tilde{J}).$$

By virtue of

$$\partial_x f|_{(x_0, y_0)} > 0, \quad \partial_y f|_{(x_0, y_0)} > 0,$$

it follows that

$$f(I, J) = [f(a, b), f(a + t, b + t)] = [f(A), f(H)]$$

as in Figure 2.

Moreover, in terms of the conditions

$$\frac{\delta}{d} < \left| \frac{\partial_x f|_{(x_0, y_0)}}{\partial_y f|_{(x_0, y_0)}} \right| < \frac{1}{\delta} \quad \text{and} \quad c \geq d = \lambda,$$




Figure 2. Basic square in Proposition 4

we conclude that

$$\begin{aligned} h_4(x, y) &= f(x + ct, y) - f(x, y + \delta t) = t(c\partial_x f - \delta\partial_y f) + o(t) \geq 0, \\ h_5(x, y) &= f(x, y + t) - f(x + \delta t, y) = t(\partial_y f - \delta\partial_x f) + o(t) \geq 0, \\ h_6(x, y) &= f(x + dt, y) - f(x, y + \delta t) = t(d\partial_x f - \delta\partial_y f) + o(t) \geq 0, \end{aligned}$$

where $o(t)/t \rightarrow 0$ uniformly as $t \rightarrow 0$, i.e. $o(t)$ is independent of the choice of (x, y) , as $\partial_x f$ and $\partial_y f$ are continuous at (x_0, y_0) . Using Lemma 1, we obtain that

$$f(\tilde{I}, \tilde{J}) = \bigcup_{1 \leq i, j \leq 3} f(I^{(i)}, J^{(j)}) = J_5 \cup J_6 \cup J_7 \cup J_8,$$

where

$$J_5 = [f(A), f(B)], \quad J_6 = [f(C), f(D)],$$

$$J_7 = [f(E), f(F)], \quad J_8 = [f(G), f(H)].$$

Hence, in order to prove $f(I, J) = f(\tilde{I}, \tilde{J})$, we only need to prove

$$\begin{cases} f(B) - f(C) \geq 0, \\ f(D) - f(E) \geq 0, \\ f(F) - f(G) \geq 0. \end{cases}$$

However, the above inequalities follow immediately from

$$\begin{aligned} f(B) - f(C) &= h_4(P) \geq 0, & f(D) - f(E) &= h_5(Q) \geq 0, \\ f(F) - f(G) &= h_6(R) \geq 0. \end{aligned}$$

☐

In the same way, we obtain the following result.

Proposition 5. *If there is a point $(x_0, y_0) \in (K \times K) \cap U$ such that*

$$\begin{cases} \partial_x f|_{(x_0, y_0)} \cdot \partial_y f|_{(x_0, y_0)} > 0, \\ \left| \frac{\partial_y f|_{(x_0, y_0)}}{\partial_x f|_{(x_0, y_0)}} \right| \in (d_1, d_2) \quad \text{or} \quad \left| \frac{\partial_x f|_{(x_0, y_0)}}{\partial_y f|_{(x_0, y_0)}} \right| \in (d_1, d_2), \end{cases}$$

then $f_U(K, K)$ contains an interval.

PROOF OF THEOREM 1.

It follows from Propositions 1–5. □

PROOF OF COROLLARY 3.

It suffices to check the conditions in Theorem 1.

(1) If $f(x, y) = x^\alpha y^\beta$ with $\alpha\beta \neq 0$, using $(x^\alpha y^\beta) = (xy^{\beta/\alpha})^\alpha$, we only need to deal with $f(x, y) = xy^\gamma$. Using the symmetry, we may assume that $|\gamma| \geq 1$. Now we have

$$\partial_x f = y^\gamma \quad \text{and} \quad \partial_y f = \gamma xy^{\gamma-1}$$

with $\left| \frac{\partial_x f}{\partial_y f} \right| = \frac{1}{|\gamma|} \left| \frac{y}{x} \right|$. If $|\gamma| = 3^k$ for some integer $k \geq 0$, we take $y = 1$ and $x = (2/3) \cdot 3^{-k}$, hence $\left| \frac{\partial_x f}{\partial_y f} \right| = 3/2 \in (1, 3)$ in this case. Otherwise, if $3^k < |\gamma| < 3^{k+1}$ for some integer $k \geq 0$, then we take $y = 1$ and $x = 3^{-(k+1)}$, then $\left| \frac{\partial_x f}{\partial_y f} \right| \in (1, 3)$. Now, $f_U(C, C)$ contains an interval for $f(x, y) = x^\alpha y^\beta$ with $\alpha\beta \neq 0$.

(2) If $f(x, y) = x^\alpha \pm y^\alpha$ with $\alpha \neq 0$, then

$$|\partial_x f| = |\alpha| x^{\alpha-1} \quad \text{and} \quad |\partial_y f| = |\alpha| y^{\alpha-1}$$

with $\left| \frac{\partial_x f}{\partial_y f} \right| = \left| \frac{x^{\alpha-1}}{y^{\alpha-1}} \right|$. When $\alpha \neq 1$, take $x, y \in C$ such that y/x is close enough to 1, then $1 < \left| \frac{\partial_x f}{\partial_y f} \right| < 3$ or $1 < \left| \frac{\partial_y f}{\partial_x f} \right| < 3$. When $\alpha = 1$, the classical result $C + C = [0, 2]$ implies there is an interval in $f(C, C) = C + C$.

(3) If $f(x, y) = x \pm y^2$, then $\partial_x f = 1, |\partial_y f| = 2y$. Take $x_0 = 8/9, y_0 = 1/3$, which implies $1 < |1/(2y_0)| < 3$.

(4) If $f(x, y) = \sin(x) \cos(y)$, then

$$|\partial_x f| = |\cos x \cos y|, \quad |\partial_y f| = |\sin x \sin y|.$$

We take $(x_0, y_0) = (2/3, 2/3)$, and obtain that

$$|\cos(2/3) \cos(2/3)| = 0.6176 \dots, \quad |\sin(2/3) \sin(2/3)| = 0.3823 \dots,$$

and thus $1 < \left| \frac{\cos(2/3) \cos(2/3)}{\sin(2/3) \sin(2/3)} \right| = 1.615 \dots < 3$.

(5) If $f(x, y) = x \sin(xy)$, then

$$|\partial_x f| = |\sin(xy) + xy \cos xy|, \quad |\partial_y f| = |x^2 \cos(xy)|.$$

We take $(x_0, y_0) = (2/3, 2/3)$, and obtain that

$$\left| \frac{\partial_x f|_{(x_0, y_0)}}{\partial_y f|_{(x_0, y_0)}} \right| = \left| 1 + \frac{9 \sin(4/9)}{4 \cos(4/9)} \right| = 2.071 \dots \in (1, 3). \quad \square$$

Remark 2. Our idea can be implemented for other overlapping self-similar sets. Moreover, in Corollary 3, for some functions, we can obtain that $f_U(C, C)$ contains infinitely many closed intervals.

References

- [1] S. AKIYAMA and K. SCHEICHER, From number systems to shift radix systems, *Nihonkai Math. J.* **16** (2005), 95–106.
- [2] J. S. ATHREYA, B. REZNICK and J. T. TYSON, Cantor set arithmetic, *Amer. Math. Monthly* **126** (2019), 4–17.
- [3] S. BAKER, Generalized golden ratios over integer alphabets, *Integers* **14** (2014), Paper No. A15, 28 pp.
- [4] C. BANDT and S. GRAF, Self-similar sets. VII. A characterization of self-similar fractals with positive Hausdorff measure, *Proc. Amer. Math. Soc.* **114** (1992), 995–1001.
- [5] J. BARRIONUEVO, R. M. BURTON, K. DAJANI and C. KRAAIKAMP, Ergodic properties of generalized Lüroth series, *Acta Arith.* **74** (1996), 311–327.
- [6] K. DAJANI and M. DE VRIES, Measures of maximal entropy for random β -expansions, *J. Eur. Math. Soc. (JEMS)* **7** (2005), 51–68.
- [7] K. DAJANI and M. DE VRIES, Invariant densities for random β -expansions, *J. Eur. Math. Soc. (JEMS)* **9** (2007), 157–176.
- [8] K. DAJANI, K. JIANG and T. KEMPTON, Self-affine sets with positive Lebesgue measure, *Indag. Math. (N.S.)* **25** (2014), 774–784.
- [9] K. DAJANI, K. JIANG, D. KONG and W. LI, Multiple expansions of real numbers with digits set $\{0, 1, q\}$, *Math. Z.* **291** (2019), 1605–1619.
- [10] K. DAJANI, K. JIANG, D. KONG and W. LI, Multiple codings for self-similar sets with overlaps, 2016, arXiv:1603.09304.
- [11] K. DAJANI and C. KRAAIKAMP, Ergodic Theory of Numbers, *Mathematical Association of America, Washington, DC*, 2002.
- [12] K. DAJANI and M. OOMEN, Random N -continued fraction expansions, *J. Approx. Theory* **227** (2018), 1–26.

- [13] M. DE VRIES and V. KOMORNIK, Unique expansions of real numbers, *Adv. Math.* **221** (2009), 390–427.
- [14] P. ERDŐS, M. HORVÁTH and I. JOÓ, On the uniqueness of the expansions $1 = \sum q^{-n_i}$, *Acta Math. Hungar.* **58** (1991), 333–342.
- [15] P. GLENDINNING and N. SIDOROV, Unique representations of real numbers in non-integer bases, *Math. Res. Lett.* **8** (2001), 535–543.
- [16] Q. GUO, H. LI, Q. WANG and L. XI, Lipschitz equivalence of a class of self-similar sets with complete overlaps, *Ann. Acad. Sci. Fenn. Math.* **37** (2012), 229–243.
- [17] K. G. HARE and N. SIDOROV, Two-dimensional self-affine sets with interior points, and the set of uniqueness, *Nonlinearity* **29** (2016), 1–26.
- [18] K. G. HARE and N. SIDOROV, On a family of self-affine sets: topology, uniqueness, simultaneous expansions, *Ergodic Theory Dynam. Systems* **37** (2017), 193–227.
- [19] M. HOCHMAN and P. SHMERKIN, Local entropy averages and projections of fractal measures, *Ann. of Math. (2)* **175** (2012), 1001–1059.
- [20] M. HOCHMAN, On self-similar sets with overlaps and inverse theorems for entropy, *Ann. of Math. (2)* **180** (2014), 773–822.
- [21] J. E. HUTCHINSON, Fractals and self-similarity, *Indiana Univ. Math. J.* **30** (1981), 713–747.
- [22] K. JIANG, Hausdorff dimension of the arithmetic sum of self-similar sets, *Indag. Math. (N.S.)* **27** (2016), 684–701.
- [23] K. JIANG, Projections of cartesian products of the self-similar sets without the irrationality assumption, 2018, arXiv:1806.01080.
- [24] K. JIANG and L. XI, Arithmetic representations of real numbers in terms of self-similar sets, 2018, arXiv:1808.09724.
- [25] R. KENYON, Projecting the one-dimensional Sierpinski gasket, *Israel J. Math.* **97** (1997), 221–238.
- [26] V. KOMORNIK, Expansions in noninteger bases, *Integers* **11B** (2011), Paper No. A9, 30 pp.
- [27] J. C. LAGARIAS and Y. WANG, Self-affine tiles in $\mathbf{R}^n$, *Adv. Math.* **121** (1996), 21–49.
- [28] C. G. T. DE A. MOREIRA and J.-C. YOCOZ, Stable intersections of regular Cantor sets with large Hausdorff dimensions, *Ann. of Math. (2)* **154** (2001), 45–96.
- [29] S.-M. NGAI and Y. WANG, Hausdorff dimension of self-similar sets with overlaps, *J. London Math. Soc. (2)* **63** (2001), 655–672.
- [30] J. PALIS and F. TAKENS, Hyperbolicity and Sensitive Chaotic Dynamics at Homoclinic Bifurcations, Fractal Dimensions and Infinitely Many Attractors, Cambridge Studies in Advanced Mathematics, Vol. **35**, Cambridge University Press, Cambridge, 1993.
- [31] Y. PERES and P. SHMERKIN, Resonance between Cantor sets, *Ergodic Theory Dynam. Systems* **29** (2009), 201–221.
- [32] H. RAO and Z.-Y. WEN, A class of self-similar fractals with overlap structure, *Adv. in Appl. Math.* **20** (1998), 50–72.
- [33] A. SCHIEF, Separation properties for self-similar sets, *Proc. Amer. Math. Soc.* **122** (1994), 111–115.
- [34] P. SHMERKIN, Personal communication, 2018.
- [35] N. SIDOROV, Almost every number has a continuum of β -expansions, *Amer. Math. Monthly* **110** (2003), 838–842.
- [36] N. SIDOROV, Expansions in non-integer bases: lower, middle and top orders, *J. Number Theory* **129** (2009), 741–754.

- [37] H. STEINHAUS, Nowa Własność Mnogości Cantora, *Wector* (1927), 1–3; English translation: A new property of the Cantor set, In: H. Steinhaus, Selected Papers, *PWN, Warszawa*, (1985), 205–207.
- [38] Y. TAKAHASHI, Products of two Cantor sets, *Nonlinearity* **30** (2017), 2114–2137.
- [39] L. TIAN, J.GU, Q. YE, L.-F. XI and K. JIANG, Multiplication on self-similar sets with overlaps, 2018, arXiv:1807.05368.

LIFENG XI
DEPARTMENT OF MATHEMATICS
NINGBO UNIVERSITY
NINGBO 315211
P. R. CHINA

E-mail: xilifeng@nbu.edu.cn
xilifengningbo@yahoo.com

KAN JIANG
DEPARTMENT OF MATHEMATICS
NINGBO UNIVERSITY
NINGBO 315211
P. R. CHINA

E-mail: jiangkan@nbu.edu.cn
kanjiangbunnik@yahoo.com

JIALI ZHU
DEPARTMENT OF MATHEMATICS
NINGBO UNIVERSITY
NINGBO 315211
P. R. CHINA

E-mail: 1811071029@nbu.edu.cn

QIYANG PEI
DEPARTMENT OF MATHEMATICS
NINGBO UNIVERSITY
NINGBO 315211
P. R. CHINA

E-mail: akevbgf@hotmail.com

(Received November 28, 2018; revised May 10, 2019)