

On the derived length of Lie solvable group algebras

By TIBOR JUHÁSZ (Debrecen)

Dedicated to Professor Adalbert Bovdi on his 70th birthday

Abstract. Let G be a nilpotent group with cyclic commutator subgroup of order p^n and let F be a field of characteristic p . It is shown here that the Lie derived length of the group algebra FG is at most $\lceil \log_2(p^n + 1) \rceil$. Furthermore, this bound is achieved if and only if one of the following conditions is satisfied: (i) p is odd; (ii) $p = 2$ and $n \leq 2$; (iii) $p = 2$, $n \geq 3$ and the nilpotency class of G is at most n .

1. Introduction

Let G be a group and F a field. The group algebra FG may be considered as a Lie algebra, with the usual bracket operation. Define the Lie derived series and the strong Lie derived series of the group algebra FG respectively as follows: let $\delta^{[0]}(FG) = \delta^{(0)}(FG) = FG$ and

$$\delta^{[n+1]}(FG) = [\delta^{[n]}(FG), \delta^{[n]}(FG)],$$

$$\delta^{(n+1)}(FG) = [\delta^{(n)}(FG), \delta^{(n)}(FG)]FG,$$

where $[X, Y]$ is the additive subgroup generated by all Lie commutators $[x, y] = xy - yx$ with $x \in X$ and $y \in Y$. We say that FG is Lie solvable if there exists $m \in \mathbb{N}$ such that $\delta^{[m]}(FG) = 0$ and the number

Mathematics Subject Classification: 16S34, 17B30.

Key words and phrases: group algebras, Lie derived length.

$\text{dl}_L(FG) = \min\{m \in \mathbb{N} \mid \delta^{[m]}(FG) = 0\}$ is called the Lie derived length of FG . Similarly, FG is said to be strongly Lie solvable of derived length $\text{dl}^L(FG) = m$ if $\delta^{(m)}(FG) = 0$ and $\delta^{(m-1)}(FG) \neq 0$. According to a result of PASSI, PASSMAN and SEHGAL [6] a group algebra FG is Lie solvable if and only if one of the following conditions holds: (i) G is abelian; (ii) $\text{char}(F) = p$ and the commutator subgroup G' of G is a finite p -group; (iii) $\text{char}(F) = 2$ and G has a subgroup H of index 2 whose commutator subgroup H' is a finite 2-group. It is easy to check that a group algebra FG is strongly Lie solvable if either G is abelian or $\text{char}(F) = p$ and G' is a finite p -group.

Let G be a group with commutator subgroup of order p^n and $\text{char}(F) = p$. SHALEV [8] showed that

$$\text{dl}_L(FG) \leq \lceil \log_2(2t(G')) \rceil,$$

where $t(G')$ denotes the nilpotent index of the augmentation ideal of FG' and $\lceil r \rceil$ the upper integral part of a real number r . Moreover, Lemma 2.2 in [8] states that if G is nilpotent of class 2 then $\text{dl}_L(FG) \leq \lceil \log_2(t(G') + 1) \rceil$. In particular, according to Proposition 2.3 in [8], if G is nilpotent of class 2 and G' is cyclic of order p^n , then

$$\text{dl}_L(FG) = \lceil \log_2(p^n + 1) \rceil.$$

In this paper our goal is to generalize the above results of SHALEV for the case when the nilpotency class of G is not necessary 2. We obtain the following.

Theorem 1. *Let G be a nilpotent group with cyclic commutator subgroup of order p^n and let F be a field of characteristic p . Then $\text{dl}_L(FG) \leq \lceil \log_2(p^n + 1) \rceil$ with equality if and only if one of the following conditions holds:*

- (i) p is odd;
- (ii) $p = 2$ and G' is of order less than 8;
- (iii) $p = 2$, $n \geq 3$ and G has nilpotency class at most n .

Moreover, if $\text{char}(F) = 2$ we can extend our result as follows.

Corollary 1. *Let G be a nilpotent group with commutator subgroup of order 2^n and let F be a field of characteristic 2. Then $\text{dl}_L(FG) = n + 1$ if and only if one of the following conditions holds:*

- (i) G' is the noncyclic group of order 4 and $\gamma_3(G) \neq 1$;
- (ii) G' is cyclic of order less than 8;
- (iii) G' is cyclic, $n \geq 3$ and G has nilpotency class at most n .

In this paper $\omega(FG)$ denotes the augmentation ideal of FG ; for a normal subgroup $H \subseteq G$ we understand by $\mathfrak{J}(H)$ the ideal $FG \cdot \omega(FH)$. For $x, y, x_1, x_2, \dots, x_n \in G$ let $x^y = y^{-1}xy$, $(x, y) = x^{-1}x^y$, and the commutator $(x_1, x_2, \dots, x_n)$ is defined inductively to be $((x_1, x_2, \dots, x_{n-1}), x_n)$. By $\zeta(G)$ we mean the center of the group G , by $\gamma_n(G)$ the n -th term of the lower central series of G with $\gamma_1(G) = G$. Furthermore, denote by C_n the cyclic group of order n . The n -th term of the upper Lie power series of FG is denoted by $(FG)^{(n)}$ which is the associative ideal generated by all Lie commutators $[x, y]$ with $x \in FG^{(n-1)}$ and $y \in FG$, where $FG^{(1)} = FG$.

We shall use freely the identities

$$[x, yz] = [x, y]z + y[x, z], \quad [xy, z] = x[y, z] + [x, z]y,$$

and for units a, b, c the commutator identities

$$(a, bc) = (a, c)(a, b)^c = (a, c)(a, b)(a, b, c);$$

$$(ab, c) = (a, c)^b(b, c) = (a, c)(a, c, b)(b, c),$$

and that $[a, b] = ba((a, b) - 1)$.

2. Preliminaries

We begin with a statement of independent interest about the strong Lie derived length of group algebras which generalizes the Corollary 4 of BAGIŃSKI's paper [1].

Proposition 1. *Let G be a nilpotent group whose commutator subgroup G' is a finite p -group and let $\text{char}(F) = p$. If $\gamma_3(G) \subseteq (G')^p$ then*

$$\text{dl}^L(FG) = \lceil \log_2(t(G') + 1) \rceil.$$

PROOF. We show by induction on n that

$$\delta^{(n)}(FG) \subseteq (FG)^{(2^n)} \quad \text{for all } n \geq 0.$$

Evidently, $\delta^{(0)}(FG) = (FG)^{(1)}$ and assume that $\delta^{(n)}(FG) \subseteq (FG)^{(2^n)}$ for some n . By elementary properties of upper Lie power series,

$$\begin{aligned} \delta^{(n+1)}(FG) &= [\delta^{(n)}(FG), \delta^{(n)}(FG)]FG \subseteq [(FG)^{(2^n)}, (FG)^{(2^n)}]FG \\ &\subseteq (FG)^{(2^{n+1})}FG = (FG)^{(2^{n+1})}. \end{aligned}$$

In view of $\gamma_3(G) \subseteq (G')^p$, Theorem 3.1(i) from [3] states that $(FG)^{(2^n)} = \mathfrak{J}(G')^{2^n-1}$. Furthermore, Lemma 2.2 in [7] asserts $\mathfrak{J}(G')^{2^n-1} \subseteq \delta^{(n)}(FG)$ for all $n \geq 1$ and we have $\delta^{(n)}(FG) = \mathfrak{J}(G')^{2^n-1}$. It is easy to see that $\delta^{(n)}(FG) = 0$ if and only if $2^n - 1 \geq t(G')$, therefore $n \geq \log_2(t(G') + 1)$, which implies the statement. $\square$

Remark 1. (i) Since $\delta^{[n]}(FG) \subseteq \delta^{(n)}(FG)$ for all n , Proposition 1 yields an upper bound on the Lie derived length. Furthermore, if G is nilpotent with cyclic commutator subgroup of order p^n , then the condition $\gamma_3(G) \subseteq (G')^p$ holds and thus

$$\text{dl}_L(FG) \leq \lceil \log_2(p^n + 1) \rceil.$$

But, as we will see, the equality does not always hold.

(ii) As the following examples show, Proposition 1 breaks down without the condition $\gamma_3(G) \subseteq (G')^p$.

- Let G be a group with $G' = C_2 \times C_2$ such that $\gamma_3(G) \neq 1$ and let $\text{char}(F) = 2$. Then $\gamma_3(G) \not\subseteq (G')^2$ and, by Theorem 3 and Theorem 6 from [5], $\text{dl}^L(FG) > 2$. So $\text{dl}^L(FG) \neq \lceil \log_2(t(G') + 1) \rceil$, because now $\lceil \log_2(t(G') + 1) \rceil = 2$.
- Let G be a group with $G' = C_3 \times C_3 \times C_3$ such that $\gamma_3(G) \neq 1$ and let $\text{char}(F) = 3$. Then $\gamma_3(G) \not\subseteq (G')^3$ and, by Theorem 2.3 from [7], $\text{dl}^L(FG) > 3$. It follows that $\text{dl}^L(FG) \neq \lceil \log_2(t(G') + 1) \rceil$, because $\lceil \log_2(t(G') + 1) \rceil = 3$.

The next lemma will be used in the proof of the theorem.

Lemma 1. *Let G be a nilpotent group with cyclic commutator subgroup of order p^n and let $\text{char}(F) = p$. Then for all $m, k \geq 1$*

- (i) $[\omega^m(FG'), \omega(FG)] \subseteq \mathfrak{J}(G')^{m+p-1}$;
- (ii) $[\mathfrak{J}(G')^m, \mathfrak{J}(G')^k] \subseteq \mathfrak{J}(G')^{m+k+1}$.

PROOF. (i) We use induction on m . For every $y \in G'$ and $g \in G$ we have

$$[y - 1, g - 1] = [y, g] = gy((y, g) - 1) \in \mathfrak{J}(\gamma_3(G)) \subseteq \mathfrak{J}(G')^p.$$

This shows that the statement (i) holds for $m = 1$, because the elements of the form $g - 1$ with $1 \neq g \in G$ constitute an F -basis of $\omega(FG)$.

Now, assume that $[\omega^m(FG'), \omega(FG)] \subseteq \mathfrak{J}(G')^{m+p-1}$ for some m . Then

$$\begin{aligned} & [\omega^{m+1}(FG'), \omega(FG)] \\ & \subseteq \omega^m(FG')[\omega(FG'), \omega(FG)] + [\omega^m(FG'), \omega(FG)]\omega(FG') \\ & \subseteq \omega^m(FG')\mathfrak{J}(G')^p + \mathfrak{J}(G')^{m+p-1}\omega(FG') \subseteq \mathfrak{J}(G')^{m+p}, \end{aligned}$$

and the proof of (i) is complete.

(ii) The statement (ii) is a consequence of (i), because

$$\mathfrak{J}(G') = \omega(FG)\omega(FG') + \omega(FG'). \quad \square$$

Let G be a group with commutator subgroup $G' = \langle x \mid x^{2^n} = 1 \rangle$, where $n \geq 3$. It is well known that the automorphism group $\text{aut}(G')$ of G' is a direct product of the cyclic group $\langle \alpha \rangle$ of order 2 and the cyclic group $\langle \beta \rangle$ of order 2^{n-2} where the action of these automorphisms on G' is given by $\alpha(x) = x^{-1}$, $\beta(x) = x^5$. For $g \in G$, let τ_g denote the restriction to G' of the inner automorphism $h \mapsto h^g$ of G . The map $G \rightarrow \text{aut}(G)$, $g \mapsto \tau_g$ is a homomorphism whose kernel coincides with the centralizer $C = C_G(G')$. Clearly, the map $\varphi : G/C \rightarrow \text{aut}(G')$ given by $\varphi(gC) = \tau_g$ is a monomorphism.

The subset

$$G_\beta = \{g \in G \mid \varphi(gC) \in \langle \beta \rangle\}$$

of G will play an important role in the sequel. It is easy to check that G_β is a subgroup of index not greater than 2 and $g \in G_\beta$ if and only if $x^g = x^{5^i}$ for some $i \in \mathbb{Z}$.

Lemma 2. *Let G be a group with cyclic commutator subgroup of order 2^n , where $n \geq 3$ and let $\text{char}(F) = 2$. Then*

- (i) $(y, g) \in (G')^4$ for all $y \in G'$ and $g \in G_\beta$;
- (ii) $[\omega^m(FG'), \omega(FG_\beta)] \subseteq \mathfrak{I}(G')^{m+3}$.

PROOF. Let $g \in G_\beta$ and $y \in G'$.

(i) Clearly, $(y, g) = y^{-1}y^g = y^{-1+5^i}$ for some $i \geq 0$ and $-1 + 5^i \equiv 0 \pmod{4}$. Therefore, $(y, g) \in (G')^4$.

(ii) Using (i) we have that

$$[y - 1, g - 1] = [y, g] = gy((y, g) - 1) \in \mathfrak{I}(G')^4,$$

from which (ii) follows for $m = 1$. One can now finish the proof by induction, as in Lemma 1(i). $\square$

Lemma 3. *Let G be a group with commutator subgroup $G' = \langle x \mid x^{2^n} = 1 \rangle$, where $n \geq 3$. Then the following are equivalent:*

- (i) $G_\beta = G$.
- (ii) G has nilpotency class at most n .

PROOF. First of all, note that G is a nilpotent group of class at most $n + 1$.

(i) $\Rightarrow$ (ii) By Lemma 2(i), $\gamma_3(G) \subseteq (G')^4$, so $|\gamma_2(G)/\gamma_3(G)| \geq 4$ and the class of G is at most n .

(ii) $\Rightarrow$ (i) Suppose that G has nilpotency class at most n , but $G_\beta \neq G$. We claim that $x^{2^{k-2}} \in \gamma_k(G)$ for all $k \geq 2$. Indeed, this is clear for $k = 2$ and assume its truth for some $k \geq 2$. If $g \in G \setminus G_\beta$ then $(x^{2^{k-2}}, g) \in \gamma_{k+1}(G)$ and $(x^{2^{k-2}}, g) = x^{2^{k-2}(-1+5^i)} = (x^{2^{k-1}})^j$ with some i and odd j . This means that $x^{2^{k-1}} \in \gamma_{k+1}(G)$, as desired. Therefore, $\gamma_{n+1}(G) \neq 1$, which is a contradiction. $\square$

Lemma 4. *Let G be a group with commutator subgroup $G' = \langle x \mid x^{2^n} = 1 \rangle$, where $n \geq 3$. If G has nilpotency class $n + 1$ then $(g, h) \in (G')^2$ for all $g, h \in G_\beta$.*

PROOF. If the lemma were not true we could choose the elements $g, h \in G_\beta$ so that $(g, h) = x$. By definition of G_β we may additionally

assume that $(g, x) = 1$. Lemma 3 states that $G \setminus G_\beta \neq \emptyset$; let y be in $G \setminus G_\beta$. Evidently, $(g, y) = x^i$ for some i . Using the equalities

$$g^h = gx, \quad g^{h^{-1}} = g(x^{-1})^{h^{-1}}, \quad g^y = gx^i, \quad g^{y^{-1}} = g(x^{-i})^{y^{-1}}$$

it is easy to check

$$\begin{aligned} g &= g^{(h,y)} = g^{h^{-1}y^{-1}hy} = (g(x^{-1})^{h^{-1}})^{y^{-1}hy} = g^{y^{-1}hy}x^{-1} \\ &= (g(x^{-i})^{y^{-1}})^{hy}x^{-1} = (gx(x^{-i})^{y^{-1}h})^yx^{-1} = gx^ix^y(x^{-i})^{y^{-1}hy}x^{-1} \\ &= g(x, y)(x^{-i}, h^y), \end{aligned}$$

which is a contradiction. Indeed, keeping in mind that $y \in G \setminus G_\beta$ and $h \in G_\beta$ we have $(x, y) = x^{-1-5^j} \in \langle x^2 \rangle \setminus \langle x^4 \rangle$ and $(x^{-i}, h^y) = x^{i(1-5^l)} \in \langle x^4 \rangle$, thus $(x, y)(x^{-i}, h^y) \neq 1$. $\square$

The author wishes to thank C. BAGIŃSKI for the elementary proof of the previous lemma.

Lemma 5. *Let G be a group with commutator subgroup $G' = \langle x \mid x^{2^n} = 1 \rangle$, where $n \geq 3$ and let $\text{char}(F) = 2$. If G has nilpotency class $n+1$ then $\text{dl}_L(FG) \leq n$.*

PROOF. Clearly, the set of the Lie commutators $[a, b]$ with $a, b \in G$ spans the F -space $\delta^{[1]}(FG)$. Since $[a, b] = g^h + g$ with $g = ba$ and $h = b$, while of course $g^h + g = [a, b]$ with $a = h^{-1}g$ and $b = h$ whenever $g, h \in G$, this spanning set for $\delta^{[1]}(FG)$ can also be described as the set of the elements $g^h + g$ with $g, h \in G$. It follows that the Lie commutators $[g_1^{h_1} + g_1, g_2^{h_2} + g_2]$, where $g_1, g_2, h_1, h_2 \in G$, span $\delta^{[2]}(FG)$. We shall compute these Lie commutators. It is easy to check that

$$\begin{aligned} [g_1^{h_1} + g_1, g_2^{h_2} + g_2] &= g_2g_1 \left(((g_1, g_2) + 1)((g_2, h_2) + 1)((g_1, h_1) + 1) \right. \\ &\quad \left. + (g_2, h_2)((g_2, h_2, g_1) + 1)((g_1, h_1) + 1) \right. \\ &\quad \left. + (g_1, g_2)(g_1, h_1)((g_1, h_1, g_2) + 1)((g_2, h_2) + 1) \right). \end{aligned} \quad (1)$$

Firstly, if neither g_1 nor g_2 are in G_β then

$$[g_1^{h_1} + g_1, g_2^{h_2} + g_2] = b\varrho_3 \quad (2)$$

for some $b \in G_\beta$ and $\varrho_3 \in \omega^3(FG')$. Indeed, it is clear from the definition of G_β that then $g_2g_1 \in G_\beta$. Furthermore, the second factor on the right-hand side of (1) always belongs to $\omega^3(FG')$, because $\gamma_3(G) \subseteq (G')^2$.

Secondly, if g_1 or g_2 , say g_1 , belongs to G_β , then we claim that

$$[g_1^{h_1} + g_1, g_2^{h_2} + g_2] = g\varrho_4 \quad (3)$$

for some $\varrho_4 \in \omega^4(FG')$ and $g \in G$.

For $g_1 \in G_\beta$, Lemma 2(i) asserts $(g_2, h_2, g_1) \in (G')^4$, therefore the right-hand side of (1) can be written as

$$\begin{aligned} [g_1^{h_1} + g_1, g_2^{h_2} + g_2] &= g_2g_1 \left(\left((g_1, g_2) + 1 \right) \left((g_1, h_1) + 1 \right) \right. \\ &\quad \left. + (g_1, g_2)(g_1, h_1) \left((g_1, h_1, g_2) + 1 \right) \right) \left((g_2, h_2) + 1 \right) + g_2g_1\varrho_4 \end{aligned} \quad (4)$$

for some $\varrho_4 \in \omega^4(FG')$. In order to prove (3) it will be sufficient to show that the element for some $\varrho_4 \in \omega^4(FG')$. In order to prove (3) it will be sufficient to show that the element

$$\begin{aligned} \vartheta &= ((g_1, g_2) + 1)((g_1, h_1) + 1) \\ &\quad + (g_1, g_2)(g_1, h_1)((g_1, h_1, g_2) + 1) \end{aligned}$$

from the right-hand side of (4) belongs to $\omega^3(FG')$.

This is clear if g_2 also belongs to G_β , because then by Lemma 4 and Lemma 2(i) both summands of ϑ are in $\omega^3(FG')$. Furthermore, if $g_2 \notin G_\beta$, then $x^{g_2} = x^{-5^l}$ for some l and we distinguish the following three cases:

Case 1: $(g_1, h_1) \in (G')^2$. Then $(g_1, h_1, g_2) = (g_1, h_1)^{-1-5^l} \in (G')^4$ and $\vartheta \in \omega^3(FG')$.

Case 2: $(g_1, g_2) \in (G')^2$. By the well-known Hall–Witt identity,

$$(g_1, h_1, g_2)^{h_1^{-1}}(h_1^{-1}, g_2^{-1}, g_1)^{g_2}(g_2, g_1^{-1}, h_1^{-1})^{g_1} = 1.$$

Lemma 2(i) ensures that the second factor on the left-hand side belongs to $(G')^4$ and this is true for the last factor too, because

$$\begin{aligned} (g_2, g_1^{-1}, h_1^{-1}) &= ((g_1, g_2)^{g_1^{-1}}, h_1^{-1}) \\ &= ((g_1, g_2)^{g_1^{-1}})^{-1}(g_1, g_2)^{g_1^{-1}h_1^{-1}} = (g_1, g_2)^{2i} \end{aligned}$$

for some i . This means that $(g_1, h_1, g_2) \in (G')^4$, which proves $\vartheta \in \omega^3(FG')$.

Case 3: $(g_1, h_1) \notin (G')^2$ and $(g_1, g_2) \notin (G')^2$. Then $\langle (g_1, h_1) \rangle = \langle (g_1, g_2) \rangle = G'$ and $(g_1, g_2) = (g_1, h_1)^k$ for some odd k . With the notation $y = (g_1, h_1)$ ϑ can be written as

$$\vartheta = (y^k + 1)(y + 1) + y^{k+1}(y^{-5^l-1} + 1) = y^{k-5^l} + 1 + y(y^{k-1} + 1).$$

Of course, if $k \equiv 1 \pmod{4}$ then $y^{-5^l-1} + 1$ and $y(y^{k-1} + 1)$ are in $\omega^4(FG')$, therefore $\vartheta \in \omega^4(FG')$. Otherwise, if $k \equiv 3 \pmod{4}$ then $y^{k-3} + 1 \in \omega^4(FG')$ which implies that

$$\begin{aligned} y(y^{k-1} + 1) &= y((y^{k-3} + 1)(y^2 + 1) + (y^{k-3} + 1) + (y^2 + 1)) \\ &\equiv y(y^2 + 1) \equiv y^2 + 1 \pmod{\omega^3(FG')}. \end{aligned}$$

Similarly, we can obtain that

$$\begin{aligned} y^{k-5^l} + 1 &= (y^{k-5^l-2} + 1)(y^2 + 1) + (y^{k-5^l-2} + 1) + (y^2 + 1) \\ &\equiv y^2 + 1 \pmod{\omega^3(FG')}. \end{aligned}$$

Hence

$$\vartheta = y^{k-5^l} + 1 + y(y^{k-1} + 1) \equiv 2(y^2 + 1) \equiv 0 \pmod{\omega^3(FG')},$$

which completes the checking of (3).

Let S be the additive subgroup generated by all elements of the form $g\varrho_4$ and $b\varrho_3$, where $g \in G$, $b \in G_\beta$ and $\varrho_3 \in \omega^3(FG')$, $\varrho_4 \in \omega^4(FG')$. We claim that $[S, S] \subseteq \mathfrak{I}(G')^8$. Indeed, the additive subgroup $[S, S]$ can be spanned by some Lie commutators of the forms $[g\varrho_4, h\varrho_3]$ and $[b_1\varrho_3, b_2\eta_3]$ with $g \in G$, $b_1, b_2 \in G_\beta$, $\varrho_3, \eta_3 \in \omega^3(FG')$, $\varrho_4 \in \omega^4(FG')$. Furthermore, by Lemma 1(i),

$$\begin{aligned} [g\varrho_4, h\varrho_3] &= g[\varrho_4, h\varrho_3] + [g, h\varrho_3]\varrho_4 \\ &= g[\varrho_4, h + 1]\varrho_3 + hg((g, h) + 1)\varrho_3\varrho_4 + h[g + 1, \varrho_3]\varrho_4 \in \mathfrak{I}(G')^8, \end{aligned}$$

and by Lemma 2(ii) and Lemma 4,

$$\begin{aligned} [b_1\varrho_3, b_2\eta_3] &= b_1[\varrho_3, b_2\eta_3] + [b_1, b_2\eta_3]\varrho_3 \\ &= b_1[\varrho_3, b_2 + 1]\eta_3 + b_2[b_1 + 1, \eta_3]\varrho_3 + b_1b_2((b_1, b_2) + 1)\eta_3\varrho_3 \end{aligned}$$

also belongs to $\mathfrak{I}(G')^8$. Therefore, $[S, S] \subseteq \mathfrak{I}(G')^8$.

From (2) and (3) we get $\delta^{[2]}(FG) \subseteq S$, so we have

$$\delta^{[3]}(FG) = [\delta^{[2]}(FG), \delta^{[2]}(FG)] \subseteq [S, S] \subseteq \mathfrak{I}(G')^8.$$

Now, we use induction on k to show that

$$\delta^{[k]}(FG) \subseteq \mathfrak{I}(G')^{2^k} \quad \text{for all } k \geq 3. \quad (5)$$

Indeed, assuming the validity of (5) for some $k \geq 3$ we have

$$\delta^{[k+1]}(FG) = [\delta^{[k]}(FG), \delta^{[k]}(FG)] \subseteq [\mathfrak{I}(G')^{2^k}, \mathfrak{I}(G')^{2^k}] \subseteq \mathfrak{I}(G')^{2^{k+1}}$$

and this proves the truth of (5) for every $k \geq 3$.

Keeping in mind that G' has order 2^n , (5) implies that $\delta^{[n]}(FG) = 0$. Hence $\text{dl}_L(FG) \leq n$ and the proof is complete. $\square$

Lemma 6. *Let G be a nilpotent group with commutator subgroup $G' = \langle x \mid x^{p^n} = 1 \rangle$, $\text{char}(F) = p$ and assume that one of the following conditions holds:*

- (i) $p = 2$, $n \geq 3$ and G has nilpotency class at most n ;
- (ii) p is odd.

Then $\text{dl}_L(FG) = \lceil \log_2(p^n + 1) \rceil$.

PROOF. Since G' is cyclic of order p^n , we can choose $a, b \in G$ such that $(a, b) = x$. First of all, we claim that

$$[b^l a^m, b^s a^t] \equiv (ms - lt)b^{l+s}a^{m+t}(x - 1) \pmod{\mathfrak{I}(G')^2} \quad (6)$$

for every $l, s, m, t \in \mathbb{Z}$. Indeed, an easy computation yields

$$\begin{aligned} [b^l a^m, b^s a^t] &= b^s a^t b^l a^m ((b^l a^m, b^s a^t) - 1) \\ &= b^{l+s} a^{m+t} (a^t, b^l)^{a^m} ((b^l a^m, b^s a^t) - 1) \\ &\equiv b^{l+s} a^{m+t} ((b^l a^m, b^s a^t) - 1) \pmod{\mathfrak{I}(G')^2}, \end{aligned} \quad (7)$$

and

$$\begin{aligned} (b^l a^m, b^s a^t) &\equiv (b^l, a^t)(a^m, b^s) \equiv (b, a)^{lt}(a, b)^{ms} \\ &\equiv x^{ms-lt} \pmod{(G')^p}, \end{aligned}$$

because $\gamma_3(G) \subseteq (G')^p$. Thus $(b^l a^m, b^s a^t) = x^{ms-lt+pi}$ for some i . In view of the identity $uv - 1 = (u - 1)(v - 1) + (u - 1) + (v - 1)$, we have

$$\begin{aligned} (b^l a^m, b^s a^t) - 1 &\equiv (ms - lt + pi)(x - 1) \\ &\equiv (ms - lt)(x - 1) \pmod{\mathfrak{I}(G')^2} \end{aligned}$$

and putting this into (7) we obtain (6).

Now, let $k \geq 1$, $l, m, s, t \in \mathbb{Z}$, $z_1, z_2 \in \mathfrak{I}(G')^{2^k}$ and set

$$f_k(l, m, s, t, z_1, z_2) = [b^l a^m (x - 1)^{2^k - 1} + z_1, b^s a^t (x - 1)^{2^k - 1} + z_2].$$

We shall show that

$$\begin{aligned} f_k(l, m, s, t, z_1, z_2) \\ \equiv (ms - lt)b^{l+s}a^{m+t}(x - 1)^{2^{k+1}-1} \pmod{\mathfrak{I}(G')^{2^{k+1}}}. \end{aligned} \quad (8)$$

Lemma 1(ii) ensures that the elements $[b^l a^m (x - 1)^{2^k - 1}, z_2]$, $[z_1, z_2]$ and $[z_1, b^s a^t (x - 1)^{2^k - 1}]$ belong to $\mathfrak{I}(G')^{2^{k+1}}$, thus

$$\begin{aligned} f_k(l, m, s, t, z_1, z_2) \\ \equiv [b^l a^m (x - 1)^{2^k - 1}, b^s a^t (x - 1)^{2^k - 1}] \pmod{\mathfrak{I}(G')^{2^{k+1}}}. \end{aligned}$$

In the case $p = 2$, Lemma 3 forces $b^l a^m, b^s a^t \in G_\beta$, so we may apply Lemma 2(ii) to obtain that

$$[b^l a^m, (x - 1)^{2^k - 1}], [(x - 1)^{2^k - 1}, b^s a^t] \in \mathfrak{I}(G')^{2^{k+1}}.$$

Furthermore, for $p > 2$ the above inclusion follows from Lemma 1(i). This implies that

$$f_k(l, m, s, t, z_1, z_2) \equiv [b^l a^m, b^s a^t](x - 1)^{2^{k+1}-2} \pmod{\mathfrak{I}(G')^{2^{k+1}}},$$

which, together with (6), proves (8).

Define the following three series inductively by:

$$u_0 = a, \quad v_0 = b, \quad w_0 = b^{-1}a^{-1},$$

and, for $k > 0$,

$$u_{k+1} = [u_k, v_k], \quad v_{k+1} = [u_k, w_k], \quad w_{k+1} = [w_k, v_k].$$

Obviously, the k -th elements of these series belong to $\delta^{[k]}(FG)$. By induction on k we show for odd k that

$$\begin{aligned} u_k &\equiv \pm ba(x-1)^{2^k-1} \pmod{\mathfrak{I}(G')^{2^k}}; \\ v_k &\equiv \pm b^{-1}(x-1)^{2^k-1} \pmod{\mathfrak{I}(G')^{2^k}}; \\ w_k &\equiv \pm a^{-1}(x-1)^{2^k-1} \pmod{\mathfrak{I}(G')^{2^k}}, \end{aligned} \tag{9}$$

and if k is even then

$$\begin{aligned} u_k &\equiv \pm a(x-1)^{2^k-1} \pmod{\mathfrak{I}(G')^{2^k}}; \\ v_k &\equiv \pm b(x-1)^{2^k-1} \pmod{\mathfrak{I}(G')^{2^k}}; \\ w_k &\equiv \pm b^{-1}a^{-1}(x-1)^{2^k-1} \pmod{\mathfrak{I}(G')^{2^k}}. \end{aligned} \tag{10}$$

Evidently, $u_1 = [a, b] = ba(x-1)$, and by (6) we have

$$v_1 = [a, b^{-1}a^{-1}] \equiv -b^{-1}(x-1) \pmod{\mathfrak{I}(G')^2},$$

and $w_1 = [b^{-1}a^{-1}, b] \equiv -a^{-1}(x-1) \pmod{\mathfrak{I}(G')^2}$. Therefore (9) holds for $k = 1$.

Now, assume that (9) is true for some odd k . According to (8) the congruences

$$\begin{aligned} u_{k+1} &= \pm f_k(1, 1, -1, 0, u_k', v_k') \\ &\equiv \pm (-1)a(x-1)^{2^{k+1}-1} \pmod{\mathfrak{I}(G')^{2^{k+1}}}; \\ v_{k+1} &= \pm f_k(1, 1, 0, -1, u_k', v_k') \\ &\equiv \pm b(x-1)^{2^{k+1}-1} \pmod{\mathfrak{I}(G')^{2^{k+1}}}; \\ w_{k+1} &= \pm f_k(0, -1, -1, 0, u_k', v_k') \\ &\equiv \pm b^{-1}a^{-1}(x-1)^{2^{k+1}-1} \pmod{\mathfrak{I}(G')^{2^{k+1}}} \end{aligned}$$

hold, where u_k', v_k', w_k' are suitable elements from $\mathfrak{I}(G')^{2^k}$. Similarly,

supposing the truth of (10) for some even k we see

$$\begin{aligned}
 u_{k+1} &= \pm f_k(0, 1, 1, 0, u_k', v_k') \\
 &\equiv \pm ba(x-1)^{2^{k+1}-1} \pmod{\mathfrak{I}(G')^{2^{k+1}}}; \\
 v_{k+1} &= \pm f_k(0, 1, -1, -1, u_k', v_k') \\
 &\equiv \pm(-1)b^{-1}(x-1)^{2^{k+1}-1} \pmod{\mathfrak{I}(G')^{2^{k+1}}}; \\
 w_{k+1} &= \pm f_k(-1, -1, 1, 0, u_k', v_k') \\
 &\equiv \pm(-1)a^{-1}(x-1)^{2^{k+1}-1} \pmod{\mathfrak{I}(G')^{2^{k+1}}}.
 \end{aligned}$$

So, (9) and (10) are valid for any $k > 0$.

Assume that $k < \lceil \log_2(p^n + 1) \rceil$. Then $2^k - 1 < p^n$ and the elements u_k, v_k, w_k are nonzero in $\delta^{[k]}(FG)$, thus $\text{dl}_L(FG) \geq \lceil \log_2(p^n + 1) \rceil$.

At the same time, Remark 1(i) says that $\text{dl}_L(FG) \leq \lceil \log_2(p^n + 1) \rceil$. $\square$

3. Proofs of Theorem 1 and Corollary 1

PROOF OF THEOREM 1. For $p = 2$ and $n < 3$ the statement is a consequence of Remark 1(i) and Theorem 3 in [5]. In the other cases Lemma 5 and Lemma 6 state the required result. The proof is complete. $\square$

PROOF OF COROLLARY 1. Clearly, if G' is cyclic the statement immediately follows from Theorem 1. Now, assume that G' is noncyclic and $\delta^{[n]}(FG) \neq 0$. We know from [2] that FG is Lie nilpotent, and as we have already seen, $\delta^{[n]}(FG) \subseteq (FG)^{(2^n)}$. Thus $(FG)^{(2^n)} \neq 0$ and Theorem 1 of [4] states that $G' = C_2 \times C_2$ and $\gamma_3(G) \neq 1$. Conversely, if $G' = C_2 \times C_2$ then $t(G') = 3$ and $\text{dl}_L(FG) \leq \lceil \log_2(2 \cdot 3) \rceil = 3$. Furthermore, when $\gamma_3(G) \neq 1$, Theorem 3 in [5] says that $\text{dl}_L(FG) \neq 2$. Therefore $\text{dl}_L(FG) = 3$ and the corollary is proved. $\square$

ACKNOWLEDGEMENTS. The author would like to thank C. BAGIŃSKI and L. G. KOVÁCS for their valuable comments and suggestions.

References

- [1] C. BAGIŃSKI, A note on the derived length of the unit group of a modular group algebra, *Comm. Algebra* **30** (2002), 4905–4913.
- [2] A. A. BOVDI and I. I. KHRIPTA, Generalized Lie nilpotent group rings, *Math. Sb. (N.S.)* **129**(171), no. 1 (1986), 154–158.
- [3] A. A. BOVDI and J. KURDICS, Lie properties of group algebra and the nilpotency class of the group of units, *J. Algebra* **212**, no. 1 (1999), 28–64.
- [4] V. BOVDI and E. SPINELLI, Modular group algebras with maximal Lie nilpotency indices, *Publ. Math. Debrecen* **65**, no. 1–2 (2004), 243–252.
- [5] F. LEVIN and G. ROSENBERGER, Lie metabelian group rings, *North-Holland Math. Stud.* **126** (1986), 153–161 (*to appear in* Group and semigroup rings (Johannesburg, 1985)).
- [6] I. B. S. PASSI, D. S. PASSMAN and S. K. SEHGAL, Lie solvable group rings, *Canad. J. Math.* **25** (1973), 748–757.
- [7] M. SAHAI, Lie solvable group algebras of derived length three, *Publ. Mat.* **39**, no. 2 (1995), 233–240.
- [8] A. SHALEV, The derived length of Lie soluble group rings I., *J. Pure Appl. Algebra* **78**, no. 3 (1992), 291–300.
- [9] A. SHALEV, The derived length of Lie soluble group rings II., *J. London Math. Soc.* (2) **49**, no. 1 (1994), 93–99.

TIBOR JUHÁSZ
 INSTITUTE OF MATHEMATICS
 UNIVERSITY OF DEBRECEN
 H-4010 DEBRECEN, P.O. BOX 12
 HUNGARY

E-mail: juhaszti@math.klte.hu

(Received March 9, 2005; revised June 17, 2005)