

**Year: 2006****Vol.: 68****Fasc.: 3-4****Title:** Criteria of supersolubility for products of supersoluble groups**Author(s):** Wenbin Guo, K. P. Shum and Alexander Skiba

Let H and T be subgroups of a group G . Then we call H conditionally permutable (or in brevity, c -permutable) with T in G if there exists an element $x \in G$ such that $HT^x = T^xH$. If H is c -permutable with T in $\langle H, T \rangle$, then we call H completely c -permutable with T in G . By using the above concepts, we will give some new criterions for the supersolubility of a finite group $G = AB$, where A and B are both supersoluble groups. In particular, we prove that a finite group G is supersoluble if and only if $G = AB$, where both A, B are nilpotent subgroups of the group G and B is completely c -permutable in G with every term in some chief series of A . We will also give some applications of our new criterions.

Address:

Wenbin Guo
Department of Mathematics
Xuzhou Normal University
Xuzhou, 221116
P.R. China
and
Department of Mathematics
University of Science and Technology of China
Hefei 230026
P. R. China
E-mail: wbguo@pub.xz.jsinfo.net

Address:

K. P. Shum
Faculty of Science
The Chinese University of Hong Kong
Shatin, Hong Kong (SAR)
P.R. China
E-mail: kpshum@math.cuhk.edu.hk

Address:

Alexander Skiba
Department of Mathematics
Gomel State University of F. Skorina
Gomel 246028
Belarus
E-mail: skiba@gsu.unibel.by