

## Estimating the defect in Jensen's Inequality

By A. M. FINK (Ames)

**Abstract.** We consider how much the difference of the two sides of Jensen's Inequality might be. It has a connection with Grüss Inequality.

Grüss' Inequality gives an estimate on the defect in the Chebyshev Inequality and has found some application elsewhere, see for example [1]. Here we consider the defect in Jensen's Inequality.

To be more precise, let all the integrals exist and  $\mu$  be a normalized measure, i.e.  $\int_a^b d\mu = 1$ . Then define

$$T(f, g) \equiv \int_a^b fg d\mu - \int_a^b f d\mu \int_a^b g d\mu. \quad (1)$$

Chebyshev showed that if  $f$  and  $g$  are both increasing (or both decreasing) then  $T(f, g) \geq 0$ . Now we know that  $T(f, g) \geq 0$  if

$$[f(x) - f(y)][g(x) - g(y)] \geq 0 \quad \text{for all pairs } x, y. \quad (2)$$

Grüss considered how positive  $T$  could be. We will cite some of the results below.

Similarly, we want to look at Jensen's Inequality,

$$\phi\left(\int_a^b f d\mu\right) \leq \int_a^b \phi(f) d\mu \quad (3)$$

which we know to hold when  $\phi$  is convex,  $f$  is in  $L_\infty$ , and  $\mu \geq 0$  and normalized. So

$$E(\phi, f, \mu) \equiv \int_a^b \phi(f) d\mu - \phi\left(\int_a^b f d\mu\right) \quad (4)$$

---

*Mathematics Subject Classification:* 26D15.

*Key words and phrases:* Jensen's Inequality, Grüss' Inequality.

is non-negative for such  $(\phi, f, \mu)$ . Then we ask, how positive can it be?

Jensen's inequality follows from the graph of a convex function lying above its tangent lines. Explicitly

$$\phi(t) \geq \phi(s) + (t - s)\phi'(s)$$

for any  $t$  and  $s$ . Here we must retain equality. So we begin with assuming that  $\phi''$  exists on  $[a, b]$  and write

$$\phi(t) = \phi(s) + (t - s)\phi'(s) + \int_s^t (t - u)\phi''(u)d\mu. \tag{5}$$

In (5) we replace  $t$  by  $f(t)$  and  $s$  by  $M(f) \equiv \int_a^b f d\mu$ , and integrate with respect to  $\mu$ . We arrive at

$$\int_a^b \phi(t)d\mu(t) = \phi(M(f)) + \int_a^b \int_{M(f)}^{f(t)} [f(t) - u]\phi''(u)dud\mu(t) \tag{6}$$

from which we obtain the representation

$$E(\phi, f, \mu) = \int_a^b \int_{M(f)}^{f(t)} [f(t) - u]\phi''(u)dud\mu(t). \tag{7}$$

The integrand of the outside integral is non-negative. For let  $A \equiv \{t \mid f(t) \geq M(f)\}$  and  $B \equiv [a, b] \setminus A$ . Then

$$\begin{aligned} E(\phi, f, \mu) &= \int_A \int_{M(f)}^{f(t)} (f(t) - u)\phi''(u)dud\mu(t) \\ &\quad + \int_B \int_{f(t)}^{M(f)} (u - f(t))\phi''(u)dud\mu(t). \end{aligned} \tag{8}$$

Let  $S(t) = \frac{t^2}{2}$  so that  $S''(t) \equiv 1$ .

**Theorem 1.** *Let  $\phi$  be convex with  $\phi''$  continuous,  $f \in L_\infty$ , and  $\mu \geq 0$  with  $\int_a^b d\mu = 1$ . Then*

$$E(\phi, f, \mu) \leq \|\phi''\|_\infty E(S, f, \mu). \tag{9}$$

*Equality hold for  $\phi = S$ .*

PROOF. The proof is immediate from (8). Since all quantities are non-negative we may majorize  $E$  by replacing  $\phi''$  with  $\|\phi''\|$ , factoring it out of the integrals, and we get  $E$  with  $\phi''$  replaced by 1, i.e.  $\phi = S$ . □

Now the quantity  $E(S, f, \mu)$  is interesting.

$$E(S, f, \mu) = \frac{1}{2} \left[ \int_a^b f^2 d\mu - \left( \int_a^b f d\mu \right)^2 \right].$$

The quantity in brackets is the defect in the Cauchy-Schwarz Inequality for  $f$  and 1. It somehow measures how much  $f$  and 1 are independent functions.

Moreover,  $E(S, f\mu) = \frac{1}{2}T(f, f)$  so there is a connection with the Chebyshev and Grüss Inequalities. We now cite the relevant things about these. GRÜSS [2] in his original paper does not notice that (with  $\mu \geq 0$  and normalized)

$$T(f, g) = \frac{1}{2} \int_a^b \int_a^b [f(x) - f(y)][g(x) - g(y)] d\mu(x) d\mu(y).$$

This was known much earlier, see the chapter on Chebyshev's Inequality in [3]. By the Cauchy-Schwarz we have  $T(f, g) \leq T(f, f)^{\frac{1}{2}} T(g, g)^{\frac{1}{2}}$ . However he arrives at this inequality in other ways and provides a couple of upper bounds. He shows that (still with  $\mu$  normalized)

$$T(f, f) \leq [\max f - M(f)][M(f) - \min f]. \tag{10}$$

Since  $\min f \leq M(f) \leq \max f$ , this last expression is at most  $\frac{1}{4}(\max f - \min f)^2$ . Furthermore, equality holds here if  $f(t) = \text{sgn}(t - \bar{\mu})$  where  $\int_a^{\bar{\mu}} d\mu = \int_{\bar{\mu}}^b d\mu = \frac{1}{2}$ .

**Corollary 1.** *Let  $\phi$  be convex  $f$  bounded and integrable, and  $\mu \geq 0$  and normalized, then*

$$\begin{aligned} E(\phi, f, \mu) &\leq \frac{1}{2} \|\phi''\|_{\infty} [\max f - M(f)][M(f) - \min f] \\ &\leq \frac{1}{8} \|\phi''\|_{\infty} [\max f - \min f]^2. \end{aligned}$$

These are all best possible constants.

For Grüss' proof and other results one may consult either [3, p. 296] or [4].

The above results are straight forward for measures for which  $\mu \geq 0$ . We know however, that there are other situations when  $E(\phi, f, \mu) \geq 0$ . Suppose that  $f$  is monotone and bounded,  $\phi$  convex, and  $\mu$  end positive, i.e.

$$L(t) \equiv \int_a^t d\mu \geq 0, \quad \text{and,} \quad R(t) \equiv \int_t^b d\mu \geq 0 \quad \text{for } a \leq t \leq b. \tag{11}$$

Then  $E(\phi, f, \mu) \geq 0$ . See e.g. [3, p. 13] or [5], but it is a result known to Steffenson in the discrete case earlier. The above arguments need to be modified since the measure is no longer non-negative and the argument of the theorem fails.

**Theorem 2.** *If  $\phi$  is convex with  $\phi''$  continuous,  $f'$  exists and is strictly one sign, and  $\mu$  is a normalized measure satisfying (11), then the estimate of Theorem 1 holds.*

PROOF. We take the case when  $f' > 0$ . There is a  $c \in (a, b)$  such that  $f(t) \leq M(f)$  on  $[a, c)$  and  $f(t) \geq M(f)$  on  $[c, b]$ . To see this, we must show that  $f(a) < M(f) < f(b)$ . Recalling that  $L(b) = R(a) = 1$  we have by interchange of integration

$$\int_a^b f d\mu = f(b) - \int_a^b f' L dt < f(b)$$

and

$$\int_a^b f d\mu = f(a) + \int_a^b f' R dt > f(a).$$

Now

$$\begin{aligned} E(\phi, f, \mu) &= \int_a^c \int_{f(t)}^{M(f)} [u - f(t)] \phi''(u) du d\mu(t) \\ &\quad + \int_c^b \int_{M(f)}^{f(t)} [f(t) - u] \phi''(u) du d\mu(t) \\ &= \int_{f(a)}^{M(f)} \int_a^{f^{-1}(u)} (u - f(t)) d\mu(t) \phi''(u) du \\ &\quad + \int_{M(f)}^{f(b)} \int_{f^{-1}(u)}^b (f(t) - u) d\mu(t) \phi''(u) du. \end{aligned} \tag{12}$$

The inner integrals are by interchange of order

$$\int_a^{f^{-1}(u)} (u - f(t)) d\mu(t) = \int_a^{f^{-1}(u)} f'(t) L(t) dt \geq 0$$

and

$$\int_{f^{-1}(u)}^b (f(t) - u) d\mu(t) = \int_{f^{-1}(u)}^b f'(t) R(t) dt \geq 0.$$

So we may again majorize  $\phi''$  by its norm as in the proof of Theorem 1. □

Again the estimate which involves  $S(t)$  is  $T(f, f)$ . Since  $(f, f)$  clearly satisfies (2), this is non-negative and we may look for a Grüss type estimate. All of the results in [2] or [3] require that  $\mu \geq 0$ . However, in [4] we looked at some results for end positive measures.

**Corollary 2.** *If  $(\phi, f, \mu)$  satisfy the hypothesis of Theorem 2, then*

$$E(\phi, f, \mu) \leq \frac{1}{2} F [f(b) - f(a)]^2 \|\phi''\|_\infty,$$

and

$$E(\phi, f, \mu) \leq N \|f'\|_\infty^2 \|\phi''\|_\infty,$$

where  $F = \max_{a \leq t \leq x \leq b} L(t)R(x)$  and  $N = \int_a^b RL_1 dx$ ,  $L_1(x) = \int_a^x L(t)dt$  Both estimates have the best possible constants.

PROOF. These are direct applications of Theorems 11 and 16 of [4].  $\square$

It is possible using the identities (7) and (12) to get estimates using Hölder's Inequality with  $\|\phi''\|_p$  but they are not so nice and it is difficult to get best possible estimates.

### References

- [1] S. S. DRAGOMIR and I. FEDALOV, A Grüss type inequality for mappings of bounded variation and applications to numerical analysis, *Funct. Anal & Appl.* **6**, no. 3 (2001), 425–438.
- [2] G. GRÜSS, Über das Maximum des absoluten Betrages von  $\frac{1}{b-a} \int_a^b f(x)g(x)dx - \frac{1}{(b-a)^2} \int_a^b f(x)dx \int_a^b g(x)dx$ , *Math. Z.* **39** (1935), 215–226.
- [3] D. S. MITRONOVIĆ, J. E. PEČARIĆ and A. M. FINK, Classical and New Inequalities in Analysis, *Kluwer Academic Publishers*, 1993.
- [4] A. M. FINK, A treatise on Grüss' Inequality, in: Analytic and Geometric Inequalities and Application, (T. M. Rassias and H. M. Srivastava, eds.), *Kluwer Academic Publishers*, 1999, 93–114.
- [5] A. M. FINK and M. JODEIT JR., Jensen inequalities for functions with higher monotonicities, *Aequationes Math.* **40** (1990), 26–43.

A. M. FINK  
 MATHEMATICS DEPARTMENT  
 IOWA STATE UNIVERSITY  
 AMES, IA 50011  
 USA

*E-mail:* fink@math.iastate.edu

(Received March 18, 2005; revised May 17, 2005)