

Conformal flatness of complex Finsler structures

By TADASHI AIKOU (Kagoshima)

Abstract. In the present paper, we shall be concerned with conformal flatness of convex Finsler structures. We introduce a complex Finsler connection and define its conformal curvature Θ . This curvature Θ is invariant by any conformal rescaling of the convex Finsler structure. Our main result is to show that this conformal curvature measures the conformal flatness.

1. Introduction

Let $\pi : E \rightarrow M$ be a holomorphic vector bundle of rank r over a complex manifold M of complex dimension n . The total space E is also a complex manifold of complex dimension $n + r$. The tangent vectors along the fibres define a holomorphic vector sub-bundle $\mathcal{V}$ of the holomorphic tangent bundle TE , that is, $\mathcal{V} = \ker d\pi$. Then we know that $\mathcal{V} \cong \pi^{-1}E$, and $\mathcal{V}$ is integrable. If a convex Finsler structure F is given on E , we can introduce a natural Hermitian structure h on $\mathcal{V}$. The Hermitian geometry of $(\mathcal{V}, h)$ has been investigated by KOBAYASHI [8], and a number of important results were obtained.

In this paper, however, we shall study the bundle $(\mathcal{V}, h)$ by using its Finsler connection, not the Hermitian connection. This connection is derived from the given Finsler structure F and a splitting of the following exact sequence of holomorphic vector bundles

$$(1.1) \quad 0 \rightarrow \mathcal{V} \xrightarrow{i} TE \rightarrow \pi^{-1}TM \rightarrow 0,$$

Mathematics Subject Classification: 53B40, 53C25.

Key words and phrases: complex Finsler structure, complex Finsler connection, conformal flatness, conformal curvature.

or equivalently its dual

$$(1.2) \quad 0 \rightarrow \pi^{-1}TM^* \rightarrow TE^* \rightarrow \mathcal{V}^* \rightarrow 0.$$

Let $\sigma : \pi^{-1}TM \rightarrow TE$ be a splitting of (1.1). Putting $\mathcal{H} := \sigma(\pi^{-1}TM)$, it defines a transversal distribution of $\mathcal{V}$ which is C^∞ isomorphic to $\pi^{-1}TM$. Let $\{s_1, \dots, s_r\}$ be a local holomorphic frame field of E on an open set U . Then it induces a local complex coordinate system $(z^1, \dots, z^n, \xi^1, \dots, \xi^r)$ on the open set $\pi^{-1}(U)$ in E . Now, $\mathcal{H}$ has local frame field X_α on $\pi^{-1}(U)$ of the form

$$(1.3) \quad \sigma\left(\frac{\partial}{\partial z^\alpha}\right) = X_\alpha = \frac{\partial}{\partial z^\alpha} - \sum_{l=1}^r N_\alpha^l \frac{\partial}{\partial \xi^l},$$

where $\{N_\alpha^l\}$ are local functions on $\pi^{-1}(U)$ satisfying some transformation law. Such a family $\{N_\alpha^l\}$ is called a *non-linear connection* on E .

If a splitting σ is given on the sequence (1.1), the co-tangent bundle TE^* has a C^∞ -splitting $TE^* = \mathcal{H}^* \oplus \mathcal{V}^*$. Then, according to this splitting, the differential operator ∂ is also decomposed as $\partial = \partial_{\mathcal{H}} + \partial_{\mathcal{V}}$, where $\partial_{\mathcal{H}}$ is the natural projection to the transversal part $\mathcal{H}^*$, and $\partial_{\mathcal{V}} = \partial - \partial_{\mathcal{H}}$. If a convex Finsler structure F is given on E , we can take a non-linear connection satisfying $\partial_{\mathcal{H}}^2 \equiv 0$, and by using this, we can introduce a canonical Finsler connection (cf. [1], [3], [4], [5]).

In a previous paper [3], we have discussed the conformal flatness of a Finsler structure in terms of Weyl connections. In this paper, we shall introduce a conformal invariant Θ which measures the conformal flatness of a complex Finsler structure, and we show that the vanishing of Θ is equivalent to the conformal flatness of F (Theorem 3.3). Our conformal invariant Θ is a natural generalization of the one in the Hermitian case (cf. [10]).

2. Finsler structures and Finsler connections

Let M be a connected complex manifold of dimension n , and E a holomorphic vector bundle of rank r over M . Each fibre E_z is a complex vector space of dimension r . In the case of $r = 1$, any complex Finsler metric is a Hermitian metric. Hence, in the sequel we assume that $\text{rank } E \geq 2$.

Definition 2.1 ([6]). A function $F(z, \xi)$ on E is said to be a *complex Finsler structure* if it satisfies the following conditions:

$$(2.1) \quad F(z, \xi) \geq 0, \text{ and } F(z, \xi) = 0 \text{ iff } \xi = 0,$$

$$(2.2) \quad F(z, \xi) \text{ is } C^\infty \text{ on the outside of the zero-section and continuous on } E,$$

$$(2.3) \quad F(z, \lambda\xi) = |\lambda|^2 F(z, \xi) \text{ for an arbitrary } \lambda \in \mathbb{C}.$$

We shall fix an open covering $\{U\}$ with holomorphic frame field $\{s_U\}$ and the induced coordinate system $\{\pi^{-1}(U), (z, \xi)\}$ on E . A complex Finsler structure F is said to be *convex* if the Hermitian matrix $(F_{i\bar{j}})$ defined by

$$F_{i\bar{j}} = \frac{\partial^2 F}{\partial \xi^i \partial \bar{\xi}^j}$$

is positive-definite. In this paper we always suppose the convexity of F . On $\pi^{-1}(U)$, the vertical bundle $\mathcal{V}$ is spanned by the *vertical* vector fields $Y_1 := \partial/\partial \xi^1, \dots, Y_r := \partial/\partial \xi^r$. We define a Hermitian structure h on $\mathcal{V}$ by

$$h(Y_i, Y_j) = F_{i\bar{j}}.$$

The transversal distribution $\mathcal{H}$ is locally spanned by the *transversal* vector fields $\{X_\alpha\}$ of the form (1.3) for a complex non-linear connection N_α^i . We shall determine a canonical non-linear connection N_α^i .

The connection form θ of the Hermitian connection ∇^h of $(\mathcal{V}, h)$ is given by

$$\theta_j^i = \sum F^{i\bar{m}} \partial F_{j\bar{m}} = \sum F^{i\bar{m}} \left(\sum \frac{\partial F_{j\bar{m}}}{\partial z^\alpha} dz^\alpha + \sum \frac{\partial F_{j\bar{m}}}{\partial \xi^k} d\xi^k \right).$$

On the other hand, $\mathcal{V}$ has a canonical holomorphic section $\epsilon : (z, \xi) \rightarrow (z, \xi; \xi)$, or in local coordinates

$$\epsilon(z, \xi) = \sum_J \xi^j Y_j.$$

Then we put

$$\nabla^h \epsilon = \sum_i \theta^i \otimes Y_i$$

where the $(1, 0)$ -form θ^j is defined by

$$\theta^i = d\xi^i + \sum_m \theta_m^i \xi^m = d\xi^i + \sum_{m, \alpha} F^{i\bar{m}} \frac{\partial F_{l\bar{m}}}{\partial z^\alpha} \xi^l dz^\alpha.$$

Here we used the identity $\sum_{k=1}^r \xi^k (\partial F_{i\bar{j}} / \partial \xi^k) \equiv 0$ which is derived from the homogeneity assumption (2.3). Now we shall define a morphism $\sigma^* : \mathcal{V}^* \rightarrow TM^*$ by

$$\sigma^*(d\xi^i) = \theta^i.$$

It is trivial that σ^* defines a splitting of the sequence (1.2). For this splitting σ^* , in local coordinates, the non-linear connection N_α^i is given by

$$(2.5) \quad N_\alpha^i = \sum F^{i\bar{m}} \frac{\partial F_{l\bar{m}}}{\partial z^\alpha} \xi^l.$$

The differential operators $\partial_{\mathcal{H}}$ and $\partial_{\mathcal{V}}$ are given by

$$\begin{aligned} \partial_{\mathcal{H}} f &= \sum_{\alpha} X_{\alpha} f dz^{\alpha} = \sum_{\alpha} \left(\frac{\partial f}{\partial z^{\alpha}} - \sum_m N_{\alpha}^m \frac{\partial f}{\partial \xi^m} \right) dz^{\alpha}, \\ \partial_{\mathcal{V}} f &= \sum_m Y_m f \theta^m = \sum_m \frac{\partial f}{\partial \xi^m} \theta^m \end{aligned}$$

for an arbitrary function f on E . By using the facts that $\sum F_{i\bar{j}} \xi^i \bar{\xi}^j = F$ and $X_{\alpha} \xi^i = -N_{\alpha}^i$, we get the following identity

$$(2.6) \quad \partial_{\mathcal{H}} F = \sum_{\alpha} \left(\frac{\partial F}{\partial z^{\alpha}} - \sum_l N_{\alpha}^l \frac{\partial F}{\partial \xi^l} \right) dz^{\alpha} \equiv 0.$$

We denote by the index $\mathbb{C}$ the complexification of vector bundles, e.g., $T^{\mathbb{C}}M = TM \oplus \overline{TM}$, $\mathcal{V}^{\mathbb{C}} = \mathcal{V} \oplus \overline{\mathcal{V}}, \dots$

Definition 2.2 ([2]). A connection $\nabla : \Gamma(\mathcal{V}) \rightarrow \Gamma(\mathcal{V} \otimes T^{\mathbb{C}}E^*)$ defined by the following two properties is called the *Finsler connection* of (E, F) or $(\mathcal{V}, h)$.

- (1) ∇ is a $(1, 0)$ -type connection,
- (2) ∇ satisfies

$$d_{\mathcal{H}} h(Z, W) = h(\nabla Z, W) + h(Z, \nabla W)$$

for $\forall Z, W \in \Gamma(\mathcal{V})$, where we put $d_{\mathcal{H}} = \partial_{\mathcal{H}} + \bar{\partial}_{\mathcal{H}}$.

Since ∇ is of $(1, 0)$ -type, we may put

$$\nabla Y_j = \sum_m \omega_j^m Y_m$$

for $(1, 0)$ -forms ω_j^m . Then we have $d_{\mathcal{H}} F_{i\bar{j}} = \sum F_{m\bar{j}} \omega_i^m + F_{i\bar{m}} \overline{\omega_j^m}$. Hence, since ω_j^i is of $(1, 0)$ -type, the connection form ω of ∇ is given by the following transversal form:

$$\omega_j^i = \sum F^{i\bar{m}} \partial_{\mathcal{H}} F_{j\bar{m}} = \sum_{\alpha} \Gamma_{j\alpha}^i dz^{\alpha},$$

where the coefficients $\Gamma_{j\alpha}^i$ are given by

$$(2.7) \quad \Gamma_{j\alpha}^i = \sum F^{i\bar{m}} X_{\alpha} F_{j\bar{m}} = \sum F^{i\bar{m}} \left(\frac{\partial F_{j\bar{m}}}{\partial z^{\alpha}} - \sum N_{\alpha}^l \frac{\partial F_{j\bar{m}}}{\partial \xi^l} \right)$$

with the functions N_{α}^i defined by (2.5). Moreover, we get easily the following relation:

$$(2.8) \quad \Gamma_{j\alpha}^i = \frac{\partial N_{\alpha}^i}{\partial \xi^j}.$$

Remark 2.1. The following relation between the connection forms θ_j^i and ω_j^i is easily obtained:

$$\theta_j^i = \omega_j^i + \sum_{k,l} C_{jk}^i \theta^k,$$

where we put $C_{jk}^i = \sum F^{i\bar{m}} Y_k F_{j\bar{m}}$. Hence the Hermitian connection ∇^h corresponds to the so-called *Cartan connection* and our connection ∇ corresponds to the so-called *Rund connection* in real Finsler geometry (cf. [9]). We note that, from (2.8), our connection ∇ also corresponds to the *Berwald connection*.

We shall compute the curvature $\Omega = d\omega + \omega \wedge \omega$ and investigate its local expressions with respect to $\{dz^{\alpha}, \theta^i\}$. Since the non-linear connection N_{α}^i is given by (2.5), we have

Lemma 2.1. $\partial_{\mathcal{H}}\omega + \omega \wedge \omega \equiv 0$.

PROOF. The proof is obtained by direct calculation. If we put

$$R_{j\alpha\beta}^i = X_{\alpha}\Gamma_{j\beta}^i - X_{\beta}\Gamma_{j\alpha}^i + \sum \Gamma_{m\alpha}^i\Gamma_{j\beta}^m - \sum \Gamma_{m\beta}^i\Gamma_{j\alpha}^m,$$

the right hand side can be written as

$$\partial_{\mathcal{H}}\omega_j^i + \sum \omega_m^i \wedge \omega_j^m = -\frac{1}{2} \sum R_{j\alpha\beta}^i dz^{\alpha} \wedge dz^{\beta}.$$

Hence we must prove $R_{j\alpha\beta}^i \equiv 0$. Since $\Gamma_{j\alpha}^i = \sum F^{i\bar{m}} X_{\alpha} F_{j\bar{m}}$,

$$\begin{aligned} R_{j\alpha\beta}^i &= \sum \left\{ X_{\alpha} F^{i\bar{m}} X_{\beta} F_{j\bar{m}} + F^{i\bar{m}} X_{\alpha} X_{\beta} F_{j\bar{m}} - X_{\beta} F^{i\bar{m}} X_{\alpha} F_{j\bar{m}} \right. \\ &\quad \left. - F^{i\bar{m}} X_{\alpha} X_{\alpha} F_{j\bar{m}} + \Gamma_{m\alpha}^i \Gamma_{j\beta}^m - \Gamma_{m\beta}^i \Gamma_{j\alpha}^m \right\} \\ &= \sum \left\{ X_{\alpha} F^{i\bar{m}} X_{\beta} F_{j\bar{m}} - X_{\beta} F^{i\bar{m}} X_{\alpha} F_{j\bar{m}} + F^{i\bar{m}} \sum R_{\alpha\beta}^l Y_l F_{j\bar{m}} \right. \\ &\quad \left. + \Gamma_{m\alpha}^i \Gamma_{j\beta}^m - \Gamma_{m\beta}^i \Gamma_{j\alpha}^m \right\} \\ &= \sum C_{jm}^i R_{\alpha\beta}^m, \end{aligned}$$

where we used $\Gamma_{j\alpha}^i = \sum F^{i\bar{m}} X_{\alpha} F_{j\bar{m}}$, and put $R_{\alpha\beta}^i = X_{\alpha} N_{\beta}^i - X_{\beta} N_{\alpha}^i$.

On the other hand, by definition of $R_{j\alpha\beta}^i$ and $\sum \xi^j \Gamma_{j\alpha}^i = N_{\alpha}^i$, we get easily $R_{\alpha\beta}^i = \sum R_{j\alpha\beta}^i \xi^j$. The equation above and $\sum C_{jk}^i \xi^j \equiv 0$ imply $R_{\alpha\beta}^i = 0$, and so $R_{j\alpha\beta}^i \equiv 0$. $\square$

By virtue of $\Omega = d\omega + \omega \wedge \omega = \bar{\partial}\omega + \partial_{\mathcal{V}}\omega + (\partial_{\mathcal{H}}\omega + \omega \wedge \omega)$ and Lemma 2.1, we have

Proposition 2.1. *The curvature form Ω of ∇ is given by $\Omega = \bar{\partial}\omega + \partial_{\mathcal{V}}\omega = \bar{\partial}_{\mathcal{H}}\omega + \bar{\partial}_{\mathcal{V}}\omega + \partial_{\mathcal{V}}\omega$:*

$$(2.9) \quad \Omega_j^i = \sum_{\alpha,\beta} R_{j\alpha\beta}^i dz^{\alpha} \wedge d\bar{z}^{\beta} + \sum_{\alpha,k} R_{j\alpha\bar{k}}^i dz^{\alpha} \wedge \bar{\theta}^k + \sum_{\alpha,k} R_{j\alpha k}^i dz^{\alpha} \wedge \theta^k,$$

where we put $R_{j\alpha\bar{\beta}}^i = -X_{\bar{\beta}}\Gamma_{ja}^i$, $R_{j\alpha\bar{k}}^i = -Y_{\bar{k}}\Gamma_{ja}^i$, $R_{j\alpha k}^i = -Y_k\Gamma_{ja}^i$.

Remark 2.2. From this lemma, we can easily infer the identity $\partial_{\mathcal{H}}^2 \equiv 0$.

Here we describe a special class of Finsler structures. A complex Finsler bundle (E, F) is said to be *modeled on a complex Minkowski space* if its connection ∇ is projectable to a connection of E , that is, its connection coefficients $\Gamma_{j\alpha}^i$ are functions of position of $z \in M$ alone. Then we have

Theorem 2.1 ([2]). *Let (E, F) be modeled on a complex Minkowski space. Then there exists a Hermitian structure h_F on E , and the Finsler connection ∇ of (E, F) is given by the pull-back of the Hermitian connection of (E, h_F) .*

It is trivial that (E, F) is modeled on a complex Minkowski space if and only if $\partial_{\mathcal{V}}\omega = \bar{\partial}_{\mathcal{V}}\omega = 0$, and, in this case, the first term of (2.9) is given by the curvature of its associated h_F .

3. Conformally flat Finsler structures

We begin with the following

Definition 3.1. A complex Finsler structure F is said to be *flat* if there exists an open covering $\{U\}$ and a suitably chosen holomorphic frame field $s_U = \{s_1, \dots, s_r\}$ on each U such that with respect to $\{U, s_U\}$ the function F is independent at the base point $z \in M : F = F(\xi)$. Such a covering $\{U, s_U\}$ is said to be *adapted*.

This notion is a complex analogue of a *locally Minkowski space* in real Finsler geometry (cf. Definition 24.1 in [9]). We shall use, however, the term *flat* since the following theorem holds:

Theorem 3.1. *A complex Finsler structure F is flat if and only if its Finsler connection ∇ is flat, that is, the curvature Ω of ∇ vanishes identically.*

PROOF. We suppose that F is independent at $z \in M$ with respect to an adapted $\{U, s_U\}$. Then, from (2.5), we have $N_{\alpha}^i = 0$, and so by virtue of (2.8) we get $\Gamma_{j\alpha}^i = 0$. Hence, with respect to an adapted $\{U, s_U\}$ the connection form ω vanishes on each U . This means the vanishing of its curvature.

Conversely we assume that the curvature Ω of ∇ vanishes identically. Then, (E, F) is modeled on a complex Minkowski space, and, by Theorem 2.1, ∇ is the Hermitian connection of the associated h_F . Since $\omega = 0$

is completely integrable, on a suitable open neighborhood U of each point $z \in M$, we can introduce a parallel frame field $s_U = \{s_1, \dots, s_r\}$. Since $ds_j = 0$, we have $\bar{\partial}s_j = 0$, and so s_U is holomorphic. Therefore the connection form ω with respect to $\{U, s_U\}$ vanishes on each U . Hence, by virtue of $\sum \xi^j \Gamma_{j\alpha}^i = N_\alpha^i$, we have $N_\alpha^i = 0$. Consequently, from (2.6) we have $F = F(\xi)$ with respect to $\{U, s_U\}$. $\square$

By this theorem and Theorem 2.1, we have

Proposition 3.1. *A convex Finsler structure F on E is flat if and only if (E, F) is modeled on a complex Minkowski space and its associated Hermitian metric h_F is flat.*

By this proposition, we know that if (E, F) is a flat Finsler vector bundle, then it admits a flat Hermitian structure h_F . Conversely, the norm function derived from a flat Hermitian structure is also a flat Finsler structure. Hence we have (cf. Proposition 4.21 on p. 14 of [7])

Theorem 3.2. *The following conditions are equivalent:*

- (1) E admits a flat Finsler structure.
- (2) E admits a flat unitary structure.
- (3) E is defined by a representation $\rho : \pi_1(M) \rightarrow U(r)$: $E \cong \tilde{M} \times_\rho \mathbb{C}^r$, where $\pi_1(M)$ is the fundamental group of M and $\tilde{M}$ is the universal covering of M .

We shall consider a conformal rescaling $F \rightarrow \tilde{F} = e^{\sigma(z)} F$ of the Finsler metric F for a differentiable function $\sigma(z)$ on M . We shall calculate the connection form $\tilde{\omega}$ of $(E, \tilde{F})$. Because of $\tilde{F}_{i\bar{j}} = e^{\sigma(z)} F_{i\bar{j}}$ and (2.5), the non-linear connection is changed as $\tilde{N}_\alpha^i = N_\alpha^i + (\partial\sigma/\partial z^\alpha) \xi^i$. Hence, by (2.8) we get

$$(3.1) \quad \tilde{\omega} = \omega + \partial\sigma \otimes I_{\mathcal{V}}$$

for the identity endomorphism $I_{\mathcal{V}}$ of $\mathcal{V}$. Then we have

Lemma 3.1. *Let (E, F) be modeled on a complex Minkowski space with an associated Hermitian structure h_F . Then, for any conformal rescaling $F \rightarrow \tilde{F} = e^{\sigma(z)} F$, $(E, \tilde{F})$ is also modeled on a complex Minkowski space, and the conformal rescaling $e^{\sigma(z)} h_F$ of h_F associates with $(E, \tilde{F})$.*

PROOF. The Finsler connection $\tilde{\nabla}$ of $(E, \tilde{F})$ is given by (3.1). Hence the first assertion is trivial. Moreover, we have

$$\tilde{\omega} = \omega + \partial\sigma \otimes I_{\mathcal{V}} = (e^{\sigma(z)} h_F)^{-1} \partial(e^{\sigma(z)} h_F) = h_{\tilde{F}}^{-1} \partial h_{\tilde{F}}.$$

Hence $e^{\sigma(z)}h_F$ associates with $(E, \tilde{F})$. $\square$

From (3.1) and Proposition 2.1, the curvature form is transformed as follows:

$$(3.2) \quad \tilde{\Omega} = \Omega + \bar{\partial}\partial\sigma \otimes I_{\mathcal{V}}.$$

From (3.1), the $(1,0)$ -form θ^j is transformed as $\tilde{\theta}^j = \theta^j + \xi^j \otimes \partial\sigma$. Then we have

Lemma 3.2. *By any conformal rescaling $F \rightarrow \tilde{F} = e^{\sigma(z)}F$, the forms $\partial_{\mathcal{V}}\omega$ and $\bar{\partial}_{\mathcal{V}}\omega$ are invariant.*

PROOF. By definition,

$$\partial_{\mathcal{V}}\omega_j^i = \sum_{\alpha,k} \frac{\partial \Gamma_{j\alpha}^i}{\partial \xi^k} \theta^k \wedge dz^\alpha.$$

With respect to the new function $\tilde{F}$, we shall compute the right hand side:

$$\begin{aligned} \sum_{\alpha,k} \frac{\partial \tilde{\Gamma}_{j\alpha}^i}{\partial \xi^k} \tilde{\theta}^k \wedge dz^\alpha &= \sum_{\alpha,k} \frac{\partial}{\partial \xi^k} \left(\Gamma_{j\alpha}^i + \frac{\partial \sigma}{\partial z^\alpha} \right) (\theta^k + \xi^k \partial\sigma) \wedge dz^\alpha \\ &= \sum_{\alpha,k} \frac{\partial \Gamma_{j\alpha}^i}{\partial \xi^k} \theta^k \wedge dz^\alpha, \end{aligned}$$

since, by the homogeneity of F , we have $\sum_k \xi^k Y_k \Gamma_{j\alpha}^i \equiv 0$. This means that the form $\partial_{\mathcal{V}}\omega$ is invariant under any conformal rescaling.

The proof for the invariance of $\bar{\partial}_{\mathcal{V}}\omega$ is similar. $\square$

By this lemma, we have

Proposition 3.2. *The $\text{End}(\mathcal{V})$ -valued $(1,1)$ -form*

$$(3.3) \quad \Theta = \Omega - \frac{1}{r} \rho \otimes I_{\mathcal{V}}$$

is invariant by any conformal rescaling, where ρ is defined by $\rho = \text{Tr}.\bar{\partial}_{\mathcal{H}}\omega$.

PROOF. By Lemma 3.1, the forms $\partial_{\mathcal{V}}\omega$ and $\bar{\partial}_{\mathcal{V}}\omega$ are invariant by any conformal rescaling. Hence the $(1,1)$ -form ρ is transformed as

$$\begin{aligned} \tilde{\rho} &= \text{Tr}.\tilde{\Omega} - \partial_{\mathcal{V}}\omega - \bar{\partial}_{\mathcal{V}}\omega = \text{Tr}.\tilde{\Omega} - \partial_{\mathcal{V}}\omega - \bar{\partial}_{\mathcal{V}}\omega \\ &= \text{Tr}.\partial_{\mathcal{H}}\omega + r\bar{\partial}\partial\sigma = \rho + r\bar{\partial}\partial\sigma. \end{aligned}$$

Thus we have $\bar{\partial}\partial\sigma = (\tilde{\rho} - \rho)/r$. Substituting this into (3.2), we see that Θ is invariant by any conformal rescaling. $\square$

We call the Θ given by (3.2) the *conformal curvature* of (E, F) . It is trivial that if Ω vanishes, then Θ also vanishes.

Definition 3.2. A Finsler structure F is (locally) *conformally flat* if every $z \in M$ has an open neighborhood U and a differentiable function $\sigma_U : U \rightarrow \mathbb{R}$ such that $\tilde{F}_U = e^{\sigma_U} F$ is a flat Finsler structure on U .

Now we shall prove our main theorem:

Theorem 3.3. *Let F be a convex Finsler structure on a holomorphic vector bundle E . Then F is conformally flat if and only if the conformal curvature Θ vanishes identically.*

PROOF. We shall fix a frame field $\{U, s_U\}$ for E , and use the local expressions with respect to $\{U, s_U\}$.

We suppose that Θ vanishes identically. Then, since $\partial_{\mathcal{V}}\omega = \bar{\partial}_{\mathcal{V}}\omega = 0$, (E, F) is modeled on a complex Minkowski space. By Theorem 2.1 there exists an associated Hermitian structure h_F , and Ω is given by the pull-back of the one Ω_F of h_F . Hence ρ is the Ricci curvature of h_F :

$$\rho = \bar{\partial}\partial \log \det(h_{i\bar{j}}),$$

where we put $h_{i\bar{j}} = h_F(s_i, s_j)$. On each U , we put $\sigma_U(z) = \frac{1}{r} \log \det(h_{i\bar{j}})$, and we consider the conformal rescaling $F \rightarrow \tilde{F}_U = e^{\sigma_U(z)} F|_U$. Then, $\tilde{F}_U$ is also modeled on a complex Minkowski space, and its curvature $\tilde{\Omega}$ is given by

$$\begin{aligned} \tilde{\Omega} &= \frac{1}{r} \tilde{\rho} \otimes I_{\mathcal{V}} = \frac{1}{r} (Tr.\Omega_F + r\bar{\partial}\partial\sigma_U) \otimes I_{\mathcal{V}} \\ &= \frac{1}{r} (-\bar{\partial}\partial\sigma_U + \bar{\partial}\partial\sigma_U) \otimes I_{\mathcal{V}} = 0, \end{aligned}$$

which shows that $\tilde{F}_U$ is flat. Hence F is conformally flat.

The converse is trivial. $\square$

The conformal flatness of a Hermitian structure has been studied in [10], where the conformal flatness of a Hermitian structure has been characterized by the vanishing of a conformally invariant curvature tensor. Our conformal curvature Θ coincides with that of Matsuo if the given Finsler structure F is the norm function associated to a Hermitian structure h , that is, $F(z, \xi) = \sum h_{i\bar{j}}(z)\xi^i\bar{\xi}^j$. Then, from Theorem 3.3, we have

Proposition 3.3. *A convex Finsler structure on E is conformally flat if and only if (E, F) is modeled on a complex Minkowski space, and its associated Hermitian structure is conformally flat.*

Let P be the $GL(r, \mathbb{C})$ -principal bundle associated to E . We denote by $PGL(r, \mathbb{C})$ the projective linear group $GL(r, \mathbb{C})/\mathbb{C}^*I_r$, where $\mathbb{C}^*I_r$ is the center of $GL(r, \mathbb{C})$. A vector bundle E is said to be *projectively flat* if the $PGL(r, \mathbb{C})$ -principal bundle $\hat{P} = P/\mathbb{C}^*I_r$ is provided with a flat structure (cf. [7]).

If the conformal curvature satisfies $\Theta \equiv 0$, then the curvature Ω of the associated h_F satisfies

$$\Omega = \frac{1}{r} \rho \otimes I_V.$$

Hence, according to Proposition 2.8 in [7], the bundle E is projectively flat, and $\hat{P}$ is defined by a representation $\rho : \pi_1(M) \rightarrow PU(r)$, where $PU(r) = U(r)/U(1)I_r$ is the projective unitary group. This means that, if we consider the universal covering space $\tilde{M}$ as a $\pi_1(M)$ -principal bundle $\tilde{M} \rightarrow M$, the bundle $\hat{P}$ is defined by the representation $\rho : \pi_1(M) \rightarrow PU(r)$. The flat structure of $\hat{P}$ is induced by the natural flat structure of $\tilde{M} \rightarrow M$.

By Proposition 3.3, we know that, if (E, F) is a conformally flat Finsler vector bundle, it admits a conformally flat Hermitian structure h_F . Conversely, the norm function defined by a conformally flat Hermitian structure is also a conformally flat Finsler structure. Hence we have (cf. Proposition 4.22 in p. 14 of [7])

Theorem 3.4. *The following conditions are equivalent:*

- (1) *E admits a conformally flat Finsler structure.*
- (2) *E admits a conformally flat Hermitian structure.*
- (3) *The bundle $\hat{P} = P/\mathbb{C}^*I_r$ is defined by a representation $\rho : \pi_1(M) \rightarrow PU(r) : \hat{P} \cong \tilde{M} \times_\rho PU(r)$.*

Example 3.1. Let M be a so-called *Hopf manifold* $\{\mathbb{C}^n - 0\}/\Delta_\lambda$, where Δ_λ is the group generated by the holomorphic transformations $(z^1, \dots, z^n) \rightarrow (\lambda z^1, \dots, \lambda z^n)$ on $\mathbb{C}^n - \{0\}$ for $\lambda \in \mathbb{C}$, $0 < |\lambda| < 1$. Then there exists a standard Hermitian structure on TM :

$$(3.3) \quad ds^2 = \frac{1}{\|z\|^2} \sum_{\alpha} dz^{\alpha} \otimes d\bar{z}^{\alpha} = e^{-\log \|z\|^2} \sum_{\alpha} dz^{\alpha} \otimes d\bar{z}^{\alpha},$$

where $\|z\|^2 = \sum_{\alpha} z^{\alpha} \bar{z}^{\alpha}$. This metric is locally conformal Kähler-flat (l.c.K₀ in short, cf. [12]). Its Hermitian connection is given by

$$(3.4) \quad \omega = -\partial(\log \|z\|^2) \otimes I_{TM}.$$

The norm function defined by the metric above is $F_0(z, \xi) = e^{-\log \|z\|^2} \|\xi\|^2$.

To obtain a conformally flat Finsler structure F , we shall modify F_0 into the form

$$(3.5) \quad F(z, \xi) = e^{-\log \|z\|^2} f(\xi)$$

for a positive function $f(\xi)$ on $\mathbb{C}^n$ satisfying $f(\lambda\xi) = |\lambda|^2 f(\xi)$ and the Hermitian matrix $(\partial^2 f / \partial \xi^i \partial \bar{\xi}^j)$ is positive definite. Since F is also invariant by the action of Δ_{λ} , it defines a convex Finsler structure on TM . It is trivial that this Finsler structure F is conformally flat. We shall check this by computing its conformal curvature Θ .

This complex Finsler manifold (M, F) is modeled on a complex Minkowski space, and its associated Hermitian metric is given by (3.1). We shall show this. If we put $f_{i\bar{j}}(\xi) = \partial^2 f / \partial \xi^i \partial \bar{\xi}^j$, we have $F_{i\bar{j}} = e^{-\log \|z\|^2} \times f_{i\bar{j}}(\xi)$. Hence, by (2.5), the non-linear connection N_{α}^i of (M, F) is given by

$$(3.6) \quad N_{\alpha}^i = -\frac{\bar{z}^{\alpha}}{\|z\|^2} \xi^i.$$

Moreover, from (2.8), the connection coefficients of ∇ are given by

$$\Gamma_{j\alpha}^i = -\frac{\partial \log \|z\|^2}{\partial z^{\alpha}} \delta_j^i.$$

Hence the Finsler connection ∇ is given by (3.4).

Since the curvature form Ω of ∇ is given by $\Omega = -\bar{\partial}\partial(\log \|z\|^2) \otimes I_{TM}$, we get $\rho = -n\bar{\partial}\partial(\log \|z\|^2)$. From these equations and the definition of Θ , we get $\Theta \equiv 0$.

Let (E, F) be a complex Finsler bundle over a compact Kähler manifold (M, g) . Assume that (E, F) is conformally flat. Then, since $\Theta = 0$, (E, F) is modeled on a complex Minkowski space, and its curvature Ω is given by $\Omega = \frac{1}{r}\rho \otimes I_{\mathcal{V}}$. Now, it is easily proved that the associated Hermitian vector bundle (E, h_F) satisfies the *weak Einstein condition*. Moreover, if (M, g) is compact Kähler, by suitable conformal rescaling $h_F \rightarrow ah_F$, we can obtain that ϕ is constant (cf. Proposition 2.4 in Chapter IV of [7]). Hence the associated Hermitian bundle (E, h_F) is Einstein–Hermitian over (M, g) . Consequently we have

Proposition 3.4. *Let (E, F) be a convex Finsler vector bundle over a compact Kähler manifold (M, g) . If (E, F) is conformally flat, then the associated Hermitian vector bundle (E, h_F) satisfies the Einstein condition.*

4. Some remarks from Hermitian geometry

By Proposition 3.3, some geometric properties of a conformally flat (E, F) are obtained from those of (E, h_F) . We shall show some results directly obtained from Hermitian or Kählerian geometry.

Let M be a complex manifold of $\dim_{\mathbb{C}} M = n$, and F a convex Finsler structure on TM . The pair (M, F) is called a *complex Finsler manifold*. Suppose that (M, F) is conformally flat. Then (TM, F) is modeled on a complex Minkowski space, and its associated Hermitian metric h_F is conformally flat. Hence, there exists an open covering $\{U\}$ and a family of local functions $\{\sigma_U\}$ such that $h_U = e^{\sigma_U} h_F$ is a flat metric on U . Moreover, if each h_U is a flat Kähler metric on U , (M, h_F) is l.c.K₀. (Example 1 is just of this type). Then, applying Theorem 2.2 in [12] (see also Theorem 6.8 in [11]), we see that the universal covering $\tilde{M}$ of M is $\mathbb{C}^n - \{0\}$, and h_F is globally conformal to the metric induced by (3.3). Applying this fact, we have

Theorem 4.1. *Let (M, F) be a compact complex Finsler manifold of $\dim_{\mathbb{C}} M = n$ which is conformally flat. Suppose that its associated Hermitian manifold (M, h_F) is (not globally) l.c.K₀. Then the universal covering $\tilde{M}$ of M is given by $\mathbb{C}^n - \{0\}$, and F is globally conformal to the Finsler structure induced by the one of the form (3.5).*

PROOF. The fact that $\tilde{M} = \mathbb{C}^n - \{0\}$ is trivial from Vaisman's theorem. We shall prove the second part of the theorem. Since the Finsler connection ∇ of (M, F) is given by the form ω in (3.2), its non-linear connection N_j^i is given by (3.6). Now, from (2.6), we have $X_\alpha F = 0$. Hence, in this case, we have

$$\frac{\partial F}{\partial z^\alpha} + \frac{\bar{z}^\alpha}{\|z\|^2} F = 0.$$

This equation implies

$$\frac{\partial}{\partial z^\alpha} \left(e^{\log \|z\|^2} F \right) = \frac{\partial}{\partial z^\alpha} (\|z\|^2 F) = \bar{z}^\alpha F + \|z\|^2 \frac{\partial F}{\partial z^\alpha} = 0.$$

In the same way as above, we have $\partial \left(e^{\log \|z\|^2} F \right) / \partial \bar{z}^\alpha = 0$. Hence we get

$$e^{\log \|z\|^2} F = f(\xi)$$

for a function f which depends only on ξ . It is trivial that f satisfies the homogeneity and the convexity conditions. Consequently F must be of the type (3.5). $\square$

In a previous paper [2], we have introduced the notion of *Finsler-Kähler manifold*. We shall recall its definition. We use the Greek letters $j, k, \dots$ for the indices of the local coordinates of M . If a complex Finsler manifold (M, F) is given, its non-linear connection N_j^i is given by (2.5):

$$N_j^i = \sum_{l, m} F^{i\bar{m}} \frac{\partial F_{l\bar{m}}}{\partial z^j} \xi^l,$$

and the connection coefficients of its Finsler connection ∇ are given by $\Gamma_{jk}^i = \partial N_k^i / \partial \xi^j$. Then (M, F) is said to be Finsler-Kähler if the condition

$$\Gamma_{jk}^i = \Gamma_{kj}^i$$

is satisfied. In [1], such a manifold is called *strongly Finsler-Kähler*. By Theorem 2.1 it is trivial that, if a Finsler-Kähler manifold (M, F) is modeled on a complex Minkowski space, then its associated h_F is Kähler.

Any conformally flat Kähler manifold is flat (cf. Theorem 4.1 in [13], see also Corollary 4.3 in [10]). In our case, we have

Theorem 4.2. *Let (M, F) be a Finsler-Kähler manifold. If (M, F) is conformally flat, then (M, F) is flat.*

PROOF. By Proposition 3.3, (M, F) is conformally flat if and only if it is modeled on a complex Minkowski space, and moreover its associated h_F is conformally flat. By the assumption of Finsler-Kähler, the associated (M, h_F) is a conformally flat Kähler manifold. Hence (M, h_F) is flat. Consequently, by Proposition 3.1, (M, F) is flat. $\square$

Acknowledgements. The author wishes to express his thanks to Prof. Dr. MAKOTO MATSUMOTO and to Prof. Dr. YOSHIHIRO ICHIJO for their several suggestions for the first version of the present paper. In particular, Prof. ICHIJO has kindly communicated YANO and MOGI's paper [13] to the author.

References

- [1] M. ABATE and G. PATRIZIO, Finsler metrics – A Global Approach with Applications to Geometric Function Theory, Lecture Notes 1591, *Springer*, 1994.
- [2] T. AIKOU, Complex manifolds modeled on a complex Minkowski space, *J. Math. Kyoto Univ.* **35** (1995), 83–101.
- [3] T. AIKOU, Some remarks on locally conformal complex Berwald spaces, *Contemporary Mathematics* **196** (1996), 109–120.
- [4] T. AIKOU, Einstein–Finsler vector bundles, *Publ. Math. Debrecen* **51** (1997), 363–384.
- [5] T. AIKOU, A partial connection on complex Finsler bundles and its applications (to appear in *Illinois J. Math.*, 1998).
- [6] S. KOBAYASHI, Negative vector bundles and complex Finsler structures, *Nagoya Math. J.* **57** (1975), 153–166.
- [7] S. KOBAYASHI, Differential Geometry of Complex Vector Bundles, *Iwanami–Princeton Univ. Press*, 1987.
- [8] S. KOBAYASHI, Complex Finsler vector bundles, *Contemporary Mathematics* **196** (1996), 145–153.
- [9] M. MATSUMOTO, Foundations of Finsler Geometry and Special Finsler Spaces, *Kaiseisha Press, Otsu, Japan*, 1986.
- [10] K. MATSUO, On local conformal Hermitian-flatness of Hermitian manifolds, *Tokyo J. Math.* **19** (1996), 499–515.
- [11] L. ORNEA, Locally Conformal Kaehler Manifolds – A survey, *Università degli di Roma “La Sapienza”, Roma, Maggio*, 1994.
- [12] I. VAISMAN, Generalized Hopf manifolds, *Geom. Dedicata* **13** (1982), 231–255.
- [13] K. YANO and I. MOGI, On real representations of Kaehlerian manifolds, *Ann. of Math.* **61** (1955), 170–189.

TADASHI AIKOU
 DEPARTMENT OF MATHEMATICS AND COMPUTER SCIENCE
 KAGOSHIMA UNIVERSITY
 1-2-35 KORIMOTO KAGOSHIMA, 890
 JAPAN

E-mail: aikou@sci.kagoshima-u.ac.jp

(Received January 16, 1998)