

Differentiable loops on the real line

By PÉTER T. NAGY (Debrecen) and IZABELLA STUHL (Debrecen)

Dedicated to Professor Adalbert Bowdi on his 70th birthday

Abstract. The paper is devoted to the study of differentiable loops L on the real line such that the group G topologically generated by the left translations is locally compact and hence it is isomorphic to the universal covering group of $PSL_2(\mathbb{R})$. Using the methods developed in [3] we introduce a class of natural parametrizations of the loop manifold L corresponding to the Iwasawa decompositions of G and find explicit expressions for the loop multiplication with respect to the given parametrizations. We characterize the differentiable curves $\mathbb{R} \rightarrow G$ consisting of the left translations of a loop L in the biinvariant Lorentzian geometry of G .

1. Introduction

An algebra $(L, \circ, e)$ with a binary operation $(x, y) \mapsto x \circ y : L \times L \rightarrow L$ and unit element $e \in L$ is called a *left loop* if for any given $a, b \in L$ the equation $a \circ x = b$ is uniquely solvable for x , and if $e \in L$ satisfies $e \circ x = x \circ e = x$ for any $x \in L$. The left loop $(L, \circ, e)$ is called a *loop* if for any given $a, b \in L$ the equation $x \circ a = b$ is uniquely solvable, too. The operation $\circ : L \times L \rightarrow L$ is called *multiplication*, the solution of the equation $a \circ x = b$ is denoted by $a \backslash b$, the solution of the equation $x \circ a = b$ is denoted by a / b . The binary *left division* operation $(a, b) \mapsto a \backslash b$ can be defined in a left loop, the binary *right division* operation $(a, b) \mapsto a / b$ can be defined in a loop. The bijective maps $\lambda_x : L \rightarrow L : y \mapsto x \circ y$, $x \in L$, are called *left translations* of $(L, \circ, e)$. The group G generated by the

Mathematics Subject Classification: 20N05, 22E15, 58D19.

Key words and phrases: $PSL_2(\mathbb{R})$, differentiable loop, group action, continuous section.

set $\Lambda = \{\lambda_x, x \in L\}$ of left translations acts transitively on L . Hence there is a bijective correspondence $L \rightarrow G/H : x \mapsto \lambda_x H$, where H is the stabilizer of $e \in L$ in the group G .

A loop $(L, \circ, e)$ is called *differentiable* of class C^r if L is a differentiable manifold of class C^r and the multiplication $\circ$, the left and the right divisions of $(L, \circ)$ are C^r -differentiable maps $L \times L \rightarrow L$. In the study of the Lie theory of differentiable loops is very fruitful the investigation of the differentiable submanifold $\Lambda = \{\lambda_x, x \in L\} \subset G$ of the group topologically generated by the left translations (cf. [3]), the mapping $x \mapsto \lambda_x : L \rightarrow G$ corresponds to a section with respect to the projection map $G \rightarrow G/H$.

Differentiable loops defined on the unit circle having a locally compact group G as the group topologically generated by the left translations, were studied in Section 18 of the monograph [3] by P. T. NAGY and KARL STRAMBACH. Particularly they proved the following assertion (Proposition 18.2, p. 235.) by an application of a theorem of BROUWER [1]:

Let L be a topological proper loop on a connected 1-manifold, such that the group G topologically generated by the left translations is locally compact. Then G is a finite covering of the group $PSL_2(\mathbb{R})$ if L is a circle, and G is the universal covering of $PSL_2(\mathbb{R})$ if L is homeomorphic to the real line $\mathbb{R}$.

Section 18 of [3] contains a detailed investigation of differentiable loops on the circle for which the group G topologically generated by the left translations is isomorphic to the group $PSL_2(\mathbb{R})$. The corresponding loop multiplication were determined by a pair of real periodic differentiable functions satisfying the conditions some differential inequality. The investigation of differentiable loops on the circle is extended to the case where the group G is isomorphic to $SL_2(\mathbb{R})$ in [4]. This paper is devoted to the study of differentiable proper loops on the real line $\mathbb{R}$ such that the group G topologically generated by the left translations is locally compact. In this case the group G is isomorphic to the universal covering group of $PSL_2(\mathbb{R})$. Modifying the description of [3] we give explicit expressions of the loop multiplications with respect to a class of natural parametrizations of the loop manifold L corresponding to the Iwasawa decompositions of G . We characterize within the biinvariant Lorentzian geometry of the group G the one-dimensional differentiable submanifolds $\Lambda = \{\lambda_x, x \in \mathbb{R}\}$ of $PSL_2(\mathbb{R})$, the point of which are the left translations of a differentiable loop on the real line $\mathbb{R}$.

2. Action of $\widetilde{PSL}_2(\mathbb{R})$ on $\mathbb{R}$

Let $\mathcal{A}$ be the algebra of real 2×2 matrices. We denote by $\widetilde{PSL}_2(\mathbb{R})$ the universal covering group of the groups

$$SL_2(\mathbb{R}) = \{A \in \mathcal{A}; \det A = 1\}, \quad PSL_2(\mathbb{R}) = \{\pm A \in \mathcal{A}; \det A = 1\}.$$

The matrices

$$\mathbf{H} = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \quad \mathbf{T} = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \mathbf{U} = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$$

form a basis of the Lie algebra $\mathfrak{sl}_2(\mathbb{R})$ of $SL_2(\mathbb{R})$. Let $\kappa : X \mapsto \frac{1}{2}\text{Trace}(X^2)$ be the normalized Cartan–Killing form on $\mathfrak{sl}_2(\mathbb{R})$ given by $\kappa(X) = -\det(X) = h^2 + t^2 - u^2$ for $X = h\mathbf{H} + t\mathbf{T} + u\mathbf{U}$, which determines a pseudo-euclidean vector space structure on $\mathfrak{sl}_2(\mathbb{R})$ such that the unit vectors $\mathbf{H}, \mathbf{T}, \mathbf{U}$ form a pseudo-orthonormal basis. We denote by $\exp : \mathfrak{sl}_2(\mathbb{R}) \rightarrow SL_2(\mathbb{R})$ the exponential map of $SL_2(\mathbb{R})$ and by $\widetilde{\exp} : \mathfrak{sl}_2(\mathbb{R}) \rightarrow \widetilde{PSL}_2(\mathbb{R})$ the exponential map of the universal covering group $\widetilde{PSL}_2(\mathbb{R})$. Let $p : \widetilde{PSL}_2(\mathbb{R}) \rightarrow SL_2(\mathbb{R})$ be the covering map. The connected 1-parameter subgroups $\{\widetilde{\exp}(t\mathbf{X}); t \in \mathbb{R}\}$ with $\mathbf{X} \neq \mathbf{0}$ are called *elliptic*, *hyperbolic* or *parabolic*, if $\kappa(\mathbf{X}) < 0$, $\kappa(\mathbf{X}) > 0$ or $\kappa(\mathbf{X}) = 0$, respectively. The covering map p induces isomorphisms $\{\widetilde{\exp}(t\mathbf{X}); t \in \mathbb{R}\} \rightarrow \{\exp(t\mathbf{X}); t \in \mathbb{R}\}$ between hyperbolic or parabolic 1-parameter subgroups, it yields proper covering homomorphisms for elliptic 1-parameter subgroups. The connected 2-dimensional subgroups of $\widetilde{PSL}_2(\mathbb{R})$ have the form $\{\widetilde{\exp}(s(\mathbf{Y} - \mathbf{Z})) \cdot \widetilde{\exp}(t\mathbf{X}); s, t \in \mathbb{R}\}$, where $(\mathbf{X}, \mathbf{Y}, \mathbf{Z})$ is a pseudo-orthonormal basis of $(\mathfrak{sl}_2(\mathbb{R}), \kappa)$, which is conjugate to the basis $(\mathbf{H}, \mathbf{T}, \mathbf{U})$ of $\mathfrak{sl}_2(\mathbb{R})$. Clearly, the covering map p induces isomorphisms

$$\{\widetilde{\exp}(s(\mathbf{Y} - \mathbf{Z})) \cdot \widetilde{\exp}(t\mathbf{X}); s, t \in \mathbb{R}\} \rightarrow \{\exp(s(\mathbf{Y} - \mathbf{Z})) \exp(t\mathbf{X}); s, t \in \mathbb{R}\}$$

between 2-dimensional subgroups.

The pseudo-euclidean scalar product induced by the Cartan–Killing form on $\mathfrak{sl}_2(\mathbb{R})$ is invariant with respect to the adjoint representation of $\widetilde{PSL}_2(\mathbb{R})$. Hence it can be extended by left or right translations of $\widetilde{PSL}_2(\mathbb{R})$ to a bi-invariant pseudo-riemannian metric $g(X, Y)$ on the manifold $\widetilde{PSL}_2(\mathbb{R})$ such that $g(X, X)$ at the identity element of $\widetilde{PSL}_2(\mathbb{R})$ coincides with $\kappa(X)$. A curve $t \mapsto \gamma(t) : \mathbb{R} \rightarrow \widetilde{PSL}_2(\mathbb{R})$ will be called *time-like* if its tangent vectors $\dot{\gamma}(t)$ satisfy $g(\dot{\gamma}(t), \dot{\gamma}(t)) < 0$ for any $t \in \mathbb{R}$.

We consider the group $G = \widetilde{PSL}_2(\mathbb{R})$, a 2-dimensional subgroup $H \subset G$ and the homogeneous space G/H diffeomorphic to $\mathbb{R}$. Let $(\mathbf{X}, \mathbf{Y}, \mathbf{Z})$ be a pseudo-orthonormal basis of $\mathfrak{sl}_2(\mathbb{R})$, for which there exists an automorphism $\beta : \mathfrak{sl}_2(\mathbb{R}) \rightarrow \mathfrak{sl}_2(\mathbb{R})$ mapping the basis $(\mathbf{X}, \mathbf{Y}, \mathbf{Z})$ onto the canonical basis $(\mathbf{H}, \mathbf{T}, \mathbf{U})$, such that

$$H = \{\widetilde{\exp}(s(\mathbf{Y} - \mathbf{Z})) \cdot \widetilde{\exp}(t\mathbf{X}); s, t \in \mathbb{R}\} \subset G.$$

Let K, N and D denote the 1-parameter subgroups $K = \{\widetilde{\exp}(t\mathbf{Z}); t \in \mathbb{R}\}$, $N = \{\widetilde{\exp}(t(\mathbf{Y} - \mathbf{Z})); t \in \mathbb{R}\}$ and $D = \{\widetilde{\exp}(t\mathbf{X}); t \in \mathbb{R}\}$. Since the vector space direct sum $\mathfrak{sl}_2(\mathbb{R}) = \mathbb{R}\mathbf{Z} + \mathbb{R}(\mathbf{Y} - \mathbf{Z}) + \mathbb{R}\mathbf{X}$ is an Iwasawa decomposition of the Lie algebra $\mathfrak{sl}_2(\mathbb{R})$, (cf. e.g. [2] p. 234), the map

$$\Delta : g \mapsto (k, n, d) : \widetilde{PSL}_2(\mathbb{R}) \rightarrow K \times N \times D \quad (1)$$

defined by $g = knd$ is an analytic diffeomorphism of $\widetilde{PSL}_2(\mathbb{R})$ onto the product manifold $K \times N \times D$. The triple $\Delta(g) = (k, n, d)$ will be called the Iwasawa decomposition of the element $g \in \widetilde{PSL}_2(\mathbb{R})$. It follows that any element $g \in \widetilde{PSL}_2(\mathbb{R})$ can be decomposed uniquely into the product $g = \widetilde{\exp}(t\mathbf{Z})\widetilde{\exp}(u(\mathbf{Y} - \mathbf{Z}))\widetilde{\exp}(v\mathbf{X})$ with $(t, u, v) \in \mathbb{R}^3$. Using the decomposition $G = K \cdot N \cdot D$ and $H = N \cdot D$ we obtain that any coset of the subgroup $H = N \cdot D$ can be represented uniquely by an element $\widetilde{\exp}(t\mathbf{Z}) \in K$, $t \in \mathbb{R}$. Hence the manifold G/H can be parametrized by elements of K . Since the mapping $t \mapsto \widetilde{\exp}(t\mathbf{Z}) : \mathbb{R} \rightarrow K$ is bijective, we obtain that

$$\phi : t \mapsto \widetilde{\exp}(t\mathbf{Z})H : \mathbb{R} \rightarrow G/H \quad (2)$$

is a diffeomorphism and hence this map gives a global real parametrization of the loop manifold which will be called an *Iwasawa parametrization*. This parametrization ϕ is not canonical in the sense that it depends on the choice of the basis $(\mathbf{X}, \mathbf{Y}, \mathbf{Z})$ of $\mathfrak{sl}_2(\mathbb{R})$.

We can define the action $\alpha_g : \mathbb{R} \mapsto \mathbb{R}$ of $g \in G$ by

$$\alpha_g t = \phi^{-1}(g \cdot \widetilde{\exp}(t\mathbf{Z})H) = \phi^{-1}(g \cdot \phi t),$$

where $kH \mapsto g \cdot kH : G/H \rightarrow G/H$ denotes the natural action of G on the factor space G/H . Hence $\alpha : G \times \mathbb{R} \rightarrow \mathbb{R}$ is a natural differentiable transformation group action of $G = \widetilde{PSL}_2(\mathbb{R})$ on $\mathbb{R}$.

Proposition 1. *The mapping $\alpha_g : \mathbb{R} \mapsto \mathbb{R}$ with $g = \widetilde{\exp}(t\mathbf{Z}) \cdot \widetilde{\exp}(u(\mathbf{Y} - \mathbf{Z})) \cdot \widetilde{\exp}(v\mathbf{X})$ with respect to the Iwasawa parametrization (2) can be expressed by*

$$\alpha_g s = \begin{cases} t + \operatorname{arccot}(2u + e^{-2v} \cot s) + i\pi, & \text{if } i\pi < s < (i+1)\pi, \\ t + i\pi, & \text{if } s = i\pi, \end{cases}$$

for any $i \in \mathbb{Z}$. In particular, $\alpha_{\widetilde{\exp}(t\mathbf{Z})} s = t + s$.

PROOF. The equation $\alpha_{\widetilde{\exp}(t\mathbf{Z})}s = t + s$ follows from the relation

$$\widetilde{\exp}(t\mathbf{Z}) \cdot \widetilde{\exp}(s\mathbf{Z}) \cdot H = \widetilde{\exp}((t+s)\mathbf{Z}) \cdot H.$$

If $t = 0$ the element $\alpha_g s$ is determined by

$$\widetilde{\exp}(u(\mathbf{Y} - \mathbf{Z})) \cdot \widetilde{\exp}(v\mathbf{X}) \cdot \widetilde{\exp}(s\mathbf{Z}) \cdot H = \widetilde{\exp}(\alpha_g s \mathbf{Z}) \cdot H.$$

Since there exists an automorphism $\beta : \mathfrak{sl}_2(\mathbb{R}) \rightarrow \mathfrak{sl}_2(\mathbb{R})$ mapping the basis $(\mathbf{X}, \mathbf{Y}, \mathbf{Z})$ onto the canonical basis $(\mathbf{H}, \mathbf{T}, \mathbf{U})$ of $\mathfrak{sl}_2(\mathbb{R})$ this equation is equivalent to

$$\widetilde{\exp}(u(\mathbf{T} - \mathbf{U})) \cdot \widetilde{\exp}(v\mathbf{H}) \cdot \widetilde{\exp}(s\mathbf{U}) \cdot \bar{H} = \widetilde{\exp}(\alpha_g s \mathbf{U}) \cdot \bar{H},$$

where $\bar{H} = \{\widetilde{\exp}(s(\mathbf{T} - \mathbf{U})) \cdot \widetilde{\exp}(t\mathbf{H}); s, t \in \mathbb{R}\} \subset G$. Applying the covering homomorphism $p : \widetilde{PSL}_2(\mathbb{R}) \rightarrow SL_2(\mathbb{R})$ we obtain

$$\begin{pmatrix} e^v & 0 \\ 2ue^v & e^{-v} \end{pmatrix} \begin{pmatrix} \cos s & \sin s \\ -\sin s & \cos s \end{pmatrix} p(\bar{H}) = \begin{pmatrix} \cos \alpha_g s & \sin \alpha_g s \\ -\sin \alpha_g s & \cos \alpha_g s \end{pmatrix} p(\bar{H})$$

or equivalently

$$\begin{pmatrix} e^v & 0 \\ 2ue^v & e^{-v} \end{pmatrix} \begin{pmatrix} \cos s & \sin s \\ -\sin s & \cos s \end{pmatrix} = \begin{pmatrix} \cos \alpha_g s & \sin \alpha_g s \\ -\sin \alpha_g s & \cos \alpha_g s \end{pmatrix} \begin{pmatrix} e^x & 0 \\ 2ye^x & e^{-x} \end{pmatrix}$$

for some $x, y \in \mathbb{R}$. Comparing the second columns of the product matrices we obtain

$$\cot \alpha_g s = \frac{2ue^v \sin s + e^{-v} \cos s}{e^v \sin s} = 2u + e^{-2v} \cot s.$$

Since the map $\alpha_g : \mathbb{R} \rightarrow \mathbb{R}$ is a bijection and hence it is monotone. The shape of the last expression shows that α_g is monotone increasing. The assertion follows from this and from $\alpha_g 0 = 0$. $\square$

3. The left multiplication curve

We consider a differentiable left loop defined on a one-manifold and assume that the group topologically generated by the left translations is isomorphic to $G = \widetilde{PSL}_2(\mathbb{R})$. Since G acts transitively, the left loop manifold may be identified with the factor space G/H , where the 2-dimensional subgroup H of G is the identity of the left loop $(G/H, \circ, H)$. Let $\pi : G \rightarrow G/H$ be the canonical projection map and let ϕ be the parametrization $\mathbb{R} \rightarrow G/H$ introduced in equation (2). We

define a differentiable left loop $(\mathbb{R}, \circ, 0)$ on the real line $\mathbb{R}$ assuming that the map $\phi : (G/H, \circ, H) \rightarrow (\mathbb{R}, \circ, 0)$ is an isomorphism. The left translations $\lambda_x, x \in \mathbb{R}$, of $(\mathbb{R}, \circ, 0)$ have the shape $\lambda_x = \alpha_{\sigma_x}$ where $\sigma_x \in G$ is the left translation on G/H corresponding to the element $\phi(x) \in G/H$. Hence σ_x is contained in $\phi(x) \in G/H$ for any $x \in \mathbb{R}$ and has the same action on G/H as the mapping given by $\phi\lambda_x\phi^{-1}$.

Definition 2. A differentiable curve $\sigma : x \mapsto \sigma_x : \mathbb{R} \rightarrow G$ is called a left multiplication curve on G if σ_x is contained in $\phi(x) \in G/H$ for any $x \in \mathbb{R}$. The map $(x, y) \mapsto x \cdot y = \phi^{-1}\sigma_x\phi(y) : \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$ is called the multiplication associated with the left multiplication curve σ with respect to the Iwasawa parametrization (2).

The multiplication $(x, y) \mapsto x \cdot y$ on $\mathbb{R}$, associated with a left multiplication curve σ determines a differentiable left loop $(\mathbb{R}, \circ, 0)$ which has $0 \in \mathbb{R}$ as unit element. One can see immediately, that the left multiplication maps of this left loop topologically generate a group which is isomorphic to G .

Now, let $(G/H, \circ, H)$ be a differentiable left loop having $G = \widetilde{PSL}_2(\mathbb{R})$ as the group topologically generated by the left translations. The mapping $x \mapsto \sigma_x : \mathbb{R} \rightarrow G$ satisfying $\lambda_x = \alpha_{\sigma_x}$ determines a left multiplication curve on G , and the multiplication associated with σ is isomorphic to the multiplication of the left loop $(G/H, \circ, H)$. According to $G = K \cdot N \cdot D$ we have the decomposition

$$\sigma_x = \widetilde{\exp}(x\mathbf{Z}) \cdot \widetilde{\exp}(\nu(x)(\mathbf{Y} - \mathbf{Z})) \cdot \widetilde{\exp}(\delta(x)\mathbf{X}),$$

where $\nu(x)$ and $\delta(x)$ are real differentiable functions having the initial value $\nu(0) = \delta(0) = 0$. The left translations of the associated multiplication $(x, y) \mapsto x \cdot y$ have the shape $\lambda_x = \alpha_{\sigma_x}$. If the left multiplication curve is not a one-parameter subgroup of G then the group topologically generated by the left translations is isomorphic to $G = \widetilde{PSL}_2(\mathbb{R})$.

We obtain from Proposition 1 the following

Proposition 3. *Let $\sigma : x \mapsto \sigma_x : \mathbb{R} \rightarrow \widetilde{PSL}_2(\mathbb{R})$ be a left multiplication curve of the shape $\sigma_x = \widetilde{\exp}(x\mathbf{Z}) \cdot \widetilde{\exp}(\nu(x)(\mathbf{Y} - \mathbf{Z})) \cdot \widetilde{\exp}(\delta(x)\mathbf{X})$, where $\nu(x)$ and $\delta(x)$ are differentiable functions having the initial value $\nu(0) = \delta(0) = 0$. The multiplication $(x, y) \mapsto x \cdot y$ associated with the left multiplication curve σ with respect to the Iwasawa parametrization (2) can be expressed by*

$$x \cdot y = \begin{cases} x + \operatorname{arccot}(2\nu(x) + e^{-2\delta(x)} \cot y) + i\pi, & \text{if } i\pi < y < (i+1)\pi, \\ x + i\pi, & \text{if } y = i\pi, \end{cases}$$

for any $i \in \mathbb{Z}$.

Now, we want characterize the loop multiplications among the differentiable left loop multiplications described in the previous proposition. Clearly, if $(\mathbb{R}, \cdot, 0)$ is a loop, then for any $y \in \mathbb{R}$ the right multiplication map $\rho_y : \mathbb{R} \rightarrow \mathbb{R}$ is a diffeomorphism. It follows that the map $\rho_y : \mathbb{R} \rightarrow \mathbb{R}$ and its inverse $\rho_y^{-1} : \mathbb{R} \rightarrow \mathbb{R}$ are monotone differentiable functions and hence the function $m(x, y) = x \cdot y$ satisfies $\frac{\partial}{\partial x} m(x, y) \neq 0$ for any $x, y \in \mathbb{R}$. Since $m(x, 0) = x$ one has $\frac{\partial}{\partial x} m(x, 0) = 1$ from which follows $\frac{\partial}{\partial x} m(x, y) > 0$ for each $x, y \in \mathbb{R}$. Now we show that this condition characterize the loop multiplications.

Proposition 4. *The multiplication $x \cdot y = \phi^{-1} \sigma_x \phi(y)$ associated with the left multiplication curve σ with respect to the Iwasawa parametrization (2) determines a differentiable loop $(\mathbb{R}, \cdot, 0)$ if and only if the function $m(x, y) = x \cdot y$ satisfies $\frac{\partial}{\partial x} m(x, y) > 0$.*

PROOF. If the inequality $\frac{\partial}{\partial x} m(x, y) > 0$ is satisfied for any $x, y \in \mathbb{R}$ then the maps $\rho_y : \mathbb{R} \rightarrow \rho_y(\mathbb{R})$ are monotone increasing diffeomorphisms of $\mathbb{R}$ onto the connected subset $\rho_y(\mathbb{R}) \subset \mathbb{R}$. We have to prove that the maps $\rho_y : \mathbb{R} \rightarrow \mathbb{R}$ are surjective. Let $y \in \mathbb{R}$ be fixed. We assume that there exists a real number a such that $\sup_{x \in \mathbb{R}} (x \cdot y) = \lim_{x \rightarrow \infty} \rho_y(x) = \lim_{x \rightarrow \infty} \lambda_x(y) = a$. We have up to a conjugation in $\widetilde{PSL}_2(\mathbb{R})$

$$\widetilde{\exp}(x\mathbf{U}) \cdot \widetilde{\exp}(\nu(x)(\mathbf{T} - \mathbf{U})) \cdot \widetilde{\exp}(\delta(x)\mathbf{H}) \cdot \widetilde{\exp}(y\mathbf{U}) \cdot \bar{H} = \widetilde{\exp}((x \cdot y)\mathbf{U}) \cdot \bar{H}.$$

Applying the continuous covering map $p : \widetilde{PSL}_2(\mathbb{R}) \rightarrow SL_2(\mathbb{R})$ we obtain

$$\begin{aligned} \begin{pmatrix} \cos x & \sin x \\ -\sin x & \cos x \end{pmatrix} \begin{pmatrix} e^{\delta(x)} & 0 \\ 2\nu(x)e^{\delta(x)} & e^{-\delta(x)} \end{pmatrix} \begin{pmatrix} \cos y & \sin y \\ -\sin y & \cos y \end{pmatrix} p(H) \\ = \begin{pmatrix} \cos(x \cdot y) & \sin(x \cdot y) \\ -\sin(x \cdot y) & \cos(x \cdot y) \end{pmatrix} p(H). \end{aligned}$$

Hence we have the decomposition

$$\begin{aligned} \begin{pmatrix} \cos(x - x \cdot y) & \sin(x - x \cdot y) \\ -\sin(x - x \cdot y) & \cos(x - x \cdot y) \end{pmatrix} \begin{pmatrix} e^{\delta(x)} & 0 \\ 2\nu(x)e^{\delta(x)} & e^{-\delta(x)} \end{pmatrix} \\ = \begin{pmatrix} e^{d(x, y)} & 0 \\ 2n(x, y)e^{d(x, y)} & e^{-d(x, y)} \end{pmatrix} \begin{pmatrix} \cos y & -\sin y \\ \sin y & \cos y \end{pmatrix}, \end{aligned}$$

where $d(x, y)$ and $n(x, y)$ are suitable continuous real functions. If $\sin y = 0$ then this equation implies $x = x \cdot y + 2k\pi$ for any $x \in \mathbb{R}$ and hence $y = 0$, in which case

trivially $\rho_0(\mathbb{R}) = \mathbb{R}$. Assuming $\sin y \neq 0$ and comparing the first rows of these product matrices we obtain

$$\begin{aligned} & \left(e^{\delta(x)} \cos(x - x \cdot y) + 2\nu(x) e^{\delta(x)} \sin(x - x \cdot y), e^{-\delta(x)} \sin(x - x \cdot y) \right) \\ &= \left(e^{d(x,y)} \cos y, -e^{d(x,y)} \sin y \right). \end{aligned}$$

From this we have $e^{2\delta(x)}(\cot(x - x \cdot y) + 2\nu(x)) = -\cot y$. Since the function $x \mapsto x - x \cdot y$ is continuous, its range is a connected subset of $\mathbb{R}$ such that $\sup_{x \in \mathbb{R}}(x - x \cdot y) = \infty$ because of $\lim_{x \rightarrow \infty} x \cdot y = a$. Hence there exists a real value $x_0 \in \mathbb{R}$ satisfying $\sin(x_0 - x_0 \cdot y) = 0$ for which the equation

$$e^{2\delta(x_0)}(\cot(x_0 - x_0 \cdot y) + 2\nu(x_0)) = -\cot y$$

gives a contradiction. $\square$

We obtain the following characterization of the left multiplication curves of loop multiplications:

Theorem 5. *Let $\sigma : x \mapsto \sigma_x : \mathbb{R} \rightarrow \widetilde{PSL}_2(\mathbb{R})$ be a left multiplication curve of the shape $\sigma_x = \widetilde{\exp}(x\mathbf{Z}) \cdot \widetilde{\exp}(\nu(x)(\mathbf{Y} - \mathbf{Z})) \cdot \widetilde{\exp}(\delta(x)\mathbf{X})$, where $\nu(x)$ and $\delta(x)$ are differentiable functions having the initial value $\nu(0) = \delta(0) = 0$. The multiplication $(x, y) \mapsto x \cdot y$ associated with the left multiplication curve σ with respect to the Iwasawa parametrization (2) determines a differentiable loop on $\mathbb{R}$ if and only if the functions $\delta(t)$ and $\nu(t)$ satisfy the differential inequality*

$$g(\dot{\sigma}_x, \dot{\sigma}_x) = \dot{\delta}(t)^2 + 2\dot{\nu}(t) + 4\nu(t)\dot{\delta}(t) - 1 < 0,$$

where $\dot{\delta} = \frac{d\delta}{dx}$, $\dot{\nu} = \frac{d\nu}{dx}$ and $\dot{\sigma}_x = \frac{d\sigma_x}{dx}$.

PROOF. The derivation of the multiplication given in Theorem 3 gives that

$$\frac{\partial}{\partial x} m(x, y) = 1 - \frac{2\dot{\nu}(x) - 2e^{-2\delta(x)}\dot{\delta}(x)\cot y}{1 + (2\nu(x) + e^{-2\delta(x)}\cot y)^2} > 0$$

if and only if $1 + (2\nu(x) + e^{-2\delta(x)}\cot y)^2 - 2\dot{\nu}(x) + 2e^{-2\delta(x)}\dot{\delta}(x)\cot y > 0$. Equivalently,

$$(1 + 4\nu^2(x) - 2\dot{\nu}(x)) + 2e^{-2\delta(x)}(2\nu(x) + \dot{\delta}(x))\cot y + e^{-4\delta^2(x)}\cot^2 y > 0$$

for all $y \in \mathbb{R}$, which gives $(2\nu(x) + \dot{\delta}(x))^2 - (1 + 4\nu^2(x) - 2\dot{\nu}(x)) < 0$ or

$$\dot{\delta}^2(x) + 2\dot{\nu}(x) + 4\nu(x)\dot{\delta}(x) - 1 < 0. \quad (3)$$

For the computation of the pseudo-Riemannian value $g(\dot{\sigma}_x, \dot{\sigma}_x)$ we apply the covering map $p : \widetilde{PSL}_2(\mathbb{R}) \rightarrow SL_2(\mathbb{R})$ to σ_x and derivate it:

$$\begin{aligned} \frac{d}{dx}p(\sigma_x) &= \begin{pmatrix} -\sin x & \cos x \\ -\cos x & -\sin x \end{pmatrix} \begin{pmatrix} e^{\delta(x)} & 0 \\ 2\nu(x)e^{\delta(x)} & e^{-\delta(x)} \end{pmatrix} \\ &\quad + \begin{pmatrix} \cos x & \sin x \\ -\sin x & \cos x \end{pmatrix} \begin{pmatrix} \dot{\delta}(x)e^{\delta(x)} & 0 \\ (2\nu(x) + 2\nu(x)\dot{\delta}(x))e^{\delta(x)} & -\dot{\delta}(x)e^{-\delta(x)} \end{pmatrix} \\ &= \begin{pmatrix} \cos x & \sin x \\ -\sin x & \cos x \end{pmatrix} \left\{ \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 2\nu(x) & 1 \end{pmatrix} \right. \\ &\quad \left. + \begin{pmatrix} \dot{\delta}(x) & 0 \\ 2\nu(x) + 2\nu(x)\dot{\delta}(x) & -\dot{\delta}(x) \end{pmatrix} \right\} \begin{pmatrix} e^{\delta(x)} & 0 \\ 0 & e^{-\delta(x)} \end{pmatrix}. \end{aligned}$$

Since $g(\dot{\sigma}_x, \dot{\sigma}_x) = \gamma(\sigma_x^{-1}\dot{\sigma}_x) = -\det(p_*(\sigma_x^{-1}\dot{\sigma}_x)) = -\det(p_*(\dot{\sigma}_x))$ we can compute

$$\begin{aligned} g(\dot{\sigma}_x, \dot{\sigma}_x) &= -\det \begin{pmatrix} 2\nu(x) + \dot{\delta}(x) & 1 \\ -1 + 2\nu(x) + 2\nu(x)\dot{\delta}(x) & -\dot{\delta}(x) \end{pmatrix} \\ &= \dot{\delta}^2(x) + 2\dot{\nu}(x) + 4\nu(x)\dot{\delta}(x) - 1. \end{aligned}$$

Hence the condition (3) is equivalent to $g(\dot{\sigma}_x, \dot{\sigma}_x) < 0$ which proves the assertion. $\square$

Corollary 6. *The multiplication $(x, y) \mapsto x \cdot y$ associated with the left multiplication curve σ is a loop multiplication if and only if the left translation curve is time-like.*

References

- [1] L. E. J. BROUWER, The Theorie der endlichen kontinuerlichen Gruppen, unabhängig von den Axiomen von Lie, *Math. Ann.* **67** (1909), 246–267.
- [2] S. HELGASON, Differential Geometry and Symmetric Spaces, *Academic Press, New York – London*, 1962.
- [3] P. T. NAGY and K. STRAMBACH, Loops in Group Theory and Lie Theory, Expositions in Mathematics 35, *Walter de Gruyter, Berlin – New York*, 2002.

- [4] P. T. NAGY and K. STRAMBACH, Coverings of topological loops, *Journal of Math. Sci.* **137** (2006), 5098–5116.

PÉTER T. NAGY
INSTITUTE OF MATHEMATICS
UNIVERSITY OF DEBRECEN
4010 DEBRECEN, P.O. BOX 12
HUNGARY

E-mail: nagypeti@math.klte.hu

IZABELLA STUHL
INSTITUTE OF MATHEMATICS
UNIVERSITY OF DEBRECEN
4010 DEBRECEN, P.O. BOX 12
HUNGARY

E-mail: stuhl@mi.uni-erlangen.de

(Received July 13, 2005)