

Prime ideals and complex ring homomorphisms on a commutative algebra

By TAKESHI MIURA (Yonezawa), SIN-EI TAKAHASI (Yonezawa)
and NORIO NIWA (Neyagawa)

Abstract. We give a characterization of prime ideals $\mathcal{P}$ of a commutative complex algebra $\mathcal{A}$ in order that $\mathcal{P}$ be the kernel of some complex ring homomorphism on $\mathcal{A}$. If, in addition, $\mathcal{A}$ is a uniform algebra on an infinite compact metric space, then we show that there are exactly 2^c complex ring homomorphisms on $\mathcal{A}$, whose kernels are non-maximal prime ideals. Moreover, it turns out that ring homomorphisms on a commutative Banach algebra are deeply connected with the existence of discontinuous homomorphisms.

1. Introduction and the statement of results

Let $\mathcal{A}$ and $\mathcal{B}$ be algebras over the complex number field $\mathbb{C}$. We say that a mapping $\rho : \mathcal{A} \rightarrow \mathcal{B}$ is a ring homomorphism, provided that

$$\rho(f + g) = \rho(f) + \rho(g)$$

$$\rho(fg) = \rho(f)\rho(g)$$

for every $f, g \in \mathcal{A}$. Moreover if ρ is homogeneous, that is $\rho(\lambda f) = \lambda\rho(f)$ for every $f \in \mathcal{A}$ and $\lambda \in \mathbb{C}$, then ρ is an ordinary homomorphism. It is obvious that if $\rho : \mathcal{A} \rightarrow \mathcal{B}$ is a ring homomorphism, then $\rho(rf) = r\rho(f)$ for every rational number r and $f \in \mathcal{A}$. If, in addition, ρ is assumed to be continuous, then we see that ρ is real linear, that is, $\rho(tf) = t\rho(f)$ for every real number t and $f \in \mathcal{A}$. So, we consider ring homomorphisms which need not be continuous. The study of ring homomorphisms between two Banach algebras has a long history. In 1944,

Mathematics Subject Classification: 46J10.

Key words and phrases: prime ideals, ring homomorphisms.

ARNOLD [1] proved that a ring isomorphism between the two Banach algebras of all bounded linear operators on two infinite dimensional Banach spaces is linear or conjugate linear. KAPLANSKY [6] generalized this result as follows: If ρ is a ring isomorphism from one semisimple Banach algebra A onto another, then A is the direct sum of closed ideals A_1 , A_2 and A_3 such that $\rho|_{A_1}$ is linear, $\rho|_{A_2}$ is conjugate linear and that A_3 is finite dimensional: The finite dimensional part is not trivial in general. In fact, KESTELMAN [7] proved that there exists a ring homomorphism $\rho : \mathbb{C} \rightarrow \mathbb{C}$ such that ρ is neither linear nor conjugate linear. Moreover, CHARNOW [2, Theorem 3] proved that there exist $2^{\aleph_k}$ ring automorphisms for every algebraically closed field k . Here and after, $\aleph_S$ denotes the cardinal number of a set S . In particular, there are $2^{\aleph_{\mathbb{C}}}$ ring automorphisms on $\mathbb{C}$. MOLNÁR [10, Theorem 1] essentially gave a representation of a ring homomorphism between two commutative C^* -algebras.

Suppose $\rho : A \rightarrow B$ is a ring homomorphism between two commutative Banach algebras A and B with the maximal ideal spaces M_A and M_B , respectively. When studying such mappings, a natural approach would be to consider ring homomorphisms $\varphi \circ \rho : A \rightarrow \mathbb{C}$ for every $\varphi \in M_B$, and patch them by a continuous mapping from a suitable subset of M_B into M_A . Indeed, some representations of ring homomorphisms, with additional conditions, are proved in this way (cf. [5, Theorem 2.3], [9, Theorem 2.6], [11, Theorem 5.1, 5.2]). Unfortunately this approach does not work in general because the kernel $\ker(\varphi \circ \rho)$ need not be a maximal ideal of A . On the other hand, Molnár [10, Theorem 2] essentially gave a representation of ring homomorphisms between two commutative C^* -algebras by another approach. Although we are concerned with *ring* homomorphism, the term ideal will mean an *algebra* ideal. Let $C(X)$ denote the commutative Banach algebra of all complex valued continuous functions on a compact Hausdorff space X . Suppose $\tau : \mathbb{C} \rightarrow \mathbb{C}$ is a ring homomorphism, and suppose $x_0 \in X$. ŠEMRL [11, Example 5.4] considered a complex ring homomorphism $\rho : C(X) \rightarrow \mathbb{C}$ of the form

$$\rho(f) = \tau(f(x_0)) \quad (f \in C(X)) \quad (*)$$

and gave the following example: If $\mathbb{N}$ is the set of all natural numbers and if K is the closure of $\{1/n : n \in \mathbb{N}\}$ with its usual topology, then there is a non-zero ring homomorphism $\phi : C(K) \rightarrow \mathbb{C}$ such that ϕ is not of the form (*). In particular, $\ker \phi$ is a non-maximal prime ideal of $C(K)$. The first author [9, Lemma 2.1] gave a characterization of a ring homomorphism ρ between two commutative Banach algebras in order that $\ker \rho$ be a maximal ideal (cf. [5, Lemma 2.2]).

In this note, we are concerned with complex ring homomorphisms ρ on a commutative complex algebra $\mathcal{A}$. If ρ is non-zero, then it is easy to see that

$\ker \rho$ is a prime ideal of $\mathcal{A}$. Recall that if $\mathcal{A}$ is a unital commutative Banach algebra, then there is a one-to-one correspondence between non-zero complex homomorphisms on $\mathcal{A}$ and maximal ideals of $\mathcal{A}$. With this in mind, one might expect that there is also a correspondence between complex ring homomorphisms and prime ideals of a complex commutative algebra $\mathcal{A}$. In this note, we give a characterization of prime ideals that can be represented as the kernels of some complex ring homomorphisms. Before we state our main result, we need some terminology. If $\mathcal{A}$ is unital, then we define $\mathcal{A}_e \stackrel{\text{def}}{=} \mathcal{A}$; otherwise, $\mathcal{A}_e$ denotes the commutative complex algebra obtained by adjunction of a unit element e to $\mathcal{A}$. As usual, we may identify $f \in \mathcal{A}$ with $(f, 0) \in \mathcal{A}_e$. Now, we are ready to state our main result.

Theorem 1.1. *Suppose $\mathcal{A}$ is a commutative complex algebra and $\mathcal{P}$ is a prime ideal of $\mathcal{A}$. Put $\mathfrak{c} = \sharp \mathbb{C}$. Then each of the following four properties implies the other three:*

- (a) *There exists a non-zero ring homomorphism $\rho : \mathcal{A} \rightarrow \mathbb{C}$ such that $\ker \rho = \mathcal{P}$.*
- (b) *The quotient algebra $\mathcal{A}/\mathcal{P}$ has the cardinal number $\mathfrak{c}$.*
- (c) *There exists a prime ideal $\tilde{\mathcal{P}}$ of $\mathcal{A}_e$ such that $\mathcal{P} = \tilde{\mathcal{P}} \cap \mathcal{A}$ and that $\mathcal{A}_e/\tilde{\mathcal{P}}$ has the cardinal number $\mathfrak{c}$.*
- (d) *There exists a non-zero ring homomorphism $\tilde{\rho} : \mathcal{A}_e \rightarrow \mathbb{C}$ such that $\mathcal{A} \cap \ker \tilde{\rho} = \mathcal{P}$.*

Let K be the closure of $\{1/n : n \in \mathbb{N}\}$ with its usual topology. As stated above, Šemrl gave a complex ring homomorphism on $C(K)$, whose kernel is a non-maximal prime ideal. In the following corollary we see that there exist $2^{\mathfrak{c}}$ such mappings. Moreover, the following is true.

Corollary 1.2. *If A is a uniform algebra on an infinite compact metric space, then there are exactly $2^{\mathfrak{c}}$ complex ring homomorphisms on A , whose kernels are non-maximal prime ideals.*

2. A proof of results

Recall that an ideal $\mathcal{P}$ of a commutative algebra is prime if $\mathcal{P}$ is proper and $fg \notin \mathcal{P}$ whenever $f \notin \mathcal{P}$ and $g \notin \mathcal{P}$.

It is advisable to note that the quotient field of an integral domain R is well defined even if R has no unit: If $a \in R \setminus \{0\}$, then the “fraction” a/a is a unit, and we may identify $b \in R$ with ab/a .

Lemma 2.1. *Suppose $\mathcal{A}$ is a commutative complex algebra and $\rho : \mathcal{A} \rightarrow \mathbb{C}$ is a non-zero ring homomorphism. Then*

- (a) *the kernel $\ker \rho$ is a prime ideal of $\mathcal{A}$, and*
- (b) *ρ is of the form $\rho = \tau \circ \pi$, where τ is a non-zero field homomorphism on the quotient field $\mathcal{F}$ of $\mathcal{A}/\ker \rho$ into $\mathbb{C}$, and $\pi : \mathcal{A} \rightarrow \mathcal{A}/\ker \rho$ is the quotient mapping.*

PROOF. Choose $a \in \mathcal{A}$ such that $\rho(a) \neq 0$: This is possible since ρ is assumed to be non-zero. (a) Pick $f \in \ker \rho$ and $\lambda \in \mathbb{C}$ arbitrarily. It follows that

$$\rho(\lambda f) \rho(a) = \rho(f) \rho(\lambda a) = 0,$$

and hence $\lambda f \in \ker \rho$. We thus obtain that $\ker \rho$ is an (algebra) ideal. Now it is obvious that $\ker \rho$ is a prime ideal.

(b) Let $\mathcal{F}$ be the quotient field of $\mathcal{A}/\ker \rho$. $\mathcal{F}$ is well defined since $\mathcal{A}/\ker \rho$ is an integral domain by (a). We define the mapping $\tau : \mathcal{F} \rightarrow \mathbb{C}$ by

$$\tau(\pi(f)/\pi(g)) = \frac{\rho(f)}{\rho(g)} \quad (\pi(f)/\pi(g) \in \mathcal{F}).$$

A simple calculation shows that τ is a well defined non-zero field homomorphism. As usual we may identify $\pi(f) \in \mathcal{A}/\ker \rho$ with $\pi(fa)/\pi(a) \in \mathcal{F}$. We get

$$\tau(\pi(f)) = \tau(\pi(fa)/\pi(a)) = \frac{\rho(fa)}{\rho(a)} = \rho(f) \quad (f \in \mathcal{A}),$$

and hence $\rho = \tau \circ \pi$. □

Lemma 2.2. *Suppose $\mathcal{A}$ is a commutative complex algebra and $\mathcal{P}$ is a prime ideal of $\mathcal{A}$. Then*

- (a) $\mathfrak{c} = \sharp \mathbb{C} \leq \sharp(\mathcal{A}/\mathcal{P})$, and
- (b) *if $a \in \mathcal{A} \setminus \mathcal{P}$, the set $\mathcal{P}_e \stackrel{\text{def}}{=} \{(f, \lambda) \in \mathcal{A}_e : fa + \lambda a \in \mathcal{P}\}$ is a prime ideal of $\mathcal{A}_e$ such that $\mathcal{P} = \mathcal{P}_e \cap \mathcal{A}$.*

PROOF. Pick $a \in \mathcal{A} \setminus \mathcal{P}$ arbitrarily.

(a) Let $\pi : \mathcal{A} \rightarrow \mathcal{A}/\mathcal{P}$ be the quotient mapping. Since $a \notin \mathcal{P}$ and since $\mathcal{P}$ is an ideal, $\pi(\lambda a) = \pi(\mu a)$ implies $\lambda = \mu$ for $\lambda, \mu \in \mathbb{C}$. This shows that the mapping $\lambda \mapsto \pi(\lambda a)$ is an injection, so that

$$\mathfrak{c} = \sharp \mathbb{C} = \sharp\{\pi(\lambda a) : \lambda \in \mathbb{C}\} \leq \sharp(\mathcal{A}/\mathcal{P}).$$

(b) It is easy to see that $\mathcal{P}_e$ is a proper ideal of $\mathcal{A}_e$ such that $\mathcal{P} = \mathcal{P}_e \cap \mathcal{A}$. To show that $\mathcal{P}_e$ is prime, suppose $(f_1, \lambda_1)(f_2, \lambda_2) \in \mathcal{P}_e$. By definition, this implies that $(f_1 f_2 + \lambda_2 f_1 + \lambda_1 f_2)a + (\lambda_1 \lambda_2)a \in \mathcal{P}$, and so we obtain $(f_1 a + \lambda_1 a)(f_2 a + \lambda_2 a) \in \mathcal{P}$. Since $\mathcal{P}$ is a prime ideal, $(f_1 a + \lambda_1 a)$ or $(f_2 a + \lambda_2 a)$ belongs to $\mathcal{P}$. This implies $(f_1, \lambda_1) \in \mathcal{P}_e$ or $(f_2, \lambda_2) \in \mathcal{P}_e$, and hence $\mathcal{P}_e$ is prime. $\square$

Let $\mathcal{K}$ be a transcendental extension field of a commutative field k , and S a subset of $\mathcal{K}$. We recall that S is said to be algebraically independent over k , if the set of all finite products of elements of S is linearly independent over k . A subset T of $\mathcal{K}$ is called a *transcendence base* of $\mathcal{K}$ over k , if T is algebraically independent over k which is maximal with respect to the inclusion ordering. The existence of a transcendence base of $\mathcal{K}$ over k is well known (cf. [8, Theorem 1.1 of Chapter X]). The maximality of T shows that $\mathcal{K}$ is algebraic over $k(T)$, the field generated by T over k .

Lemma 2.3. *Let $\mathbb{Q}$ be the rational number field and k a transcendental extension field of $\mathbb{Q}$ such that $\sharp k = \mathfrak{c}$. If T is a transcendence base of k over $\mathbb{Q}$, then $\sharp T = \mathfrak{c}$.*

PROOF. Suppose T is a transcendence base of k over $\mathbb{Q}$. Let $\mathbb{Q}(T)$ be the field generated by T over $\mathbb{Q}$. Since $T \subset \mathbb{Q}(T) \subset k$, we obtain $\sharp T \leq \sharp \{\mathbb{Q}(T)\} \leq \mathfrak{c}$. So, we show that $\mathfrak{c} \leq \sharp T$. Since k is algebraic over $\mathbb{Q}(T)$, each element of k is a zero point of some function in $\wp$, the set of all monic polynomials over $\mathbb{Q}(T)$. Note that for each monic polynomial, its zero points in k is at most finite. Put $\mathfrak{a} = \sharp \mathbb{Q}$, then we have

$$\mathfrak{c} = \sharp k \leq (\sharp \wp) \times \mathfrak{a} \leq (\sharp \{\mathbb{Q}(T)\} \times \mathfrak{a}) \times \mathfrak{a} = \sharp \{\mathbb{Q}(T)\},$$

and hence $\mathfrak{c} \leq \sharp \{\mathbb{Q}(T)\}$. We thus obtain

$$\mathfrak{c} = \sharp \{\mathbb{Q}(T)\} \leq \mathfrak{a} \times \sharp T. \quad (**)$$

If T were finite, then we would have $\sharp \{\mathbb{Q}(T)\} = \mathfrak{a}$, in contradiction to $\sharp \{\mathbb{Q}(T)\} = \mathfrak{c}$. It follows that $\sharp T \geq \mathfrak{a}$, and so $\mathfrak{a} \times \sharp T = \sharp T$. By $(**)$ we get $\mathfrak{c} \leq \sharp T$, proving $\sharp T = \mathfrak{c}$. $\square$

PROOF OF THEOREM 1.1. Let $\pi : \mathcal{A} \rightarrow \mathcal{A}/\mathcal{P}$ be the quotient mapping and fix $a \in \mathcal{A} \setminus \mathcal{P}$.

(a) $\Rightarrow$ (b) By (a) of Lemma 2.2, we obtain $\mathfrak{c} \leq \sharp(\mathcal{A}/\mathcal{P})$. To prove the opposite inequality, let $\mathcal{F}$ be the quotient field of $\mathcal{A}/\mathcal{P}$. By (b) of Lemma 2.1, we can write $\rho = \tau \circ \pi$, where $\tau : \mathcal{F} \rightarrow \mathbb{C}$ is a field homomorphism, and hence injective. It

follows that $\sharp\mathcal{F} \leq \sharp\mathbb{C} = \mathfrak{c}$. If we regard $\mathcal{A}/\mathcal{P}$ as a subset of $\mathcal{F}$, it follows that $\sharp(\mathcal{A}/\mathcal{P}) \leq \sharp\mathcal{F} \leq \mathfrak{c}$, proving $\sharp(\mathcal{A}/\mathcal{P}) = \mathfrak{c}$.

(b) $\Rightarrow$ (c) Let $\mathcal{P}_e$ be the prime ideal of $\mathcal{A}_e$ as in (b) of Lemma 2.2. Let $\tilde{\pi} : \mathcal{A}_e \rightarrow \mathcal{A}_e/\mathcal{P}_e$ be the quotient mapping. Identification of f and $(f, 0)$ shows that $\pi(f) = \pi(g)$ if and only if $\tilde{\pi}(f, 0) = \tilde{\pi}(g, 0)$ for $f, g \in \mathcal{A}$, and so $\mathfrak{c} = \sharp(\mathcal{A}/\mathcal{P}) \leq \sharp(\mathcal{A}_e/\mathcal{P}_e)$. To show the opposite inequality, we define the mapping $\psi : \mathcal{A}_e/\mathcal{P}_e \rightarrow \mathcal{A}/\mathcal{P}$ by

$$\psi(\tilde{\pi}(f, \lambda)) = \pi(fa + \lambda a) \quad (\tilde{\pi}(f, \lambda) \in \mathcal{A}_e/\mathcal{P}_e).$$

A simple calculation shows that ψ is a well defined injection. Hence $\sharp(\mathcal{A}_e/\mathcal{P}_e) \leq \sharp(\mathcal{A}/\mathcal{P}) = \mathfrak{c}$, proving $\sharp(\mathcal{A}_e/\mathcal{P}_e) = \mathfrak{c}$.

(c) $\Rightarrow$ (d) Let $\tilde{\mathcal{F}}$ be the quotient field of $\mathcal{A}_e/\tilde{\mathcal{P}}$. Then

$$\mathfrak{c} = \sharp(\mathcal{A}_e/\tilde{\mathcal{P}}) \leq \sharp\tilde{\mathcal{F}} \leq \sharp(\mathcal{A}_e/\tilde{\mathcal{P}}) \times \sharp(\mathcal{A}_e/\tilde{\mathcal{P}}) = \mathfrak{c},$$

so that $\tilde{\mathcal{F}}$ also has the cardinal number $\mathfrak{c}$. Note that $\tilde{\mathcal{F}}$ is a transcendental extension of $\mathbb{Q}$ since $\tilde{\mathcal{F}}$ contains a unital algebra $\mathcal{A}_e/\tilde{\mathcal{P}} \supset \mathbb{C}$.

Let T and $\tilde{T}$ be transcendence bases of $\mathbb{C}$ and $\tilde{\mathcal{F}}$ over $\mathbb{Q}$, respectively. By Lemma 2.3, we see that $\sharp T = \mathfrak{c} = \sharp\tilde{T}$. Thus we can find a bijection $\theta : \tilde{T} \rightarrow T$. Since T is algebraically independent over $\mathbb{Q}$, the mapping θ is naturally extended to a field homomorphism $\tilde{\theta} : \mathbb{Q}(\tilde{T}) \rightarrow \mathbb{Q}(T)$ so that $\tilde{\theta}(r) = r$ for every $r \in \mathbb{Q}$. Since $\tilde{\mathcal{F}}$ is an algebraic extension of $\mathbb{Q}(\tilde{T})$ and since $\mathbb{C}$ is algebraically closed, $\tilde{\theta}$ can be extended to a field homomorphism on $\tilde{\mathcal{F}}$ into $\mathbb{C}$ (cf. [8, Theorem 2.8 of Chapter VII]), which is also denoted by $\tilde{\theta}$. Define $\tilde{\rho} = \tilde{\theta} \circ \tilde{\pi}$, where $\tilde{\pi} : \mathcal{A}_e \rightarrow \mathcal{A}_e/\tilde{\mathcal{P}}$ is the quotient mapping. Then $\tilde{\rho} : \mathcal{A}_e \rightarrow \mathbb{C}$ is a ring homomorphism whose kernel is equal to $\tilde{\mathcal{P}}$, proving $\mathcal{A} \cap \ker \tilde{\rho} = \mathcal{A} \cap \tilde{\mathcal{P}} = \mathcal{P}$.

(d) $\Rightarrow$ (a) Put $\rho = \tilde{\rho}|_{\mathcal{A}}$. Then $\rho : \mathcal{A} \rightarrow \mathbb{C}$ is a non-zero ring homomorphism such that $\ker \rho = \mathcal{P}$. $\square$

PROOF OF COROLLARY 1.2. Suppose A is a uniform algebra on an infinite compact metric space X . Let Δ be the set of all non-maximal prime ideals of A . It is well known [4, Corollary 1] that there exist exactly $2^{\mathfrak{c}}$ non-maximal prime ideals of A , and hence $\sharp\Delta = 2^{\mathfrak{c}}$. Since X is separable, $\sharp A = \mathfrak{c}$: For if X_0 is a countable dense subset of X , then the restriction map $f \mapsto f|_{X_0}$ ($f \in A$) is injective since each element of A is continuous, and hence $\mathfrak{c} \leq \sharp A \leq \mathfrak{a} \times \mathfrak{c} = \mathfrak{c}$. So, there exist exactly $2^{\mathfrak{c}}$ functions on A into $\mathbb{C}$, which need not be continuous nor ring homomorphic. This implies that there are at most $2^{\mathfrak{c}}$ complex ring homomorphisms on A .

Conversely pick $P \in \Delta$ arbitrarily. By (a) of Lemma 2.2, we have

$$\mathfrak{c} \leq \sharp(A/P) \leq \sharp A = \mathfrak{c},$$

and hence $\sharp(A/P) = \mathfrak{c}$. So, by Theorem 1.1, to each $P \in \Delta$ there corresponds a complex ring homomorphism $\rho_P : A \rightarrow \mathbb{C}$ such that $\ker \rho_P = P$. Suppose $\rho_{P_1} = \rho_{P_2}$ for $P_1, P_2 \in \Delta$. It follows from $\ker \rho_{P_j} = P_j$ for $j = 1, 2$ that $P_1 = P_2$, and hence the mapping $P \mapsto \rho_P$ is an injection. We conclude that

$$2^{\mathfrak{c}} = \sharp \Delta \leq \sharp \{\rho_P : P \in \Delta\},$$

and the proof is complete. $\square$

Remark. Let A be a commutative Banach algebra. It is well-known (cf. [3, Theorem 5.7.32]) that under the continuum hypothesis there is a discontinuous homomorphism on A into some Banach algebra whenever there is a non-maximal prime ideal P of A with $\sharp(A/P) = \mathfrak{c}$. It follows from Theorem 1.1 that under the continuum hypothesis a discontinuous homomorphisms on A exists whenever there is a non-zero ring homomorphism $\rho : A \rightarrow \mathbb{C}$ such that $\ker \rho$ is non-maximal.

Example 2.1. Let $\bar{\mathbb{D}}$ denote the closure of the open unit disk $\mathbb{D}$ in $\mathbb{C}$. The disk algebra $A(\bar{\mathbb{D}})$ is a typical example of uniform algebras. HATORI, ISHII with the first and second author ([5, Corollary 5.3]) proved that if $\rho : A(\bar{\mathbb{D}}) \rightarrow A(\bar{\mathbb{D}})$ is a ring homomorphism whose range contains a non-constant function, then ρ is linear or conjugate linear.

Here, let us consider complex ring homomorphisms on $A(\bar{\mathbb{D}})$, that is, the range contains only constant functions. It is well known that the set of all non-zero complex homomorphisms on $A(\bar{\mathbb{D}})$ can be identified with $\bar{\mathbb{D}}$. So, there are $\mathfrak{c}$ complex homomorphisms on $A(\bar{\mathbb{D}})$. On the other hand, by Corollary 1.2, we see that there are $2^{\mathfrak{c}}$ ring homomorphisms whose kernels are non-maximal prime ideals.

Finally, we give a pathological feature of complex ring homomorphisms (cf. [5, Corollary 5.2]).

Example 2.2. If $H(\Omega)$ is the algebra of all analytic functions on a region $\Omega \subset \mathbb{C}$, then, as we shall show, $H(\Omega)$ is a subring of $\mathbb{C}$. In particular, it will follow that every subalgebra $\mathcal{A}$ of $H(\Omega)$, which is with or without unit, is a subring of $\mathbb{C}$. In fact, the ideal (0) containing only zero is a prime ideal of $H(\Omega)$. Moreover $\sharp H(\Omega) = \mathfrak{c}$ since $\mathfrak{c} = \sharp \mathbb{C} \leq \sharp H(\Omega) \leq \sharp C(\Omega) = \mathfrak{c}$, and so by Theorem 1.1 there exists a non-zero complex ring homomorphism ρ on $H(\Omega)$ such that $\ker \rho = (0)$. Therefore, ρ is an injective complex ring homomorphism on $H(\Omega)$.

References

- [1] B. H. ARNOLD, Rings of operators on vector spaces, *Ann. of Math.* **45** (1944), 24–49.
- [2] A. CHARNOW, The automorphisms of an algebraically closed field, *Canad. Math. Bull.* **18** (1970), 95–97.
- [3] H. G. DALES, Banach Algebras and Automatic Continuity, London Mathematical Society Monographs 24, *Clarendon Press, Oxford*, 2000.
- [4] W. E. DIETRICH, JR., Prime ideals in uniform algebras, *Proc. Amer. Math. Soc.* **42** (1974), 171–174.
- [5] O. HATORI, T. ISHII, T. MIURA and S.-E. TAKAHASI, Characterizations and automatic linearity for ring homomorphisms on algebras of functions, *Contemp. Math.* **328** (2003), 201–215.
- [6] I. KAPLANSKY, Ring isomorphisms of Banach algebras, *Canad. J. Math.* **6** (1954), 374–381.
- [7] H. KESTELMAN, Automorphisms of the field of complex numbers, *Proc. London Math. Soc.* (2) **53** (1951), 1–12.
- [8] S. LANG, Algebra (2nd edn), *Addison-Wesley, California*, 1984.
- [9] T. MIURA, A representation of ring homomorphisms on commutative Banach algebras, *Sci. Math. Jpn.* **53** (2001), 515–523.
- [10] L. MOLNÁR, The range of a ring homomorphism from a commutative C^* -algebra, *Proc. Amer. Math. Soc.* **124** (1996), 1789–1794.
- [11] P. ŠEMRL, Non linear perturbations of homomorphisms on $C(X)$, *Quart. J. Math. Oxford Ser. (2)* **50** (1999), 87–109.

TAKESHI MIURA
 DEPARTMENT OF BASIC TECHNOLOGY
 APPLIED MATHEMATICS AND PHYSICS
 YAMAGATA UNIVERSITY
 YONEZAWA 992-8510
 JAPAN

E-mail: miura@yz.yamagata-u.ac.jp

SIN-EI TAKAHASI
 DEPARTMENT OF BASIC TECHNOLOGY
 APPLIED MATHEMATICS AND PHYSICS
 YAMAGATA UNIVERSITY
 YONEZAWA 992-8510
 JAPAN

E-mail: sin-ei@emperor.yz.yamagata-u.ac.jp

NORIO NIWA
 FACULTY OF ENGINEERING
 OSAKA ELECTRO-COMMUNICATION UNIVERSITY
 NEYAGAWA 572-8530
 JAPAN

(Received September 16, 2005; revised January 12, 2006)