

Title: Locally graded Bell groups

Author(s): Costantino Delizia, Primož Moravec and Chiara Nicotera

For any integer $n \neq 0, 1$, a group is said to be n -Bell if it satisfies the law $[x^n, y] = [x, y^n]$. In this paper we prove that every finitely generated locally graded n -Bell group embeds into the direct product of a finite n -Bell group and a torsion-free nilpotent group of class ≤ 2 . We prove that n -Bell groups which are not locally graded always have infinite simple sections of finite exponent. Additionally, we obtain similar results for varieties of n -Levi groups and n -abelian groups defined by the laws $[x^n, y] = [x, y]^n$ and $(xy)^n = x^n y^n$, respectively. We give characterizations of these groups in the locally graded case.

Address:

Costantino Delizia
Dipartimento di Matematica e Informatica
Università di Salerno
Via Ponte don Melillo
84084 - Fisciano (SA)
Italy
E-mail: cdelizia@unisa.it

Address:

Primož Moravec
Department of Mathematics
University of Ljubljana
Jadranska 19
1000 Ljubljana
Slovenia
E-mail: primoz.moravec@fmf.uni-lj.si

Address:

Chiara Nicotera
Dipartimento di Matematica e Informatica
Università di Salerno
Via Ponte don Melillo
84084 - Fisciano (SA)
Italy
E-mail: cnicotera@unisa.it