

On the Diophantine equation $X^2 - (2^{2m} + 1)Y^4 = -2^{2m}$

By BO HE (Neijiang), ALAIN TOGBÉ (Westville) and P. GARY WALSH (Ottawa)

*This paper is dedicated to Professor Paulo Ribenboim on the occasion
of his 80th birthday*

Abstract. Using a recent result of Akhtari on quartic Thue equations, it is shown that the quartic equation $X^2 - (2^{2m} + 1)Y^4 = -2^{2m}$ has at most 12 solutions in odd positive integers $X, Y > 1$.

1. Introduction

In [3], LJUNGGREN proved that the quartic equation

$$X^2 - 2Y^4 = -1$$

has only the positive integer solutions $(x, y) = (1, 1), (239, 13)$. Also, in [4], LJUNGGREN proved that the only positive integer solutions to

$$X^2 - 5Y^4 = -4$$

are $(X, Y) = (1, 1)$. Since the case $m = 1$ was already studied, we need to consider $m \geq 2$.

Mathematics Subject Classification: 11D25, 11D41, 11B39.

Key words and phrases: elliptic curve, linear recurrence, diophantine equation.

The first author is supported by the Natural Science Foundation of Education Department of Sichuan Province (No.2006C057). The second author is partially supported by Purdue University North Central. The third author gratefully acknowledges support from the Natural Sciences and Engineering Research Council of Canada.

The goal of the present paper is to consider the family of equations

$$X^2 - (2^{2m} + 1)Y^4 = -2^{2m}.$$

It will first shown that a positive integer solution to this equation in positive integers gives rise to an integer solution to a certain Thue equation, and then apply a recent result of AKHTARI [1] in order to deduce an upper bound for the number of integer solutions to $X^2 - (2^{2m} + 1)Y^4 = -2^{2m}$, which is independent of m . As the specific cases $m = 0$ and $m = 1$ have already been dealt with by Ljunggren, we state our main theorems only for $m \geq 2$.

Theorem 1.1. *For any integer $m \geq 2$, the Diophantine equation*

$$X^2 - (2^{2m} + 1)Y^4 = -2^{2m} \tag{1}$$

has at most 12 solutions in odd positive integers $X, Y > 1$.

This result is almost certainly not the best possible. An extensive computation has only found the integer solution $(X, Y) = (103, 5)$, with $m = 2$ and $X > 1$.

2. Proof of Theorem 1.1

All coprime integer solutions (x, y) to the quadratic equation

$$x^2 - (2^{2m} + 1)y^2 = -2^{2m}$$

are given by

$$x + y\sqrt{2^{2m} + 1} = \pm \left(\pm 1 + \sqrt{2^{2m} + 1} \right) \left(2^m + \sqrt{2^{2m} + 1} \right)^{2i}$$

for some $i \geq 0$. We refer to Lemma 2 of [2] for this fact.

For brevity, let $a = 2^{m-1}$, and let

$$\alpha = T + U\sqrt{1 + 4a^2} = 2^m + \sqrt{2^{2m} + 1}.$$

For $i \geq 0$, define sequences $\{T_i\}$ and $\{U_i\}$ by

$$\alpha^i = T_i + U_i\sqrt{1 + 4a^2}.$$

Therefore, a positive integer solution to $X^2 - (2^{2m} + 1)Y^4 = -2^{2m}$ is equivalent to a solution to

$$Y^2 = T_{2k} \pm U_{2k}$$

for some $k \geq 0$. By the well known identities $T_{2k} = T_k^2 + (1 + 4a^2)U_k^2$ and

$U_{2k} = 2T_k U_k$, this gives

$$Y^2 = (T_k \pm U_k)^2 + (2aU_k)^2,$$

and it is evident that the terms involved in this equality are pairwise coprime. Thus, there are coprime non-negative integers r and s , of opposite parity, for which

$$Y = r^2 + s^2, \quad T_k \pm U_k = r^2 - s^2, \quad 2aU_k = 2rs.$$

We will assume that r is even, as the argument for the other case is identical. Letting $R = r/a$, solving each of these expressions for T_k and U_k , substituting the result into $T_k^2 - (1 + 4a^2)U_k^2 = \pm 1$, and then simplifying leads to the equation

$$s^4 \pm 2s^3 R - 6a^2 R^2 s^2 \mp 2a^2 R^3 S + a^4 R^4 = \pm 1.$$

Now putting $x = \pm s$ and $y = R$ gives the Thue equation

$$x^4 - 2x^3 y - 6a^2 x^2 y^2 + 2a^2 x y^3 + a^4 y^4 = \pm 1.$$

There roots of the dehomogenized quartic polynomial

$$p_a(x) = x^4 - 2x^3 - 6a^2 x^2 + 2a^2 x + a^4, \quad (2)$$

are given explicitly by

$$\begin{aligned} \beta_1 &= \frac{1}{2} \left(1 + \varepsilon + \sqrt{2} \sqrt{\varepsilon^2 + \varepsilon} \right), \\ \beta_2 &= \frac{1}{2} \left(1 + \varepsilon - \sqrt{2} \sqrt{\varepsilon^2 + \varepsilon} \right), \\ \beta_3 &= \frac{1}{2} \left(1 - \varepsilon + \sqrt{2} \sqrt{\varepsilon^2 - \varepsilon} \right), \\ \beta_4 &= \frac{1}{2} \left(1 - \varepsilon - \sqrt{2} \sqrt{\varepsilon^2 - \varepsilon} \right), \end{aligned}$$

where $\varepsilon = \sqrt{1 + 4a^2}$. We see therefore that $p_a(x)$ is irreducible, and that all four roots of the polynomial $p_a(x)$ are real.

The j -invariant of a quartic polynomial $a_0 x^4 + a_1 x^3 + a_2 x^2 + a_3 x + a_4$, is defined to be the expression $j = 2a_2^3 - 9a_1 a_2 a_3 + 27a_1^2 a_4 - 72a_0 a_2 a_4 + 27a_0 a_3^2$, which happens to vanish in the case of $p_a(x)$.

We can now apply a recent result of AKHTARI (see Theorem 1.1 of [1]), which states that if $F(x, y)$ is an irreducible homogeneous quartic polynomial with integer coefficients, whose roots are all real, and for which the j -invariant of the

dehomogenized quartic of $F(x, y)$ vanishes, then the equation $|F(x, y)| = 1$ has at most 12 solutions in integers (x, y) , where the solution $(-x, -y)$ is identified with the solution (x, y) . In particular, the equation

$$x^4 - 2x^3y - 6a^2x^2y^2 + 2a^2xy^3 + a^4y^4 = \pm 1$$

has at most 12 solutions in integers x, y (with $(-x, -y)$ identified with (x, y)), and we note that if a solution (x, y) to this Thue equation gives rise to a positive integer solution to $Y^2 = T_{2k} \pm U_{2k}$, then $(-x, -y)$ gives rise to the same solution. This completes the proof of Theorem 1.1.

References

- [1] S. AKHTARI, The method of Thue and Siegel for binary quartic forms, *International Journal of Number Theory* (to appear).
- [2] M.-H. LE, A note on the number of solutions of the generalized Ramanujan Nagell equation $x^2 - D = k^n$, *Acta Arithmetica* **78** (1996), 11–18.
- [3] W. LJUNGGREN, Zur Theorie der Gleichung $x^2 + 1 = Dy^4$, *Avh. Norsk. Vid. Akad. Oslo* (1942), 1–27.
- [4] W. LJUNGGREN, On the Diophantine equation $Ax^4 - By^2 = C$ ($C = 1, 4$), *Math. Scand.* **21** (1967), 149–158.

BO HE
KEY LABORATORY OF NUMERICAL SIMULATION OF SICHUAN PROVINCE
NEIJANG NORMAL UNIVERSITY
NEIJANG, SICHUAN 641112
P.R. CHINA

E-mail: hebo-one@hotmail.com

ALAIN TOGBÉ
DEPARTMENT OF MATHEMATICS
PURDUE UNIVERSITY NORTH CENTRAL
1401 S. U.S. 421
WESTVILLE, IN 46391
USA

E-mail: atogbe@pnc.edu

P. GARY WALSH
DEPARTMENT OF MATHEMATICS
UNIVERSITY OF OTTAWA
585 KING EDWARD ST.
OTTAWA, ONTARIO, K1N 6N5
CANADA

E-mail: gwalsh@uottawa.ca

(Received May 5, 2008; revised June 25, 2008)