

Title: Annihilators on co-commutators with generalized derivations on Lie ideals

Author(s): Luisa Carini, Vincenzo De Filippis and Basudeb Dhara

Let R be a prime ring of characteristic different from 2, U the Utumi quotient ring of R , $C = Z(U)$ the extended centroid of R , L a non-central Lie ideal of R , H and G non-zero generalized derivations of R . Suppose that there exists $0 \neq a \in R$ such that $a(H(u)u - uG(u)) = 0$, for all $u \in L$, then one of the following holds:

- (1) there exist $b', c' \in U$ such that $H(x) = b'x + xc'$, $G(x) = c'x$ with $ab' = 0$;
- (2) R satisfies s_4 and there exist $b', c', q' \in U$ such that $H(x) = b'x + xc'$, $G(x) = c'x + xq'$, with $a(b' - q') = 0$.

Address:

Luisa Carini
Dipartimento di Matematica
Università di Messina
Messina
Italia

Address:

Vincenzo De Filippis
Dipartimento di Scienze per l'Ingegneria e per l'Architettura
Sezione di Matematica e Eidomatica
Università di Messina
Messina
Italia

Address:

Basudeb Dhara
Department of Mathematics
Belda College, Belda
Paschim Medinipur, 721424 (W.B.)
India