

L_1 -factorization of Pietsch integral operators

By RAFFAELLA CILIA (Catania) and JOAQUIN M. GUTIÉRREZ (Madrid)

Abstract. Given a compact Hausdorff space K and a regular positive finite, Borel measure μ on K , we characterize the operators on $C(K)$ admitting a factorization through the natural inclusion of $C(K)$ into $L_1(K, \mu)$. We also characterize the operators on $L_\infty(\Omega, \nu)$, with ν a positive finite measure, that factor through the natural inclusion of $L_\infty(\Omega, \nu)$ into $L_1(\Omega, \nu)$.

Throughout, E and F will denote Banach spaces, B_{E^*} will stand for the closed unit ball of the dual space E^* endowed with the weak-star topology, and $\mathcal{L}(E, F)$ will be the space of all (linear bounded) operators from E into F endowed with the supremum norm. The symbol k_F will be used for the canonical isometric embedding of F into F^{**} .

An operator $T \in \mathcal{L}(E, F)$ is *Pietsch integral* [DU, Definition VI.3.8] if there exists a countably additive F -valued, Borel measure $\mathcal{G}$ of bounded variation on B_{E^*} such that

$$T(x) = \int_{B_{E^*}} x^*(x) d\mathcal{G}(x^*) \quad (x \in E).$$

The *Pietsch integral norm* of T is given by

$$\|T\|_{\text{PI}} := \inf |\mathcal{G}|(B_{E^*}),$$

where $|\mathcal{G}|$ denotes the variation of $\mathcal{G}$ and the infimum is taken over all vector measures $\mathcal{G}$ satisfying the definition.

Mathematics Subject Classification: Primary: 47B10; Secondary: 46B20.

Key words and phrases: Pietsch integral operator, factorization theorem, $L_1(\mu)$ -spaces, $C(K)$ -spaces, $L_\infty(\mu)$ -spaces.

The first named author was supported in part by GNAMPA (Italy)

Both authors were supported in part by Dirección General de Investigación, MTM 2006–03531 (Spain).

In the theory of Pietsch integral operators and its development, a fundamental rôle is played by the following well-known factorization theorem.

Theorem 1 ([DU, Theorem VI.3.11]). *An operator $T \in \mathcal{L}(E, F)$ is Pietsch integral if and only if there are a compact Hausdorff space K , a regular Borel measure μ on K , and operators $S \in \mathcal{L}(L_1(K, \mu), F)$ and $R \in \mathcal{L}(E, C(K))$ giving rise to the commutative diagram*

$$\begin{array}{ccc} E & \xrightarrow{T} & F \\ R \downarrow & & \uparrow S \\ C(K) & \xrightarrow{J} & L_1(K, \mu) \end{array}$$

where J denotes the natural inclusion of $C(K)$ into $L_1(K, \mu)$.

It is well-known that K may be chosen to be any weak-star compact norming subset of B_{E^*} , denoting by R the natural isometric embedding of E into $C(K)$ [DJT, Theorem 5.6].

VILLANUEVA [V, pages 58–59] noticed that the result remains true if K is any compact space (not necessarily contained in B_{E^*}) such that there is an isomorphic embedding of E into $C(K)$. So we can state:

Theorem 2. *Let $T \in \mathcal{L}(E, F)$ be an operator, and let K be a compact Hausdorff space such that there is an isomorphic embedding $R \in \mathcal{L}(E, C(K))$. Then T is Pietsch integral if and only if there exist a regular Borel measure μ on K and an operator $S \in \mathcal{L}(L_1(K, \mu), F)$ such that the following diagram is commutative*

$$\begin{array}{ccc} E & \xrightarrow{T} & F \\ R \downarrow & & \uparrow S \\ C(K) & \xrightarrow{J} & L_1(K, \mu) \end{array}$$

where J denotes the natural inclusion of $C(K)$ into $L_1(K, \mu)$.

Therefore, we can select the compact space K and the operator $R \in \mathcal{L}(E, C(K))$, provided that it be an isomorphism. Then it seems natural to ask if we can also select the regular Borel measure μ on K .

It is easy to see that the answer is negative. Moreover, we give characterizations of the operators on $C(K)$ spaces that factor through the natural inclusion of $C(K)$ into $L_1(K, \mu)$ when μ is a given regular positive finite, Borel measure

on K . We also characterize the operators on $L_\infty(\Omega, \nu)$, with ν a positive finite measure, that factor through the natural inclusion of $L_\infty(\Omega, \nu)$ into $L_1(\Omega, \nu)$.

The following simple example shows that there exists a functional on $C[0, 1]$ (trivially, Pietsch integral) that cannot be factored through the natural inclusion of $C[0, 1]$ into $L_1([0, 1], \mu)$, where μ is Lebesgue measure.

Example 3. Consider $\delta_{1/2} \in C[0, 1]^*$. Assume that it factors through the natural inclusion $J : C[0, 1] \rightarrow L_1([0, 1], \mu)$. Then $\delta_{1/2} = \xi \circ J$, where $\xi \in L_\infty([0, 1], \mu)$. For each natural number n , let $f_n \in C[0, 1]$ be a function such that

$$f_n(t) = \begin{cases} 0, & \text{if } t \leq \frac{1}{2} - \frac{1}{n} \\ 1, & \text{if } t = \frac{1}{2} \\ 0, & \text{if } t \geq \frac{1}{2} + \frac{1}{n}. \end{cases}$$

Then

$$1 = f_n \left(\frac{1}{2} \right) = |\delta_{1/2}(f_n)| = |\xi \circ J(f_n)| \leq \|\xi\| \|J(f_n)\|_{L_1} \xrightarrow{n \rightarrow \infty} 0,$$

a contradiction.

In fact, this example can be deduced easily from Theorem 5 below.

Given a compact Hausdorff space K , let Σ be the σ -algebra of the Borel subsets of K . Denote by $B(\Sigma)$ the space of all uniform limits of simple measurable functions on K , endowed with the supremum norm.

Let

$$\lambda : B(\Sigma) \longrightarrow C(K)^{**}$$

be the isometric embedding given by

$$\langle \lambda(f), \nu \rangle := \int_K f d\nu \quad (\nu \in C(K)^*, f \in B(\Sigma)),$$

where the integral is defined as in [DU, Definition I.1.12], that is, first on the simple measurable functions and then extended to $B(\Sigma)$.

Denote by

$$i : C(K) \longrightarrow B(\Sigma)$$

the natural embedding [D, Corollary §14.5.1].

Given an operator $T \in \mathcal{L}(C(K), F)$, the representing measure $m_T : \Sigma \rightarrow F^{**}$ associated with T is defined by

$$m_T(A) := T^{**}(\lambda(\chi_A)) \quad (A \in \Sigma)$$

(see the proof of [DU, Theorem VI.2.1]).

If μ is a finite positive measure on K , denote by

$$J : C(K) \longrightarrow L_1(K, \mu)$$

and

$$\overline{J} : B(\Sigma) \longrightarrow L_1(K, \mu)$$

the natural inclusions. Note that $\overline{J} \circ i = J$ (see the diagram below).

$$\begin{array}{ccc} B(\Sigma) & \xrightarrow{\quad \overline{J} \quad} & L_1(K, \mu) \\ & \nwarrow i \quad \nearrow J & \\ & C(K) & \end{array}$$

Indeed, given $\omega \in K$ and $f \in C(K)$, since J , $\overline{J}$, and i are natural inclusions, we have

$$J(f)(\omega) = f(\omega) = i(f)(\omega) = \overline{J}(i(f))(\omega) = (\overline{J} \circ i(f))(\omega),$$

so $\overline{J} \circ i = J$.

The following lemma is well-known (see the proof of [DU, Example VI.3.6]), but we have not found its proof in the literature. We include it for completeness.

Lemma 4. *Let K be a compact Hausdorff space and let μ be a regular finite positive, Borel measure on K . Then the representing measure m_J of $J : C(K) \rightarrow L_1(K, \mu)$ is given by*

$$m_J(A) := \chi_A \quad (A \in \Sigma).$$

PROOF. Define a vector measure $\mathcal{G} : \Sigma \rightarrow L_1(K, \mu)$ by

$$\mathcal{G}(A) := \chi_A \quad (A \in \Sigma).$$

Using the regularity of μ , and choosing $\xi \in L_\infty(K, \mu)$, it is easy to show that $\xi \circ \mathcal{G}$ is a regular set function. It is also easy to see that it is countably additive and has bounded variation.

By [DS, Theorem VI.7.3], there is a (weakly compact) operator

$$T : C(K) \longrightarrow L_1(K, \mu)$$

given by

$$T(f) := \int_K f d\mathcal{G} \quad (f \in C(K)),$$

whose representing measure is $\mathcal{G}$.

By [DU, Theorem I.1.13], there is an operator

$$U : B(\Sigma) \longrightarrow L_1(K, \mu)$$

given by

$$U(f) := \int_K f d\mathcal{G} \quad (f \in B(\Sigma)),$$

with representing measure $\mathcal{G}$.

Using the definitions of U and T , we obtain $U \circ i = T$.

For $A \in \Sigma$, we have

$$U(\chi_A) = \int_K \chi_A d\mathcal{G} = \int_A d\mathcal{G} = \mathcal{G}(A) = \chi_A = \overline{J}(\chi_A).$$

Given $f \in C(K)$, there is a sequence (f_n) of simple measurable functions such that $f_n \rightarrow i(f)$ in $B(\Sigma)$, under the supremum norm [D, Corollary §14.5.1]. Therefore,

$$T(f) = U \circ i(f) = \lim_n U(f_n) = \lim_n \overline{J}(f_n) = \overline{J} \circ i(f) = J(f),$$

so $T = J$ and $m_J = \mathcal{G}$. □

Given an operator $T \in \mathcal{L}(C(K), F)$, the following diagram is commutative:

$$\begin{array}{ccccc} B(\Sigma) & \xrightarrow{\lambda} & C(K)^{**} & & \\ \uparrow i & \nearrow k_{C(K)} & \downarrow T^{**} & & \\ C(K) & \xrightarrow{T} & F & \xrightarrow{k_F} & F^{**} \end{array}$$


Indeed, it is enough to show that $\lambda \circ i = k_{C(K)}$. For $f \in C(K)$ and $\nu \in C(K)^*$, we have

$$\langle \lambda \circ i(f), \nu \rangle = \int_K i(f) d\nu = \int_K f d\nu = \langle \nu, f \rangle = \langle k_{C(K)}(f), \nu \rangle,$$

and the claim is proved.

Theorem 5. *Let K be a compact Hausdorff space, let μ be a regular positive finite, Borel measure on K , and let $T \in \mathcal{L}(C(K), F)$ be an operator. Then the following assertions are equivalent:*

- (a) *There is an operator $S \in \mathcal{L}(L_1(K, \mu), F)$ such that $T = S \circ J$ (see the diagram below).*



- (b) *There exists a constant $C > 0$ such that*

$$\|m_T(A)\| \leq C\mu(A) \quad (A \in \Sigma).$$

- (c) *There exists a constant $C > 0$ such that*

$$|m_T|(A) \leq C\mu(A) \quad (A \in \Sigma).$$

- (d) *There exists a constant $C > 0$ such that*

$$\|T(f)\| \leq C \|J(f)\|_{L_1} \quad (f \in C(K)).$$

Moreover, if (b), (c) or (d) holds, then T is Pietsch integral, and

$$\|T\|_{PI} \leq C\mu(K).$$

PROOF. (a) $\Rightarrow$ (b). For $A \in \Sigma$, we have

$$\begin{aligned}
 \|m_T(A)\| &= \|T^{**}(\lambda(\chi_A))\| = \|S^{**} \circ J^{**}(\lambda(\chi_A))\| \\
 &\leq \|S\| \|m_J(A)\| = \|S\| \|\chi_A\|_{L_1} \quad (\text{by Lemma 4}) \\
 &= \|S\| \mu(A).
 \end{aligned}$$

(b) $\Rightarrow$ (c). By the definition of variation, given $A \in \Sigma$, we have

$$|m_T|(A) = \sup_{\pi} \sum_{N \in \pi} \|m_T(N)\|,$$

where the supremum is taken over all partitions π of A into a finite number of pairwise disjoint members of Σ . Then,

$$\sup_{\pi} \sum_{N \in \pi} \|m_T(N)\| \leq C \sup_{\pi} \sum_{N \in \pi} \mu(N) = C\mu(A).$$

(c) $\Rightarrow$ (d). Let

$$g := \sum_{i=1}^n a_i \chi_{A_i}$$

be a simple measurable function on K , with $(A_i)_{i=1}^n$ a disjoint family of members of Σ . Then,

$$\begin{aligned} \|T^{**}(\lambda(g))\| &= \left\| \sum_{i=1}^n a_i T^{**}(\lambda(\chi_{A_i})) \right\| = \left\| \sum_{i=1}^n a_i m_T(A_i) \right\| \\ &\leq \sum_{i=1}^n |a_i| \|m_T(A_i)\| \leq \sum_{i=1}^n |a_i| |m_T|(A_i) \\ &\leq C \sum_{i=1}^n |a_i| \mu(A_i) = C \int_K \left| \sum_{i=1}^n a_i \chi_{A_i} \right| d\mu = C \|\bar{J}(g)\|_{L_1}. \end{aligned}$$

Given $f \in C(K)$, let (f_n) be a sequence of simple measurable functions on K such that $i(f) = \lim_n f_n$ in $B(\Sigma)$. Then,

$$\begin{aligned} \|T(f)\| &= \|k_F \circ T(f)\| = \|T^{**}(\lambda(i(f)))\| \\ &= \lim_n \|T^{**}(\lambda(f_n))\| \leq C \lim_n \|\bar{J}(f_n)\| \\ &= C \|\bar{J}(i(f))\| = C \|J(f)\|_{L_1}. \end{aligned}$$

(d) $\Rightarrow$ (a). Define an operator $S : J(C(K)) \rightarrow F$ by $S(J(f)) := T(f)$, for all $f \in C(K)$. We have

$$\|S(J(f))\| = \|T(f)\| \leq C \|J(f)\|_{L_1} \quad (f \in C(K)),$$

so S is continuous on $J(C(K))$ endowed with the L_1 -norm. Since $J(C(K))$ is dense in $L_1(K, \mu)$ [DS, Lemma IV.8.19], S has an extension, denoted also by S , to $L_1(K, \mu)$ such that $S \circ J = T$ and $\|S\| \leq C$.

Suppose now that T satisfies (b), (c) or (d), for a constant $C > 0$. Then T satisfies also (a), with $\|S\| \leq C$. Since J is Pietsch integral, so is $T = S \circ J$, and

$$\|T\|_{\text{PI}} \leq \|S\| \|J\|_{\text{PI}} \leq C \mu(K),$$

where we have used [DU, Example VI.3.10]. $\square$

We now study the operators on $L_\infty(\Omega, \nu)$. We first give an example of a functional on $L_\infty([0, 1], \mu)$, where μ is Lebesgue measure, that cannot be factored through the canonical inclusion into $L_1([0, 1], \mu)$.

Example 6. For every $n \in \mathbb{N}$, let

$$g_n := n(n+1)\chi_{[\frac{1}{n+1}, \frac{1}{n}]}$$

Obviously, $g_n \in L_1([0, 1], \mu)$ and $\|g_n\|_{L_1([0, 1], \mu)} = 1$. Define a linear form

$$H : L_\infty([0, 1], \mu) \longrightarrow \mathbb{R}$$

by

$$H(f) := \sum_{n=1}^{+\infty} \frac{1}{n^{3/2}} \int_{[0, 1]} f g_n d\mu \quad \text{for } f \in L_\infty([0, 1], \mu).$$

Since

$$|H(f)| \leq \sum_{n=1}^{+\infty} \frac{1}{n^{3/2}} \|f\|_{L_\infty([0, 1], \mu)},$$

H is continuous. Suppose that there exists $B \in L_1([0, 1], \mu)^*$ such that the following diagram is commutative

$$\begin{array}{ccc} L_\infty([0, 1], \mu) & \xrightarrow{H} & \mathbb{R} \\ & \searrow J \quad \nearrow B & \\ & L_1([0, 1], \mu) & \end{array}$$

where J is the natural inclusion of $L_\infty([0, 1], \mu)$ into $L_1([0, 1], \mu)$. For each $m \in \mathbb{N}$, consider the function

$$f_m := \chi_{[\frac{1}{m+1}, \frac{1}{m}]}$$

We have

$$B \circ J(f_m) = H(f_m) = \sum_{n=1}^{+\infty} \frac{1}{n^{3/2}} \int_{[0, 1]} \chi_{[\frac{1}{m+1}, \frac{1}{m}]} n(n+1) \chi_{[\frac{1}{n+1}, \frac{1}{n}]} d\mu = \frac{1}{m^{3/2}}.$$

Therefore

$$\frac{1}{m^{3/2}} = |B \circ J(f_m)| \leq \|B\| \|f_m\|_{L_1([0, 1], \mu)} = \|B\| \frac{1}{m(m+1)},$$

so

$$\frac{m(m+1)}{m^{3/2}} \leq \|B\| \quad \text{for all } m \in \mathbb{N},$$

a contradiction.

The following result is mentioned in [LM, Exercise 10.10]. We give a proof for completeness.

Lemma 7. *Let (Ω, Σ, μ) be a measure space. Then the space of simple measurable functions is dense in $L_\infty(\Omega, \mu)$.*

PROOF. Let $f \in L_\infty(\Omega, \mu)$. Suppose first that $f \geq 0$. There is a subset $M \subset \Omega$ with $\mu(M) = 0$ such that f is bounded on $\Omega \setminus M$. By [R, Theorem 1.17], there is a nondecreasing sequence (f_n) of simple measurable functions such that

$$\lim_n f_n(\omega) = f(\omega) \quad (\omega \in \Omega \setminus M).$$

Since f is bounded, the convergence is uniform on $\Omega \setminus M$ (see the comment after [R, Theorem 1.17]). Therefore, the L_∞ -norm of $(f_n - f)$ tends to zero.

If f is arbitrary, we decompose $f = f^+ - f^-$ as usual. There are sequences (g_n) and (h_n) of simple measurable functions converging respectively to f^+ and f^- . Then, the sequence of simple measurable functions $(g_n - h_n)$ converges to f in $L_\infty(\Omega, \mu)$. $\square$

The following result is contained in [DJT, Examples 2.9 and Corollary 5.22].

Lemma 8. *Let (Ω, Σ, μ) be a finite measure space. Then the natural inclusion*

$$J : L_\infty(\Omega, \mu) \longrightarrow L_1(\Omega, \mu)$$

is Pietsch integral with norm $\|J\|_{PI} = \mu(\Omega)$.

Given an operator $T \in \mathcal{L}(L_\infty(\Omega, \mu), F)$, its representing measure $m_T : \Sigma \rightarrow F$ is defined by

$$m_T(A) := T(\chi_A) \quad (A \in \Sigma)$$

[DU, page 148].

Theorem 9. *Let (Ω, Σ, μ) be a positive finite measure space and let*

$$T \in \mathcal{L}(L_\infty(\Omega, \mu), F)$$

be an operator. Then the following assertions are equivalent:

- (a) *There is an operator $S \in \mathcal{L}(L_1(\Omega, \mu), F)$ such that $T = S \circ J$ (see the diagram below).*

$$\begin{array}{ccc} L_\infty(\Omega, \mu) & \xrightarrow{T} & F \\ & \searrow J \quad \nearrow S & \\ & L_1(\Omega, \mu) & \end{array}$$

(b) *There exists a constant $C > 0$ such that*

$$\|m_T(A)\| \leq C\mu(A) \quad (A \in \Sigma).$$

(c) *There exists a constant $C > 0$ such that*

$$|m_T|(A) \leq C\mu(A) \quad (A \in \Sigma).$$

(d) *There exists a constant $C > 0$ such that*

$$\|T(f)\| \leq C \|J(f)\|_{L_1} \quad (f \in L_\infty(\Omega, \mu)).$$

Moreover, if (b), (c) or (d) holds, then T is Pietsch integral, and

$$\|T\|_{PI} \leq C\mu(\Omega).$$

PROOF. (a) $\Rightarrow$ (b). For $A \in \Sigma$, we have

$$\|m_T(A)\| = \|T(\chi_A)\| = \|S \circ J(\chi_A)\| \leq \|S\| \|\chi_A\|_{L_1} = \|S\| \mu(A).$$

(b) $\Rightarrow$ (c) as in Theorem 5.

(c) $\Rightarrow$ (d). Let

$$g := \sum_{i=1}^n a_i \chi_{A_i}$$

be a simple measurable function on Ω , with $(A_i)_{i=1}^n$ a disjoint family of members of Σ . Then,

$$\begin{aligned} \|T(g)\| &= \left\| \sum_{i=1}^n a_i T(\chi_{A_i}) \right\| = \left\| \sum_{i=1}^n a_i m_T(A_i) \right\| \\ &\leq C \int_{\Omega} \left| \sum_{i=1}^n a_i \chi_{A_i} \right| d\mu \quad (\text{as in the proof of Theorem 5}) \\ &= C \|J(g)\|_{L_1}. \end{aligned}$$

Let $f \in L_\infty(\Omega, \mu)$. By Lemma 7, there is a sequence (f_n) of simple measurable functions on Ω such that $\lim_n f_n = f$ in $L_\infty(\Omega, \mu)$. Then,

$$\|T(f)\| = \lim_n \|T(f_n)\| \leq C \lim_n \|J(f_n)\|_{L_1} = C \|J(f)\|_{L_1}.$$

(d) $\Rightarrow$ (a) as in the proof of Theorem 5, bearing in mind that $J(L_\infty(\Omega, \mu))$ is dense in $L_1(\Omega, \mu)$ [R, Theorem 3.13].

Suppose now that T satisfies (b), (c) or (d), for a constant $C > 0$. Then T satisfies also (a), with $\|S\| \leq C$. Since J is Pietsch integral, so is $T = S \circ J$, and

$$\|T\|_{PI} \leq \|S\| \|J\|_{PI} \leq C\mu(\Omega),$$

where we have used Lemma 8. $\square$

References

- [DJT] J. DIESTEL, H. JARCHOW and A. TONGE, Absolutely Summing Operators, Vol. 43, Cambridge Stud. Adv. Math., *Cambridge University Press, Cambridge*, 1995.
- [DU] J. DIESTEL and J. J. UHL, JR., Vector Measures, Vol. 15, Math. Surveys Monographs, *American Mathematical Society, Providence RI*, 1977.
- [D] V. DINULEANU, Vector Measures, *Pergamon Press, Oxford*, 1967.
- [DS] N. DUNFORD and J. T. SCHWARTZ, Linear Operators, I: General Theory, *Wiley, New York*, 1988.
- [LM] J. LUKEŠ and J. MALÝ, Measure and Integral, *Matfyzpress, Prague*, 1995.
- [R] W. RUDIN, Real and Complex Analysis, 3rd ed., *McGraw-Hill, New York*, 1987.
- [V] I. VILLANUEVA, Integral mappings between Banach spaces, *J. Math. Anal. Appl.* **279** (2003), 56–70.

RAFFAELLA CILIA
DIPARTIMENTO DI MATEMATICA
FACOLTÀ DI SCIENZE
UNIVERSITÀ DI CATANIA
VIALE ANDREA DORIA 6
95100 CATANIA
ITALY

E-mail: cilia@dmf.unict.it

JOAQUÍN M. GUTIÉRREZ
DEPARTAMENTO DE MATEMÁTICA APLICADA
ETS DE INGENIEROS INDUSTRIALES
UNIVERSIDAD POLITÉCNICA DE MADRID
C. JOSÉ GUTIÉRREZ ABASCAL 2
28006 MADRID
SPAIN

E-mail: jgutierrez@etsii.upm.es

(Received October 14, 2008)